


PATENT
4001.002282

U.S. PATENT APPLICATION
for
CANCER TREATMENT KITS USING
ANTIBODIES TO AMINOPHOSPHOLIPIDS
by
Philip E. Thorpe
and
Sophia Ran

6621.00 2481.5660

| EXPRESS MAIL MAILING LABEL | |
|--|-------------------|
| NUMBER | EL 332 810 590 US |
| DATE OF DEPOSIT | July 12, 1999 |
| I hereby certify that this paper or fee is being deposited with the United States Postal Service "EXPRESS MAIL POST OFFICE TO ADDRESSEE" service on the date indicated above and is addressed to: Assistant Commissioner for Patents, Washington D.C. 20231. | |
|  | |
| Anita Cheyne | |

BACKGROUND OF THE INVENTION

The present application claims priority to first provisional application Serial No. 60/092,672, filed July 13, 1998, and second provisional application Serial No. 60/110,608, filed December 02, 1998, the entire text and figures of which applications are incorporated herein by reference without disclaimer. The U.S. Government owns rights in the present invention pursuant to grant numbers 1RO1CA74951-01 and 5RO1CA54168-05 from the National Institutes of Health.

1. Field of the Invention

The present invention relates generally to the fields of blood vessels and tumor biology. More particularly, it embodies the surprising findings that aminophospholipids, such as phosphatidylserine and phosphatidylethanolamine, are specific and stable markers of tumor blood vessels and that administration of anti-aminophospholipid antibodies alone is sufficient to induce thrombosis and tumor regression. The invention thus provides safe and effective methods and compositions for the specific targeting and destruction of tumor blood vessels and for the treatment of solid tumors. The use of unconjugated anti-phosphatidylserine antibodies is a particular advantage, although the invention provides various effective compositions and combinations thereof.

2. Description of the Related Art

Tumor cell resistance to chemotherapeutic agents represents a significant problem in clinical oncology. In fact, this is one of the main reasons why many of the most prevalent forms of human cancer still resist effective chemotherapeutic intervention, despite certain advances in the field of chemotherapy.

A significant problem to address in tumor treatment regimens is the desire for a "total cell kill". This means that the more effective treatment regimens come closer to a total cell kill of all so-called "clonogenic" malignant cells, *i.e.*, cells that have the ability to grow uncontrolled and replace any tumor mass that might be removed by the therapy. Due to the goal of

Another effective version of the vascular targeting approach is to target a coagulation factor to tumor vasculature (Huang *et al.*, 1997; U.S. Patents 5,877,289, ^{6,004,555 and 6,093,329} ~~5, _____ and 5, _____~~ (U.S. Applications Serial Nos. 08/487,427 and 08/482,369; Issue Fees paid)). The use of antibodies and other targeting agents to deliver coagulants to tumor vasculature has the further advantages of reduced immunogenicity and even lower risk of toxic side effects. As disclosed in U.S. Patent 5,877,289, a preferred coagulation factor for use in such tumor-specific thrombogens, or "coaguligands", is a truncated version of the human coagulation-inducing protein, Tissue Factor (TF). TF is the major initiator of blood coagulation (Ruf *et al.*, 1991; Edgington *et al.*, 1991; Ruf and Edgington, 1994). Treatment of tumor-bearing mice with such coaguligands results in significant tumor necrosis and even complete tumor regression in many animals (Huang *et al.*, 1997; U.S. Patents 5,877,289, ^{6,004,555 and 6,093,329} ~~5, _____ and 5, _____~~ (U.S. Applications Serial Nos. 08/487,427 and 08/482,369; Issue Fees paid)).

Although the specific delivery of therapeutic agents, such as anti-cellular agents, toxins and coagulation factors, to tumor vessels represents a significant advance in tumor treatment protocols, there is still room for additional or even alternative vascular targeting therapies. The identification of additional targets to allow specific tumor vessel destruction *in vivo* would naturally be of benefit in expanding the number of targeting options. More particularly, as the previously described vascular targeting constructs and coaguligands are two-component systems, involving the targeting agent and the effector portion, the development of a one component agent for tumor vasculature destruction would represent a major advance. Should the preparation of this type of agent prove possible, this would also likely speed the progress of anti-vascular therapy to the clinic, given the simplicity of the new therapeutic agent.

SUMMARY OF THE INVENTION

The present invention addresses the needs of the prior art by providing new, simplified therapeutic methods for specific tumor destruction. The invention is based, in part, on the finding that aminophospholipids, such as phosphatidylserine and phosphatidylethanolamine, are accessible and stably targetable markers of tumor vasculature. More particularly, the invention

embodies the unexpected discovery that naked antibodies against aminophospholipid components are capable of specifically inducing tumor blood vessel destruction and tumor necrosis *in vivo*.

5 Certain preferred aspects of the invention were developed from the surprising discovery that antibodies against the aminophospholipid, phosphatidylserine (PS), specifically localize to the vasculature of solid tumors and, even more surprisingly, exert a tumor destructive effect in the absence of conjugation to effector molecules, such as toxins or coagulants. Single component therapeutics directed against aminophospholipids thus represent a breakthrough in vascular
10 targeting and provide safe and effective methods for the treatment of solid tumors.

 An underlying surprising feature of the invention is that translocation of aminophospholipids, such as PS, to the surface of tumor vascular endothelial cells occurs, at least in a significant part, independently of cell damage and apoptotic or other cell-death mechanisms.
15 PS surface expression in the tumor environment is therefore not a consequence of, or a trigger for, cell death and destruction, but occurs on morphologically intact vascular endothelial cells. This means that PS expression is not transient, but rather is stable enough to provide a target for therapeutic intervention.

 The methods of the invention thus provide for killing, or specifically killing, tumor vascular endothelial cells, and comprise administering to an animal or patient having a vascularized tumor at least one dose of a biologically effective amount of at least a first pharmaceutical composition comprising a naked or unconjugated antibody, or antigen-binding region thereof, that binds to at least a first aminophospholipid expressed on the luminal surface
20 of tumor vascular endothelial cells. The "biologically effective amount" is an amount of the naked or unconjugated antibody effective to specifically kill at least a portion, and preferably a significant portion, of the tumor vascular endothelial cells, as opposed to endothelial cells in normal vessels, upon binding to an aminophospholipid expressed on the luminal surface of the tumor vascular endothelial cells. As such, it is an "endothelial cell killing amount" or a "tumor
25

vascular endothelial cell killing amount" of a naked or unconjugated anti-aminophospholipid antibody or antigen-binding region thereof.

As used throughout the entire application, the terms "a" and "an" are used in the sense that they mean "at least one", "at least a first", "one or more" or "a plurality" of the referenced components, except in instances wherein an upper limit is thereafter specifically stated. Therefore "an anti-aminophospholipid antibody" means "at least a first anti-aminophospholipid antibody". The operable limits and parameters of combinations, as with the amounts of any single agent, will be known to those of ordinary skill in the art in light of the present disclosure.

10

The "a" and "an" terms are also used to mean "at least one", "at least a first", "one or more" or "a plurality" of steps in the recited methods, except where specifically stated. This is particularly relevant to the administration steps in the treatment methods. Thus, not only may different doses be employed with the present invention, but different numbers of doses, *e.g.*, injections, may be used, up to and including multiple injections.

15

An "aminophospholipid", as used herein, means a phospholipid that includes within its structure at least a first primary amino group. Preferably, the term "aminophospholipid" is used to refer to a primary amino group-containing phospholipid that occurs naturally in mammalian cell membranes. However, this is not a limitation on the meaning of the term "aminophospholipid", as this term also extends to non-naturally occurring or synthetic aminophospholipids that nonetheless have uses in the invention, *e.g.*, as an immunogen in the generation of anti-aminophospholipid antibodies ("cross-reactive antibodies") that do bind to aminophospholipids of mammalian plasma membranes. The aminophospholipids of U.S. Patent No. 5,767,298, incorporated herein by reference, are appropriate examples.

20

25

The prominent aminophospholipids found in mammalian biological systems are the negatively-charged phosphatidylserine ("PS") and the neutral or zwitterionic phosphatidylethanolamine ("PE"), which are therefore preferred aminophospholipids for targeting by the present invention. However, the invention is by no means limited to the

30

targeting of phosphatidylserines and phosphatidylethanolamines, and any other aminophospholipid target may be employed (White *et al.*, 1978; incorporated herein by reference) so long as it is expressed, accessible or complexed on the luminal surface of tumor vascular endothelial cells.

5

All aminophospholipid-, phosphatidylserine- and phosphatidylethanolamine-based components are encompassed as targets of the invention irrespective of the type of fatty acid chains involved, including those with short, intermediate or long chain fatty acids, and those with saturated, unsaturated and polyunsaturated fatty acids. Preferred compositions for raising antibodies for use in the present invention may be aminophospholipids with fatty acids of C18, with C18:1 being more preferred (Levy *et al.*, 1990; incorporated herein by reference). To the extent that they are accessible on tumor vascular endothelial cells, aminophospholipid degradation products having only one fatty acid (lyso derivatives), rather than two, may also be targeted (Oamar *et al.*, 1990; incorporated herein by reference).

15

Another group of potential aminophospholipid targets include, for example, phosphatidal derivatives (plasmalogens), such as phosphatidylserine and phosphatidylethanolamine (having an ether linkage giving an alkenyl group, rather than an ester linkage giving an acyl group). Indeed, the targets for therapeutic intervention by the present invention include any substantially lipid-based component that comprises a nitrogenous base and that is present, expressed, translocated, presented or otherwise complexed in a targetable form on the luminal surface of tumor vascular endothelial cells, not excluding phosphatidylcholine ("PC"). Lipids not containing glycerol may also form appropriate targets, such as the sphingolipids based upon sphingosine and derivatives.

25

The biological basis for including a range of lipids in the group of targetable components lies, in part, with the observed biological phenomena of lipids and proteins combining in membranous environments to form unique lipid-protein complexes. Such lipid-protein complexes extend to antigenic and immunogenic forms of lipids such as phosphatidylserine, phosphatidylethanolamine and phosphatidylcholine with, *e.g.*, proteins such as β_2 -glycoprotein I, prothrombin, kininogens and prekallikrein. Therefore, as proteins and polypeptides can have

30

one or more free primary amino groups, it is contemplated that a range of effective "aminophospholipid targets" may be formed *in vivo* from lipid components that are not aminophospholipids in the strictest sense. Nonetheless, all such targetable complexes that comprise lipids and primary amino groups constitute an "aminophospholipid" within the scope of the present invention.

The inventive methods also act to arrest blood flow, or specifically arrest blood flow, in tumor vasculature. This is achieved by administering to an animal or patient having a vascularized tumor at least one dose of at least a first pharmaceutical composition comprising a coagulation-inducing amount, or a vessel-occluding amount, of at least a first naked or unconjugated antibody, or antigen-binding region thereof, that binds to an aminophospholipid, preferably phosphatidylserine or phosphatidylethanolamine, translocated to the luminal surface of tumor vasculature.

A "coagulation-inducing amount" or "vessel-occluding amount" is an amount of the naked or unconjugated antibody effective to specifically induce or promote coagulation in, and hence occlude, at least a portion, and preferably a significant portion, of tumor or intratumoral blood vessels, as opposed to normal blood vessels, upon binding to an aminophospholipid, preferably phosphatidylserine or phosphatidylethanolamine, translocated to the luminal surface of tumor blood vessels. The "vessel-occluding amount" is therefore a functionally effective amount, and is not a physical mass of antibody sufficient to span the breadth of a vessel.

Methods for destroying, or specifically destroying, tumor vasculature are provided that comprise administering to an animal or patient having a vascularized tumor one or more doses of at least a first pharmaceutical composition comprising a tumor-destructive amount of at least a first naked or unconjugated antibody, or antigen-binding region thereof, that binds to an aminophospholipid, preferably phosphatidylserine or phosphatidylethanolamine, presented on the luminal surface of tumor vasculature. The "tumor-destructive amount" is an amount of the naked or unconjugated antibody effective to specifically destroy or occlude at least a portion, and preferably a significant portion, of tumor blood vessels, as opposed to normal blood vessels,

upon binding to an aminophospholipid, preferably phosphatidylserine or phosphatidylethanolamine, presented on the luminal surface of the vascular endothelial cells of the tumor blood vessels.

5 The invention further encompasses methods for treating cancer and solid tumors, comprising administering to an animal or patient having a vascularized tumor a tumor necrosis-inducing amount or amounts of at least a first pharmaceutical composition comprising at least a first naked or unconjugated antibody, or antigen-binding fragment thereof, that binds to an aminophospholipid, preferably phosphatidylserine or phosphatidylethanolamine, on the luminal
10 surface of blood vessels of the vascularized tumor. The "tumor necrosis-inducing amount" is an amount of the naked or unconjugated antibody effective to specifically induce hemorrhagic necrosis in at least a portion, and preferably a significant portion, of the tumor upon binding to an aminophospholipid, preferably phosphatidylserine or phosphatidylethanolamine, complexed at the luminal surface of the vascular endothelial cells of the tumor blood vessels, while exerting
15 little adverse side effects on normal, healthy tissues.

20 The methods of the invention may thus be summarized as methods for treating an animal or patient having a vascularized tumor, comprising administering to the animal or patient at least a first dose of a therapeutically effective amount of at least a first pharmaceutical composition comprising at least a first naked or unconjugated antibody, or antigen-binding fragment thereof, that binds to an aminophospholipid (preferably phosphatidylserine or phosphatidylethanolamine) present, expressed, translocated, presented or complexed at the luminal surface of blood transporting vessels of the vascularized tumor.

25 The essence of the invention may also be defined as a composition comprising at least a first naked or unconjugated anti-aminophospholipid antibody, preferably an anti-phosphatidylserine or anti-phosphatidylethanolamine antibody, or antigen-binding fragment thereof, for use in the preparation of a medicament for use in tumor vasculature destruction and for human tumor treatment. This can also be defined as a composition comprising at least a first
30 naked or unconjugated anti-aminophospholipid antibody, preferably an anti-PS or anti-PE

antibody, or antigen-binding fragment thereof, for use in the preparation of a medicament for use in binding to an aminophospholipid, preferably phosphatidylserine or phosphatidylethanolamine, present, expressed, translocated, presented or complexed at the luminal surface of blood transporting vessels of a vascularized tumor and for use in inducing tumor vasculature destruction and for human tumor treatment.

In the methods, medicaments and uses of the present invention, one of the advantages lies in the fact that the provision of a simple naked or unconjugated anti-aminophospholipid antibody composition, preferably anti-phosphatidylserine or anti-phosphatidylethanolamine, into the systemic circulation of an animal or patient results in the preferential or specific destruction of the tumor vasculature and the induction of tumor necrosis. The invention therefore solves the problem of the complex preparative methods of the multicomponent anti-vascular agents of the prior art.

The terms "naked" and "unconjugated" antibody, as used herein, are intended to refer to an antibody that is not conjugated, operatively linked or otherwise physically or functionally associated with an effector moiety, such as a cytotoxic or coagulative agent. It will be understood that the terms "naked" and "unconjugated" antibody do not exclude antibody constructs that have been stabilized, multimerized, humanized or in any other way manipulated, other than by the attachment of an effector moiety.

Accordingly, all post-translationally modified naked and unconjugated antibodies are included herewith, including where the modifications are made in the natural antibody-producing cell environment, by a recombinant antibody-producing cell, and are introduced by the hand of man after initial antibody preparation. Of course, the term "naked" antibody does not exclude the ability of the antibody to form functional associations with effector cells and/or molecules after administration to the body, as some such interactions are necessary in order to exert a biological effect. The lack of associated effector group is therefore applied in definition to the naked antibody *in vitro*, not *in vivo*.

In the context of the present invention, the term "a vascularized tumor" most preferably means a vascularized, malignant tumor, solid tumor or "cancer". The invention is particularly advantageous in treating vascularized tumors of at least about intermediate size, and in treating large vascularized tumors - - although this is by no means a limitation on the invention. The invention may therefore be used in the treatment of any tumor that exhibits aminophospholipid-positive blood vessels, preferably phosphatidylserine- and/or phosphatidylethanolamine-positive blood vessels.

In preferred embodiments, the tumors to be treated by the present invention will exhibit a killing effective number of aminophospholipid-positive blood vessels. "A killing effective number of aminophospholipid-positive blood vessels", as used herein, means that at least about 3% of the total number of blood vessels within the tumor will be positive for aminophospholipid expression, preferably phosphatidylserine and/or phosphatidylethanolamine expression. Preferably, at least about 5%, at least about 8%, or at least about 10% or so, of the total number of blood vessels within the tumor will be positive for aminophospholipid expression. Given the aminophospholipid-negative, particularly PS-negative, nature of the blood vessels within normal tissues, the tumor vessels will act as sink for the administered antibodies. Furthermore, as destruction of only a minimum number of tumor vessels can cause widespread thrombosis, necrosis and an avalanche of tumor cell death, antibody localization to all, or even a majority, of the tumor vessels is not necessary for effective therapeutic intervention.

Nonetheless, in more preferred embodiments, tumors to be treated by this invention will exhibit a significant number of aminophospholipid-positive blood vessels. "A significant number of aminophospholipid-positive blood vessels", as used herein, means that at least about 10-12% of the total number of blood vessels within the tumor will be positive for aminophospholipid expression, preferably phosphatidylserine and/or phosphatidylethanolamine expression. Even more preferably, the % of aminophospholipid-expressing tumor vessels will be at least about 15%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, or at least about 80% or so of the total number of blood vessels within the tumor, up to and including even at least about 90% or 95% of the vessels.

5 The "therapeutically effective amounts" for use in the invention are amounts of naked or unconjugated anti-aminophospholipid antibodies, preferably anti-PS or anti-PE antibodies, effective to specifically kill at least a portion of tumor vascular endothelial cells; to specifically promote coagulation in at least a portion of tumor blood vessels; to specifically occlude or destroy at least a portion of blood transporting vessels of the tumor; to specifically induce necrosis in at least a portion of a tumor; and/or to induce tumor regression or remission upon administration to selected animals or patients. Such effects are achieved while exhibiting little or no binding to, or little or no killing of, vascular endothelial cells in normal, healthy tissues; little or no coagulation in, occlusion or destruction of blood vessels in healthy, normal tissues; and exerting negligible or manageable adverse side effects on normal, healthy tissues of the animal or patient.

15 The terms "preferentially" and "specifically", as used herein in the context of promoting coagulation in, or destroying, tumor vasculature, and/or in the context of causing tumor necrosis, thus mean that anti-aminophospholipid antibodies function to achieve coagulation, destruction and/or tumor necrosis that is substantially confined to the tumor vasculature and tumor site, and does not substantially extend to causing coagulation, destruction and/or tissue necrosis in normal, healthy tissues of the animal or subject. The structure and function of healthy cells and tissues is therefore maintained substantially unimpaired by the practice of the invention.

20 Therapeutic benefits may be realized by the administration of at least two, three or more naked or unconjugated anti-aminophospholipid antibodies; bispecific antibodies; chimeric antibodies; and/or dimeric, trimeric or multimeric antibodies. The anti-aminophospholipid antibodies may also be combined with other therapies to provide combined therapeutically effective amounts, as disclosed herein.

Although understanding the mechanism of action is not necessary to the practice of the present anti-aminophospholipid antibody treatment invention, the methods may operate to induce cell-mediated cytotoxicity, complement-mediated lysis and/or apoptosis. The cytotoxic methods

may also be based upon antibody-induced cell signaling (direct signaling), or mimicking or altering signal transduction pathways (indirect signaling). The ability of the anti-aminophospholipid antibodies to localize and bind to a component of the membrane itself may also be relevant, as opposed to previous therapies that are generally directed to binding to a protein component or complex, which may be sterically distinct or distant from the membrane surface itself.

The treatment methods thus include administering to an animal or patient having a vascularized tumor at least a first pharmaceutical composition comprising an amount of at least a first antibody construct effective to induce, or specifically induce, cell-mediated cytotoxicity of at least a portion of the tumor vascular endothelial cells. Herein, the first antibody construct is a naked or unconjugated antibody, or effective fragment thereof, that binds to an aminophospholipid, preferably phosphatidylserine or phosphatidylethanolamine, present, expressed, translocated, presented or complexed at the luminal surface of tumor vascular endothelial cells and that induces cell-mediated cytotoxicity of at least a portion of the tumor vascular endothelial cells, as opposed to endothelial cells in normal vessels. As used herein, "cell-mediated cytotoxicity or destruction" includes ADCC (antibody-dependent, cell-mediated cytotoxicity) and NK (natural killer) cell killing.

The methods further include administering to an animal or patient having a vascularized tumor at least a first pharmaceutical composition comprising an amount of at least a first antibody construct effective to induce, or specifically induce, complement-mediated lysis of at least a portion of the tumor vascular endothelial cells. Herein, the first antibody construct is a naked or unconjugated antibody, or effective fragment thereof, that binds to an aminophospholipid, preferably phosphatidylserine or phosphatidylethanolamine, present, expressed, translocated, presented or complexed at the luminal surface of tumor vascular endothelial cells and that induces complement-mediated lysis of at least a portion of the tumor vascular endothelial cells, as opposed to endothelial cells in normal vessels. As used herein, "complement-mediated or complement-dependent lysis or cytotoxicity" means the process by which the complement-dependent coagulation cascade is activated, multi-component complexes

are assembled, ultimately generating a lytic complex that has direct lytic action, causing cell permeabilization. Anti-aminophospholipid antibodies for use in inducing complement-mediated lysis will generally include the Fc portion of the antibody.

5 The complement-based mechanisms by which the present invention may operate further include "complement-activated ADCC". In such aspects, the administered naked or unconjugated antibodies, or fragments thereof, bind to an aminophospholipid, preferably phosphatidylserine or phosphatidylethanolamine, present, expressed, translocated, presented or complexed at the luminal surface of tumor vascular endothelial cells and induce complement-
10 activated ADCC of at least a portion of the tumor vascular endothelial cells, as opposed to endothelial cells in normal vessels. "Complement-activated ADCC" is used to refer to the process by which complement, not an antibody Fc portion *per se*, holds a multi-component complex together and in which cells such as neutrophils lyse the target cell.

15 In other embodiments, the methods include administering to an animal or patient having a vascularized tumor at least a first pharmaceutical composition comprising an amount of at least a first antibody construct effective to induce, or specifically induce, apoptosis in at least a portion of the tumor vascular endothelial cells. Herein, the first antibody construct is a naked or unconjugated antibody, or antigen-binding fragment thereof, that binds to an aminophospholipid, preferably phosphatidylserine or phosphatidylethanolamine, present, expressed, translocated, presented or complexed at the luminal surface of tumor vascular endothelial cells and that induces apoptosis in at least a portion of the tumor vascular endothelial cells, as opposed to endothelial cells in normal vessels. As used herein, "induces apoptosis" means induces the process of programmed cell death that, during the initial stages, maintains the integrity of the cell
20 membrane, yet transmits the death-inducing signals into the cell. This is opposed to the mechanisms of cell necrosis, during which the cell membrane loses its integrity and becomes leaky at the onset of the process.

25 The anti-aminophospholipid antibody treatment methods will generally involve the administration of at least one dose of the pharmaceutically effective composition to the animal

systemically, such as via intravenous injection. However, any route of administration that allows the antibody localize to the tumor vascular endothelial cells and to induce cell-mediated cytotoxicity, complement-mediated lysis and/or apoptosis thereof will be acceptable.

5 "Administration", as used herein, therefore means provision or delivery of anti-aminophospholipid antibodies in an amount(s) and for a period of time(s) effective to allow binding to an aminophospholipid, preferably phosphatidylserine or phosphatidylethanolamine, present, expressed, translocated, presented or complexed at the luminal surface of blood transporting vessels of the vascularized tumor, and to exert a tumor vasculature destructive and
10 tumor-regressive effect. The passive administration of proteinaceous antibodies is generally preferred, in part, for its simplicity and reproducibility.

However, the term "administration" is herein used to refer to any and all means by which anti-aminophospholipid antibodies are delivered or otherwise provided to the tumor vasculature.

15 "Administration" therefore includes the provision of cells, such as hybridomas, that produce the anti-aminophospholipid antibodies in a manner effective to result in the delivery of the anti-aminophospholipid antibodies to the tumor vasculature, and/or their localization to such vasculature. In such embodiments, it may be desirable to formulate or package the antibody-producing cells in a selectively permeable membrane, structure or implantable device, generally
20 one that can be removed to cease therapy.

"Antibody administration", as used herein, also extends to all methods by which anti-aminophospholipid antibodies are generated in a patient ("endogenous antibodies"), allowing them to circulate and localize to the tumor vasculature. Therefore, "administration" also includes
25 the active immunization of a patient with an immunogenically effective amount of an aminophospholipid sample, antigen or immunogen. All methods of human immunization are appropriate for use in such embodiments, as exemplified by those described below in the context of generating an anti-aminophospholipid antibody response in an animal in order to obtain the antibody therefrom. Exogenous antibody administration will still generally be preferred over cell-

and immunization-based delivery, as this represents a less invasive method that allows the dose to be closely monitored and controlled.

5 The "antibody administration methods" of the invention also extend to the provision of nucleic acids that encode anti-aminophospholipid antibodies in a manner effective to result in the expression of the anti-aminophospholipid antibodies in the vicinity of the tumor vasculature, and/or in the expression of anti-aminophospholipid antibodies that can localize to the tumor vasculature. Any gene therapy technique may be employed, such as naked DNA delivery, recombinant genes and vectors, cell-based delivery, including *ex vivo* manipulation of patients' 10 cells, and the like.

One of the benefits of the present invention is that aminophospholipids, particularly phosphatidylserine and phosphatidylethanolamine, are generally expressed or available throughout the tumor vessels. Aminophospholipid expression on established, intratumoral blood 15 vessels is advantageous as targeting and destroying such vessels will rapidly lead to anti-tumor effects. However, so long as the administered anti-aminophospholipid antibodies bind to at least a portion of the blood transporting vessels, significant anti-tumor effects will ensue. This will not be problematical as aminophospholipids, such as phosphatidylserine and phosphatidylethanolamine, are expressed on the large, central vessels, and also on veins, 20 venules, arteries, arterioles and blood transporting capillaries in all regions of the tumor.

In any event, the ability of the anti-aminophospholipid antibodies to destroy the tumor vasculature means that tumor regression can be achieved, rather than only tumor stasis. Tumor stasis is most often the result of anti-angiogenic therapies that target only the budding vessels at 25 the periphery of a solid tumor and stop the vessels proliferating. Even if the present invention targeted more of the peripheral regions of the tumor in certain tumor types, which is not currently believed to be the case, destruction of the blood transporting vessels in such areas would still lead to widespread thrombosis and tumor necrosis.

In any of the foregoing methods, the terms anti-aminophospholipid "antibody, naked antibody and unconjugated antibody", as used herein, refer broadly to any immunologic binding agent, such as polyclonal and monoclonal IgG, IgM, IgA, IgD and IgE antibodies. Generally, IgG and/or IgM are preferred because they are the most common antibodies in the physiological situation and because they are most easily made in a laboratory setting.

Polyclonal anti-aminophospholipid antibodies, obtained from antisera, may be employed in the invention. However, the use of monoclonal anti-aminophospholipid antibodies (MAbs) will generally be preferred. MAbs are recognized to have certain advantages, *e.g.*, reproducibility and large-scale production, that makes them suitable for clinical treatment. The invention thus provides monoclonal antibodies of the murine, human, monkey, rat, hamster, rabbit and even frog or chicken origin. Due to the ease of preparation and ready availability of reagents, murine monoclonal antibodies will be used in certain embodiments.

As will be understood by those in the art, the immunologic binding reagents encompassed by the term "anti-aminophospholipid antibody" extend to all naked and unconjugated anti-aminophospholipid antibodies from all species, and antigen binding fragments thereof, including dimeric, trimeric and multimeric antibodies; bispecific antibodies; chimeric antibodies; human and humanized antibodies; recombinant and engineered antibodies, and fragments thereof.

The term "anti-aminophospholipid antibody" is thus used to refer to any anti-aminophospholipid antibody-like molecule that has an antigen binding region, and includes antibody fragments such as Fab', Fab, F(ab')₂, single domain antibodies (DABs), Fv, scFv (single chain Fv), and the like. The techniques for preparing and using various antibody-based constructs and fragments are well known in the art.

In certain embodiments, the antibodies employed will be "humanized" or human antibodies. "Humanized" antibodies are generally chimeric monoclonal antibodies from mouse, rat, or other non-human species, bearing human constant and/or variable region domains ("part-human chimeric antibodies"). Mostly, humanized monoclonal antibodies for use in the present

invention will be chimeric antibodies wherein at least a first antigen binding region, or complementarity determining region (CDR), of a mouse, rat or other non-human anti-aminophospholipid monoclonal antibody is operatively attached to, or "grafted" onto, a human antibody constant region or "framework".

5

"Humanized" monoclonal antibodies for use herein may also be anti-aminophospholipid monoclonal antibodies from non-human species wherein one or more selected amino acids have been exchanged for amino acids more commonly observed in human antibodies. This can be readily achieved through the use of routine recombinant technology, particularly site-specific mutagenesis.

10

Entirely human, rather than "humanized", anti-aminophospholipid antibodies may also be prepared and used in the present invention. Such human antibodies may be polyclonal antibodies, as obtained from human patients that have any one or more of a variety of diseases, disorders or clinical conditions associated with the production of anti-aminophospholipid antibodies. Such antibodies may be concentrated, partially purified or substantially purified for use herein.

15

A range of techniques are also available for preparing human monoclonal antibodies. As human patients with anti-aminophospholipid antibody-producing diseases exist, the anti-aminophospholipid antibody-producing cells from such patients may be obtained and manipulated *in vitro* to provide a human monoclonal antibody. The *in vitro* manipulations or techniques include fusing to prepare a monoclonal antibody-producing hybridoma, and/or cloning the gene(s) encoding the anti-aminophospholipid antibody from the cells ("recombinant human antibodies").

20

25

Human anti-aminophospholipid antibody-producing cells may also be obtained from human subjects without an anti-aminophospholipid antibody-associated disease, *i.e.* "healthy subjects" in the context of the present invention. To achieve this, one would simply obtain a population of mixed peripheral blood lymphocytes from a human subject, including antigen-presenting and antibody-producing cells, and stimulate the cell population *in vitro* by admixing

30

with an immunogenically effective amount of an aminophospholipid sample. Again, the human anti-aminophospholipid antibody-producing cells, once obtained, could be used in hybridoma and/or recombinant antibody production.

5 Further techniques for human monoclonal antibody production include immunizing a transgenic animal, preferably a transgenic mouse, that comprises a human antibody library with an immunogenically effective amount of an aminophospholipid sample. This also generates human anti-aminophospholipid antibody-producing cells for further manipulation in hybridoma and/or recombinant antibody production, with the advantage that spleen cells, rather than peripheral blood
10 cells, can be readily obtained from the transgenic animal or mouse.

As used herein, the term "anti-aminophospholipid antibody" is used co-extensively with "naked and unconjugated" to mean anti-aminophospholipid antibodies, and antigen binding fragments thereof, that are not conjugated to, or operatively associated with, an effector
15 molecule, such as a cytotoxic agent or coagulant. In addition to non-effector modifications of the antibody, and *in vivo* interactions, the term "naked" in no way excludes combinations of the antibody with other therapeutic agents, as disclosed in detail herein.

20 In contrast, anti-aminophospholipid antibodies, bispecific antibodies, and antigen binding fragments thereof, that are conjugated to, or operatively associated with, cytotoxic or anticellular agents are referred to herein as "anti-aminophospholipid immunotoxins". Likewise, anti-aminophospholipid antibodies, bispecific antibodies, and antigen binding fragments thereof, that are conjugated to, or operatively associated with, coagulation factors are herein referred to as "anti-aminophospholipid coaguligands". The use of anti-aminophospholipid immunotoxins and
25 anti-aminophospholipid coaguligands in tumor treatment is also contemplated by the present inventors and is disclosed and claimed in first and second provisional applications Serial Nos. 60/092,589 (filed July 13, 1998) and 60/110,600 (filed December 02, 1998) and in co-filed U.S. and PCT patent applications (Attorney Docket Nos. 4001.002300, 4001.002382, 4001.002383 and 4001.002210), each specifically incorporated herein by reference.

Preferred anti-aminophospholipid antibodies for use in the present invention are anti-phosphatidylserine (anti-PS) and anti-phosphatidylethanolamine (anti-PE) antibodies. Anti-PS antibodies will generally recognize, bind to or have immunospecificity for the PS molecule present, expressed, translocated, presented or complexed at the luminal surface of tumor vascular endothelial cells. Suitable antibodies will thus bind to phosphatidyl-L-serine (Umeda *et al.*, 1989; incorporated herein by reference). Anti-PE antibodies will generally recognize, bind to or have immunospecificity for the PE molecule present, expressed, translocated, presented or complexed at the luminal surface of tumor vascular endothelial cells.

Administering anti-aminophospholipid antibodies to an animal with a tumor will result in specific binding of the antibody to the aminophospholipid molecules present, expressed or translocated to the luminal surface of the tumor blood vessels, *i.e.*, the antibodies will bind to the aminophospholipid molecules in a natural, biological environment. Therefore, no particular manipulation will be necessary to ensure antibody binding.

However, it is of scientific interest to note that aminophospholipids may be most frequently recognized, or bound, by anti-aminophospholipid antibodies when the aminophospholipid molecules are associated with one or more proteins or other non-lipid biological components. For example, anti-PS antibodies that occur as a sub-set of anti-phospholipid (anti-PL) antibodies in patients with certain diseases and disorders are now believed to bind to PS in combination with proteins such as β_2 -glycoprotein I (β_2 -GPI or apolipoprotein H, apoH) and prothrombin (U.S. Patent No. 5,344,758; Rote, 1996; each incorporated herein by reference). Similarly, anti-PE antibodies that occur in disease states are now believed to bind to PE in combination with proteins such as low and high molecular weight kininogen (HK), prekallikrein and even factor XI (Sugi and McIntyre, 1995; 1996a; 1996b; each incorporated herein by reference).

This is the meaning of the terms "presented" and "complexed at" the luminal surface of tumor blood vessels, as used herein, which mean that the aminophospholipid molecules are present at the surface of tumor blood vessels in an antibody-binding competent state,

irrespective of the molecular definition of that particular state. PS may even be targeted as a complex with factor II/IIa, VII/VIIa, IX/IXa and X/Xa. Moreover, the nature of the aminophospholipid target may change during practice of the invention, as the initial aminophospholipid antibody binding, anti-endothelial cell and anti-tumor effects may result in biological changes that alter the number, conformation and/or type of the aminophospholipid target epitope(s).

The term "anti-aminophospholipid antibody", as used in the context of the present invention, therefore means any naked or unconjugated anti-aminophospholipid antibody, immunological binding agent or antisera; monoclonal, human, humanized, dimeric, trimeric, multimeric, chimeric, bispecific, recombinant or engineered antibody; or Fab', Fab, F(ab')₂, DABs, Fv or scFv antigen binding fragment thereof; that at least binds to a lipid and amino group-containing complex or aminophospholipid target, preferably a phosphatidylserine- or phosphatidylethanolamine-based target.

The requirement that the antibody "at least bind to an aminophospholipid target" is met by the antibody binding to any and/or all physiologically relevant forms of aminophospholipids, including so-called "hexagonal" and "hexagonal phase II" PS and PE (HexII PS and HexII PE) (Rauch *et al.*, 1986; Rauch and Janoff, 1990; Berard *et al.*, 1993; each incorporated herein by reference) and PS and PE in combination with any other protein, lipid, membrane component, plasma or serum component, or any combination thereof. Thus, an "anti-aminophospholipid antibody" is an antibody that binds to an aminophospholipid in the tumor blood vessels, notwithstanding the fact that bilayer or micelle aminophospholipids may be considered to be immunogenically neutral.

The anti-aminophospholipid antibodies may recognize, bind to or have immunospecificity for aminophospholipid molecules, or an immunogenic complex thereof (including hexagonal aminophospholipids and protein combinations), to the exclusion of other phospholipids or lipids. Such antibodies may be termed "aminophospholipid-specific or aminophospholipid-restricted antibodies", and their use in the invention will often be preferred.

"Aminophospholipid-specific or aminophospholipid-restricted antibodies" will generally exhibit significant binding to aminophospholipids, while exhibiting little or no significant binding to other lipid components, such as phosphatidylinositol (PI), phosphatidylglycerol (PG) and even phosphatidylcholine (PC) in certain embodiments.

5

"PS-specific or PS-restricted antibodies" will generally exhibit significant binding to PS, while exhibiting little or no significant binding to lipid components such as phosphatidylethanolamine and cardiolipin (CL), as well as PC, PI and PG. "PE-specific or PE-restricted antibodies" will generally exhibit significant binding to PE, while exhibiting little or no significant binding to lipid components such as phosphatidylserine and cardiolipin, as well as PC, PI and PG. The preparation of specific anti-aminophospholipid antibodies is readily achieved, *e.g.*, as disclosed by Rauch *et al.* (1986); Umeda *et al.* (1989); Rauch and Janoff (1990); and Rote *et al.* (1993); each incorporated herein by reference.

10

09351862-071299

15

"Cross-reactive anti-aminophospholipid antibodies" that recognize, bind to or have immunospecificity for an aminophospholipid molecule, or an immunogenic complex thereof (including hexagonal aminophospholipids and protein combinations), in addition to exhibiting lesser but detectable binding to other phospholipid or lipid components are by no means excluded from use in the invention. Such "cross-reactive anti-aminophospholipid antibodies" may be employed so long as they bind to an aminophospholipid present, expressed, translocated, presented or complexed at the luminal surface of tumor vascular endothelial cells and exert an anti-tumor effect upon administration *in vivo*.

20

Further suitable aminophospholipid-specific or aminophospholipid-restricted antibodies are those anti-aminophospholipid antibodies that bind to both PS and PE. While clearly being specific or restricted to aminophospholipids, as opposed to other lipid components, antibodies exist that bind to each of the preferred targets of the present invention. Examples of such antibodies for use in the invention include, but are not limited to, PS3A, PSF6, PSF7, PSB4, PS3H1 and PS3E10 (Igarashi *et al.*, 1991; incorporated herein by reference)

30

Further exemplary anti-PS antibodies for use in the invention include, but are not limited to BA3B5C4, PS4A7, PS1G3 and 3SB9b; with PS4A7, PS1G3 and 3SB9b generally being preferred. Monoclonal antibodies, humanized antibodies and/or antigen-binding fragments based upon the 3SB9b antibody (Rote *et al.*, 1993; incorporated herein by reference) are currently most preferred.

Although aminophospholipids, such as PS and PE, in bilayer or micelle form have been reported to be non- or weakly antigenic, or non- or weakly-immunogenic, the scientific literature has reported no difficulties in generating anti-aminophospholipid antibodies, such as anti-PS and anti-PE antibodies. Anti-aminophospholipid antibodies or monoclonal antibodies may therefore be readily prepared by preparative processes and methods that comprise:

- (a) preparing an anti-aminophospholipid antibody-producing cell; and
- (b) obtaining an anti-aminophospholipid antibody or monoclonal antibody from the antibody-producing cell.

The processes of preparing anti-aminophospholipid antibody-producing cells and obtaining anti-aminophospholipid antibodies therefrom may be conducted *in situ* in a given patient. That is, simply providing an immunogenically effective amount of an immunogenic aminophospholipid sample to a patient will result in anti-aminophospholipid antibody generation. Thus, the anti-aminophospholipid antibody is still "obtained" from the antibody-producing cell, but it does not have to be isolated away from a host and subsequently provided to a patient, being able to spontaneously localize to the tumor vasculature and exert its biological anti-tumor effects.

As disclosed herein, anti-aminophospholipid antibody-producing cells may be obtained, and antibodies subsequently isolated and/or purified, from human patients with anti-aminophospholipid antibody-producing diseases, from stimulating peripheral blood lymphocytes with aminophospholipids *in vitro*, and also by immunization processes and methods. The latter of which generally comprise:

- 5
- (a) immunizing an animal by administering to the animal at least one dose, and optionally more than one dose, of an immunogenically effective amount of an immunogenic aminophospholipid sample (such as a hexagonal, or hexagonal phase II form of an aminophospholipid), preferably an immunogenic PS or PE sample; and
 - (b) obtaining an anti-aminophospholipid antibody-producing cell from the immunized animal.

10

The immunogenically effective amount of the aminophospholipid sample or samples may be a *Salmonella*-coated aminophospholipid sample (Umeda *et al.*, 1989; incorporated herein by reference); an aminophospholipid micelle, liposome, lipid complex or lipid formulation sample; or an aminophospholipid sample fabricated with SDS. Any such aminophospholipid sample may be administered in combination with any suitable adjuvant, such as Freund's complete adjuvant (Rote *et al.*, 1993; incorporated herein by reference). Any empirical technique or variation may be employed to increase immunogenicity, and/or hexagonal or hexagonal phase II forms of the aminophospholipids may be administered.

20 The immunization may be based upon one or more intrasplenic injections of an immunogenically effective amount of an aminophospholipid sample (Umeda *et al.*, 1989; incorporated herein by reference).

25 Irrespective of the nature of the immunization process, or the type of immunized animal, anti-aminophospholipid antibody-producing cells are obtained from the immunized animal and, preferably, further manipulated by the hand of man. "An immunized animal", as used herein, is a non-human animal, unless otherwise expressly stated. Although any antibody-producing cell may be used, most preferably, spleen cells are obtained as the source of the antibody-producing cells. The anti-aminophospholipid antibody-producing cells may be used in a preparative process that
30 comprises:

- (a) fusing an anti-aminophospholipid antibody-producing cell with an immortal cell to prepare a hybridoma that produces an anti-aminophospholipid monoclonal antibody and

5

- (b) obtaining an anti-aminophospholipid monoclonal antibody from the hybridoma.

Hybridoma-based monoclonal antibody preparative methods thus include those that comprise:

10

- (a) immunizing an animal by administering to the animal at least one dose, and optionally more than one dose, of an immunogenically effective amount of an immunogenic aminophospholipid sample (such as a hexagonal, or hexagonal phase II form of an aminophospholipid), preferably an immunogenic PG or PE sample;

15

- (b) preparing a collection of monoclonal antibody-producing hybridomas from the immunized animal;

20

- (c) selecting from the collection at least a first hybridoma that produces at least a first anti-aminophospholipid monoclonal antibody, and preferably, at least a first aminophospholipid-specific monoclonal antibody; and

25

- (d) culturing the at least a first anti-aminophospholipid-producing or aminophospholipid-specific hybridoma to provide the at least a first anti-aminophospholipid monoclonal antibody or aminophospholipid-specific monoclonal antibody; and preferably

- (e) obtaining the at least a first anti-aminophospholipid monoclonal antibody or aminophospholipid-specific monoclonal antibody from the cultured at least a first hybridoma.

5 As non-human animals are used for immunization, the anti-aminophospholipid monoclonal antibodies obtained from such a hybridoma will often have a non-human make up. Such antibodies may be optionally subjected to a humanization process, grafting or mutation, as known to those of skill in the art and further disclosed herein. Alternatively, transgenic animals, such as mice, may be used that comprise a human antibody gene library. Immunization of such
10 animals will therefore directly result in the generation of human anti-aminophospholipid antibodies.

After the production of a suitable antibody-producing cell, most preferably a hybridoma, whether producing human or non-human antibodies, the monoclonal antibody-encoding nucleic
15 acids may be cloned to prepare a "recombinant" monoclonal antibody. Any recombinant cloning technique may be utilized, including the use of PCR to prime the synthesis of the antibody-encoding nucleic acid sequences. Therefore, yet further appropriate monoclonal antibody preparative methods include those that comprise using the anti-aminophospholipid antibody-producing cells as follows:

- 20
- (a) obtaining at least a first anti-aminophospholipid antibody-encoding nucleic acid molecule or segment from an anti-aminophospholipid antibody-producing cell, preferably a hybridoma; and
- 25 (b) expressing the nucleic acid molecule or segment in a recombinant host cell to obtain a recombinant anti-aminophospholipid monoclonal antibody.

However, other powerful recombinant techniques are available that are ideally suited to the preparation of recombinant monoclonal antibodies. Such recombinant techniques include the
30 phagemid library-based monoclonal antibody preparative methods comprising:

5 (a) immunizing an animal by administering to the animal at least one dose, and optionally more than one dose, of an immunogenically effective amount of an immunogenic aminophospholipid sample (such as a hexagonal, or hexagonal phase II form of an aminophospholipid), preferably an immunogenic PS or PE sample;

10 (b) preparing a combinatorial immunoglobulin phagemid library expressing RNA isolated from the antibody-producing cells, preferably from the spleen, of the immunized animal;

15 (c) selecting from the phagemid library at least a first clone that expresses at least a first anti-aminophospholipid antibody, and preferably, at least a first aminophospholipid-specific antibody;

20 (d) obtaining anti-aminophospholipid antibody-encoding nucleic acids from the at least a first selected clone and expressing the nucleic acids in a recombinant host cell to provide the at least a first anti-aminophospholipid antibody or aminophospholipid-specific antibody; and preferably

(e) obtaining the at least a first anti-aminophospholipid antibody or aminophospholipid-specific antibody expressed by the nucleic acids obtained from the at least a first selected clone.

25 Again, in such phagemid library-based techniques, transgenic animals bearing human antibody gene libraries may be employed, thus yielding recombinant human monoclonal antibodies.

30 Irrespective of the manner of preparation of a first anti-aminophospholipid antibody nucleic acid segment, further suitable anti-aminophospholipid antibody nucleic acid segments

may be readily prepared by standard molecular biological techniques. In order to confirm that any variant, mutant or second generation anti-aminophospholipid antibody nucleic acid segment is suitable for use in the present invention, the nucleic acid segment will be tested to confirm expression of an antibody that binds to an aminophospholipid. Preferably, the variant, mutant or
5 second generation anti-aminophospholipid antibody nucleic acid segment will also be tested to confirm hybridization to an anti-aminophospholipid antibody nucleic acid segment under standard, more preferably, standard stringent hybridization conditions. Exemplary suitable hybridization conditions include hybridization in about 7% sodium dodecyl sulfate (SDS), about 0.5 M NaPO₄, about 1 mM EDTA at about 50°C; and washing with about 1% SDS at about
10 42°C.

As a variety of recombinant monoclonal antibodies, whether human or non-human in origin, may be readily prepared, the treatment methods of the invention may be executed by providing to the animal or patient at least a first nucleic acid segment that expresses a biologically
15 effective amount of at least a first anti-aminophospholipid antibody in the patient. The "nucleic acid segment that expresses an anti-aminophospholipid antibody" will generally be in the form of at least an expression construct, and may be in the form of an expression construct comprised within a virus or within a recombinant host cell. Preferred gene therapy vectors of the present invention will generally be viral vectors, such as comprised within a recombinant retrovirus, herpes simplex virus (HSV), adenovirus, adeno-associated virus (AAV), cytomegalovirus
20 (CMV), and the like.

In certain embodiments, the vasculature of the vascularized tumor of the animal or patient to be treated may be first imaged. Generally this is achieved by first administering to the
25 animal or patient a diagnostically effective amount of at least a first pharmaceutical composition comprising at least a first detectably-labeled aminophospholipid binding construct, such as an anti-aminophospholipid antibody-detectable agent construct, that binds to and identifies an aminophospholipid, preferably phosphatidylserine or phosphatidylethanolamine, present, expressed, translocated, presented or complexed at the luminal surface of blood vessels of the
30 vascularized tumor. The invention thus further provides compositions for use in, and methods

of, distinguishing between tumor and/or intratumoral blood vessels and normal blood vessels. The "distinguishing" is achieved by administering one or more of the detectably-labeled aminophospholipid binding constructs described.

- 5 The detectably-labeled aminophospholipid binding construct or anti-aminophospholipid antibody-detectable agent construct may comprise an X-ray detectable compound, such as bismuth (III), gold (III), lanthanum (III) or lead (II); a radioactive ion, such as copper⁶⁷, gallium⁶⁷, gallium⁶⁸, indium¹¹¹, indium¹¹³, iodine¹²³, iodine¹²⁵, iodine¹³¹, mercury¹⁹⁷, mercury²⁰³, rhenium¹⁸⁶, rhenium¹⁸⁸, rubidium⁹⁷, rubidium¹⁰³, technetium^{99m} or yttrium⁹⁰; a nuclear magnetic
- 10 spin-resonance isotope, such as cobalt (II), copper (II), chromium (III), dysprosium (III), erbium (III), gadolinium (III), holmium (III), iron (II), iron (III), manganese (II), neodymium (III), nickel (II), samarium (III), terbium (III), vanadium (II) or ytterbium (III); or rhodamine or fluorescein .

Pre-imaging before tumor treatment may thus be carried out by.

00361062-071200

- 15 (a) administering to the animal or patient a diagnostically effective amount of a pharmaceutical composition comprising at least a first detectably-labeled aminophospholipid binding construct that comprises a diagnostic agent operatively attached to an antibody, binding protein or ligand, or aminophospholipid binding fragment thereof, that binds to an aminophospholipid, preferably phosphatidylserine or phosphatidylethanolamine, present, expressed, translocated, presented or complexed at the luminal surface of blood vessels or intratumoral blood vessels of the vascularized tumor; and
- 25 (b) subsequently detecting the detectably-labeled aminophospholipid binding construct bound to an aminophospholipid, preferably phosphatidylserine or phosphatidylethanolamine, on the luminal surface of tumor or intratumoral blood vessels, thereby obtaining an image of the tumor vasculature.

Cancer treatment may also be carried out by:

(a) forming an image of a vascularized tumor by administering to an animal or patient having a vascularized tumor a diagnostically minimal amount of at least a first detectably-labeled aminophospholipid binding construct comprising a diagnostic agent operatively attached to an antibody, binding protein or ligand, or aminophospholipid binding fragment thereof, that binds to an aminophospholipid, preferably phosphatidylserine or phosphatidylethanolamine, on the luminal surface of tumor or intratumoral blood vessels of the vascularized tumor, thereby forming a detectable image of the tumor vasculature; and

(b) subsequently administering to the same animal or patient a therapeutically optimized amount of at least a first naked antibody, or antigen-binding fragment thereof, that binds to an aminophospholipid, preferably phosphatidylserine or phosphatidylethanolamine, on the tumor blood vessel luminal surface and thereby destroys the tumor vasculature.

Imaging and treatment formulations or medicaments are thus provided, which generally comprise:

(a) a first pharmaceutical composition comprising a diagnostically effective amount of a detectably-labeled aminophospholipid binding construct that comprises a detectable agent operatively attached to an antibody, binding protein or ligand, or aminophospholipid binding fragment thereof, that binds to an aminophospholipid, preferably phosphatidylserine or phosphatidylethanolamine, on the luminal surface of tumor or intratumoral blood vessels of the vascularized tumor; and

(b) at least a second pharmaceutical composition comprising a therapeutically effective amount of at least a second, unconjugated anti-aminophospholipid

antibody, preferably anti-phosphatidylserine or anti-phosphatidylethanolamine, or antigen binding fragment thereof.

In such methods and medicaments, advantages will be realized wherein the first and
5 second pharmaceutical compositions comprise the same targeting agents, *e.g.*, anti-aminophospholipid antibodies, or fragments thereof, from the same antibody preparation, or preferably, from the same antibody-producing hybridoma.

In the vasculature imaging aspects of the invention, it is recognized that the administered
10 detectably-labeled aminophospholipid binding construct, or anti-aminophospholipid antibody-detectable agent, may itself have a therapeutic effect. Whilst this would not be excluded from the invention, the amounts of the detectably-labeled constructs to be administered would generally be chosen as "diagnostically effective amounts", which are typically lower than the amounts required for therapeutic benefit.

In the imaging embodiments, the targeting agent may be either antibody-based or binding
15 ligand- or binding protein-based. Although not previously connected with tumors or tumor vasculature, detectably labeled aminophospholipid binding ligand compositions are known in the art and can now, in light of this motivation and the present disclosure, be used in the combined
20 imaging and treatment aspects of the present invention. The detectably-labeled annexins of U.S. Patent No. 5,627,036; WO 95/19791; WO 95/27903; WO 95/34315; WO 96/17618; and WO 98/04294; each incorporated herein by reference; may thus be employed.

The foregoing imaging and treatment formulations or medicaments may also further
25 comprise one or more anti-cancer agents. That is, the present invention encompasses imaging and combination treatment formulations and medicaments that generally comprise (a) diagnostically effective amounts of detectably-labeled aminophospholipid binding constructs; (b) therapeutically effective amounts of unconjugated anti-aminophospholipid antibodies, preferably anti-phosphatidylserine or anti-phosphatidylethanolamine, or antigen binding
30 fragments thereof; and (c) therapeutically effective amounts of at least other anti-cancer agent(s).

In still further embodiments, the animals or patients to be treated by the present invention are further subjected to surgery or radiotherapy, or are provided with a therapeutically effective amount of at least a first anti-cancer agent. The "at least a first anti-cancer agent" in this context means "at least a first anti-cancer agent in addition to the naked anti-aminophospholipid antibody" (preferably anti-phosphatidylserine or anti-phosphatidylethanolamine). The "at least a first anti-cancer agent" may thus be considered to be "at least a second anti-cancer agent", where the naked anti-aminophospholipid antibody is a first anti-cancer agent. However, this is purely a matter of semantics, and the practical meaning will be clear to those of ordinary skill in the art.

The at least a first anti-cancer agent may be administered to the animal or patient substantially simultaneously with the anti-aminophospholipid antibody, or antigen-binding fragment thereof; such as from a single pharmaceutical composition or from two pharmaceutical compositions administered closely together.

Alternatively, the at least a first anti-cancer agent may be administered to the animal or patient at a time sequential to the administration of the at least a first anti-aminophospholipid antibody, or antigen-binding fragment thereof. "At a time sequential", as used herein, means "staggered", such that the at least a first anti-cancer agent is administered to the animal or patient at a time distinct to the administration of the at least a first anti-aminophospholipid antibody. Generally, the two agents are administered at times effectively spaced apart to allow the two agents to exert their respective therapeutic effects, *i.e.*, they are administered at "biologically effective time intervals".

The at least a first anti-cancer agent may be administered to the animal or patient at a biologically effective time prior to the anti-aminophospholipid antibody or fragment thereof, or at a biologically effective time subsequent to the anti-aminophospholipid antibody fragment. Administration of one or more non-aminophospholipid targeted anti-cancer agents at a therapeutically effective time subsequent to an anti-aminophospholipid antibody may be particularly desired wherein the anti-cancer agent is an anti-tumor cell immunotoxin designed to

kill tumor cells at the outermost rim of the tumor, and/or wherein the anti-cancer agent is an anti-angiogenic agent designed to prevent micrometastasis of any remaining tumor cells. Such considerations will be known to those of skill in the art.

5 Administration of one or more non-aminophospholipid targeted anti-cancer agents at a therapeutically effective time prior to an anti-aminophospholipid antibody may be particularly employed where the anti-cancer agent is designed to increase aminophospholipid expression. This may be achieved by using anti-cancer agents that injure, or induce apoptosis in, the tumor endothelium. Exemplary anti-cancer agent include, *e.g.*, taxol, vincristine, vinblastine,
10 neomycin, combretastatin(s), podophyllotoxin(s), TNF- α , angiostatin, endostatin, vasculostatin, $\alpha_v\beta_3$ antagonists, calcium ionophores, calcium-flux inducing agents, any derivative or prodrug thereof.

09351862-071209
15 The one or more additional anti-cancer agents may be chemotherapeutic agents, radiotherapeutic agents, cytokines, anti-angiogenic agents, apoptosis-inducing agents or anti-cancer immunotoxins or coaguligands. "Chemotherapeutic agents", as used herein, refer to classical chemotherapeutic agents or drugs used in the treatment of malignancies. This term is used for simplicity notwithstanding the fact that other compounds may be technically described as chemotherapeutic agents in that they exert an anti-cancer effect. However, "chemotherapeutic" has
20 come to have a distinct meaning in the art and is being used according to this standard meaning.

A number of exemplary chemotherapeutic agents are described herein. Those of ordinary skill in the art will readily understand the uses and appropriate doses of chemotherapeutic agents, although the doses may well be reduced when used in combination with the present invention. A
25 new class of drugs that may also be termed "chemotherapeutic agents" are agents that induce apoptosis. Any one or more of such drugs, including genes, vectors and antisense constructs, as appropriate, may also be used in conjunction with the present invention.

Anti-cancer immunotoxins or coaguligands are further appropriate anti-cancer agents.
30 "Anti-cancer immunotoxins or coaguligands", or targeting-agent/therapeutic agent constructs, are

based upon targeting agents, including antibodies or antigen binding fragments thereof, that bind to a targetable component of a tumor cell, tumor vasculature or tumor stroma, and that are operatively attached to a therapeutic agent, generally a cytotoxic agent (immunotoxin) or coagulation factor (coaguligand). A "targetable component" of a tumor cell, tumor vasculature or tumor stroma, is preferably a surface-expressed, surface-accessible or surface-localized component, although components released from necrotic or otherwise damaged tumor cells or vascular endothelial cells may also be targeted, including cytosolic and/or nuclear tumor cell antigens.

Both antibody and non-antibody targeting agents may be used, including growth factors, such as VEGF and FGF; peptides containing the tripeptide R-G-D, that bind specifically to the tumor vasculature; and other targeting components such as annexins and related ligands.

Anti-tumor cell immunotoxins or coaguligands may comprise antibodies exemplified by the group consisting of B3 (ATCC HB 10573), 260F9 (ATCC HB 8488), D612 (ATCC HB 9796) and KS1/4, said KS1/4 antibody obtained from a cell comprising the vector pGKC2310 (NRRL B-18356) or the vector pG2A52 (NRRL B-18357).

Anti-tumor stroma immunotoxins or coaguligands will generally comprise antibodies that bind to a connective tissue component, a basement membrane component or an activated platelet component; as exemplified by binding to fibrin, RIBS or LIBS.

Anti-tumor vasculature immunotoxins or coaguligands may comprise ligands, antibodies, or fragments thereof, that bind to a surface-expressed, surface-accessible or surface-localized component of the blood transporting vessels, preferably the intratumoral blood vessels, of a vascularized tumor. Such antibodies include those that bind to surface-expressed components of intratumoral blood vessels of a vascularized tumor, including aminophospholipids themselves, and intratumoral vasculature cell surface receptors, such as endoglin (TEC-4 and TEC-11 antibodies), a TGF β receptor, E-selectin, P-selectin, VCAM-1, ICAM-1, PSMA, a VEGF/VPF receptor, an FGF receptor, a TIE, $\alpha_v\beta_3$ integrin, pleiotropin, endosialin and MHC Class II

proteins. The antibodies may also bind to cytokine-inducible or coagulant-inducible components of intratumoral blood vessels.

Other anti-tumor vasculature immunotoxins or coaguligands may comprise antibodies, or
5 fragments thereof, that bind to a ligand or growth factor that binds to an intratumoral vasculature
cell surface receptor. Such antibodies include those that bind to VEGF/VPF (GV39 and GV97
antibodies), FGF, TGF β , a ligand that binds to a TIE, a tumor-associated fibronectin isoform,
scatter factor/hepatocyte growth factor (HGF), platelet factor 4 (PF4), PDGF and TIMP. The
antibodies, or fragments thereof, may also bind to a ligand:receptor complex or a growth
10 factor:receptor complex, but not to the ligand or growth factor, or to the receptor, when the
ligand or growth factor or the receptor is not in the ligand:receptor or growth factor:receptor
complex.

Anti-tumor cell, anti-tumor stroma or anti-tumor vasculature antibody-therapeutic agent
15 constructs may comprise anticellular, cytostatic or cytotoxic agents such as plant-, fungus- or
bacteria-derived toxins (immunotoxins). Ricin A chain and deglycosylated ricin A chain will
often be preferred, and gelonin and angiopoietins are also contemplated. Anti-tumor cell, anti-
tumor stroma or anti-tumor vasculature antibody-therapeutic agent constructs may comprise
coagulation factors or second antibody binding regions that bind to coagulation factors
20 (coaguligands). The operative association with Tissue Factor or Tissue Factor derivatives, such
as truncated Tissue Factor, will often be preferred.

The present invention yet further provides a series of novel therapeutic kits, medicaments
and/or cocktails for use in conjunction with the methods of the invention. The kits, medicaments
25 and/or cocktails generally comprise a combined effective amount of an anti-cancer agent and an
antibody, or an antigen-binding fragment thereof, that binds to an aminophospholipid, preferably
phosphatidylserine or phosphatidylethanolamine.

Where the primary purpose of a kit of the invention is in combination therapy, the kit
30 may nonetheless still further comprise an imaging component, generally a diagnostically

effective amount of a detectably-labeled aminophospholipid binding construct, such as a labeled anti-aminophospholipid antibody or antigen binding fragment thereof.

5 The kits and medicaments will comprise, preferably in suitable container means, a biologically effective amount of at least a first antibody, or an antigen-binding fragment thereof, that binds to an aminophospholipid, preferably phosphatidylserine or phosphatidylethanolamine; in combination with a biologically effective amount of at least a first anti-cancer agent. The components of the kits and medicaments may be comprised within a single container or container means, or comprised within distinct containers or container means. The cocktails will generally
10 be admixed together for combined use.

15 The entire range of anti-aminophospholipid antibodies, as described above, may be employed in the kits, medicaments and/or cocktails, with anti-PS, anti-PE, human, humanized and monoclonal antibodies, or fragments thereof, being preferred. The anti-cancer agents are also those as described above, including chemotherapeutic agents, radiotherapeutic agents, anti-angiogenic agents, apoptotic agents, immunotoxins and coaguligands. Agents formulated for intravenous administration will often be preferred.

BRIEF DESCRIPTION OF THE DRAWINGS

20 The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

25 **FIG. 1A and FIG. 1B.** Activity of cell-bound anti-VCAM-1•tTF *in vitro*. **FIG. 1A.** Binding of anti-VCAM-1•tTF coaguligand to unstimulated (control) and IL-1 α -activated bEnd.3 cells. **FIG. 1B.** Generation of factor Xa by cell-bound anti-VCAM-1•tTF coaguligand.

30 **FIG. 2.** Retardation of growth of L540 tumors in mice treated with anti-VCAM-1•tTF. L540 tumor bearing mice were injected i.v. with either saline, 20 μ g of anti-VCAM-1•tTF, 4 μ g

of unconjugated tTF or 20 µg of control IgG•tTF. Injections were repeated on day 4 and 8 after the first treatment. Tumors were measured daily. Mean tumor volume and SD of 8 mice per group is shown.

5 **FIG. 3.** Annexin V blocks coaguligand activation of Factor X *in vitro*. IL-1α-stimulated bEnd.3 cells were incubated with anti-VCAM•tTF coaguligand in 96-well microtiter plates, as described in Example V. Annexin V was added at concentrations ranging from 0.1 to 10 µg/ml (as shown) and cells were incubated for 30 min. before addition of diluted Proplex T. The amount of Factor Xa generated in the presence or absence of Annexin V was determined using a
10 chromogenic substrate, as described in Example V.

15 **FIG. 4A and FIG. 4B.** Anti-tumor effects of naked anti-PS antibodies in animals with syngeneic and xenogeneic tumors. 1x10⁷ cells of murine colorectal carcinoma Colo 26 (**FIG. 4A**) or human Hodgkin's lymphoma L540 (**FIG. 4B**) were injected subcutaneously into the right flank of Balb/c mice (**FIG. 4A**) or male CB17 SCID mice (**FIG. 4B**), respectively. Tumors were allowed to grow to a size of about 0.6-0.9 cm³ and then the mice (4 animals per group) were injected i.p. with 20 µg of naked anti-PS antibody (open squares) or saline (open circles) (control mouse IgM gave similar results to saline.). Treatment was repeated 3 times with a 48 hour interval. Animals were monitored daily for tumor measurements and body weight.
20 Tumor volume was calculated as described in Example VII. Mice were sacrificed when tumors had reached 2 cm³, or earlier if tumors showed signs of necrosis or ulceration.

DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

A. Tumor Destruction using VCAM-1 Coaguligand

25 Solid tumors and carcinomas account for more than 90% of all cancers in man. Although the use of monoclonal antibodies and immunotoxins has been investigated in the therapy of lymphomas and leukemias (Vitetta *et al.*, 1991), these agents have been disappointingly ineffective in clinical trials against carcinomas and other solid tumors (Abrams and Oldham, 1985). A principal reason for the ineffectiveness of antibody-based treatments is that

macromolecules are not readily transported into solid tumors. Even once within a tumor mass, these molecules fail to distribute evenly due to the presence of tight junctions between tumor cells, fibrous stroma, interstitial pressure gradients and binding site barriers (Dvorak *et al.*, 1991).

5 In developing new strategies for treating solid tumors, the methods that involve targeting the vasculature of the tumor, rather than the tumor cells, offer distinct advantages. An effective destruction or blockade of the tumor vessels arrests blood flow through the tumor and results in an avalanche of tumor cell death. Antibody-toxin and antibody-coagulant constructs have already been used to great effect in the specific targeting and destruction of tumor vessels,
10 resulting in tumor necrosis (Burrows *et al.*, 1992; Burrows and Thorpe, 1993; WO 93/17715; WO 96/01653; Huang *et al.*, 1997; each incorporated herein by reference).

Tumor vasculature-targeted cytotoxic agents are described in the following patents and patent applications; U.S. Patents Nos. 5,855,866; 5,776,427; 5,863,538; ~~5,660,827~~ ^{5,660,827; 6,004,554;} ~~5,917,113 and 6,051,250~~; and U.S. Application Serial No. ~~07/846,349~~ ^{107/846,349} Applications Serial No. 07/846,349; 08/295,868 (Patent No. 5,____; Issue Fee paid); 08/350,212 (Patent No. 5,____; Issue Fee paid); and 08/457,869 (Notice of Allowance Received); each incorporated herein by reference. Tumor targeted coagulants are described in the following patents and patent applications: U.S. Patent Nos. ~~5,855,866 and 5,877,289~~ ^{5,855,866; 5,877,289;} ~~5,965,132; 6,004,553; 6,036,955; and 6,043,398~~; and U.S. Application Serial No. ~~07/846,349~~ ^{107/846,349} Applications Serial Nos. 07/846,349; 08/350,212 (Patent No. 5,____; Issue Fee paid); 08/482,369 (U.S. Patent 5,____; Issue Fee paid); 08/487,427 (U.S. Patent 5,____; Issue Fee paid); and 08/479,727 (U.S. Patent 5,____; Issue Fee paid); each incorporated herein by reference.

25 Where antibodies, growth factors or other binding ligands are used to specifically deliver a coagulant to the tumor vasculature, such agents are termed "coaguligands". A currently preferred coagulant for use in coaguligands is truncated Tissue Factor (tTF) (Huang *et al.*, 1997; WO 96/01653; U.S. Patent 5,877,289. TF is the major initiator of blood coagulation (Ruf *et al.*, 1991). At sites of injury, Factor VII/VIIa in the blood comes into contact with, and binds to, TF on cells in the perivascular tissues. The TF:VIIa complex, in the presence of the phospholipid

surface, activates factors IX and X. This, in turn, leads to the formation of thrombin and fibrin and, ultimately, a blood clot (Ruf and Edgington, 1994).

5 The recombinant, truncated form of tissue factor (tTF), lacking the cytosolic and transmembrane domains, is a soluble protein that has about five orders of magnitude lower coagulation inducing ability than native TF (Stone *et al.*, 1995; Huang *et al.*, 1997). This is because TF needs to be associated with phospholipids for the complex with VIIa to activate IXa or Xa efficiently. However, when tTF is delivered to tumor vascular endothelium by means of a targeting antibody or agent, it is brought back into proximity to a lipid surface and regains
10 thrombogenic activity (Huang *et al.*, 1997; U.S. Patents 5,877,289, ~~5,877,289~~^{6,004,555 and 6,093,319} and ~~5,877,289~~^{6,004,555 and 6,093,319} (U.S. Applications Serial Nos. 08/487,427 and 08/482,369; Issue Fees paid)). A coaguligand is thus created that selectively thromboses tumor vasculature.

15 Truncated TF has several advantages that commend its use in vascular targeted coaguligands: human tTF is readily available, and the human protein will have negligible or low immunogenicity in man; human tTF is fully functional in experimental animals, including mice; and targeted tTF is highly potent because it triggers the activation of a cascade of coagulation proteins, giving a greatly amplified effect (U.S. Patents 5,877,289, ~~5,877,289~~^{6,004,555 and 6,093,319} and ~~5,877,289~~^{6,004,555 and 6,093,319} (U.S. Applications Serial Nos. 08/487,427 and 08/482,369; Issue Fees paid)).

20 A range of suitable target molecules that are available on tumor endothelium, but largely absent from normal endothelium, have been described. For example, expressed targets may be utilized, such as endoglin, E-selectin, P-selectin, VCAM-1, ICAM-1, PSMA, a TIE, a ligand reactive with LAM-1, a VEGF/VPF receptor, an FGF receptor, $\alpha_v\beta_3$ integrin, pleiotropin or
25 endosialin (U.S. Patents 5,855,866 5,877,289; Burrows *et al.*, 1992; Burrows and Thorpe, 1993; Huang *et al.*, 1997; Liu *et al.*, 1997; Ohizumi *et al.*, 1997; each incorporated herein by reference).

30 Adsorbed targets are another suitable group, such as VEGF, FGF, TGF β , HGF, PF4, PDGF, TIMP, a ligand that binds to a TIE or a tumor-associated fibronectin isoform (U.S. Patents 5,877,289, ~~5,877,289~~^{5,965,132 and 6,004,555} and ~~5,877,289~~^{5,965,132 and 6,004,555}; corresponding to U.S. Serial Nos. 08/350,212 and

A

~~08/487,427; Issue Fees paid~~; each incorporated herein by reference). Fibronectin isoforms are ligands that bind to the integrin family of receptors. Tumor-associated fibronectin isoforms are targetable components of both tumor vasculature and tumor stroma. The monoclonal antibody BC-1 (Carnemolla *et al.*, 1989) specifically binds to tumor-associated fibronectin isoforms.

5

A

Other targets inducible by the natural tumor environment or following intervention by man are also targetable entities, as described in U.S. Patent Nos. 5,776,427, 5,863,538 and ~~5,____~~ ^{6,036,956 (each} (U.S. Serial No. ~~08/479,727; Issue Fee paid~~; each incorporated herein by reference). When used

A

10

in conjunction with prior suppression in normal tissues and tumor vascular induction, MHC Class II antigens may also be employed as targets (U.S. Patent Nos. 5,776,427; 5,863,538; ~~5,____~~ ^{6,004,554 and} and ~~5,____~~ ^{6,036,955} (U.S. Serial Nos. ~~08/295,868 and 08/479,727; Issue Fees paid~~); each incorporated herein by reference).

A

15

One currently preferred target for clinical applications is vascular endothelial adhesion molecule-1 (VCAM-1) (U.S. Patents 5,855,866, 5,877,289, ~~5,____~~ ^{6,004,555 and 6,093,319} and ~~5,____~~ (corresponding to U.S. Serial Nos. ~~08/482,369 and 08/487,427; Issue Fees paid~~); each incorporated herein by reference). VCAM-1 is a cell adhesion molecule that is induced by inflammatory cytokines IL-1 α , IL-4 (Thornhill *et al.*, 1990) and TNF α (Munro, 1993) and whose role *in vivo* is to recruit leukocytes to sites of acute inflammation (Bevilacqua, 1993).

A

08/487,427; Issue Fees paid; each incorporated herein by reference). VCAM-1 is a cell adhesion molecule that is induced by inflammatory cytokines IL-1 α , IL-4 (Thornhill *et al.*, 1990) and TNF α (Munro, 1993) and whose role *in vivo* is to recruit leukocytes to sites of acute inflammation (Bevilacqua, 1993).

20

VCAM-1 is present on vascular endothelial cells in a number of human malignant tumors including neuroblastoma (Patey *et al.*, 1996), renal carcinoma (Droz *et al.*, 1994), non-small lung carcinoma (Staal-van den Brekel *et al.*, 1996), Hodgkin's disease (Patey *et al.*, 1996), and angiosarcoma (Kuzu *et al.*, 1993), as well as in benign tumors, such as angioma (Patey *et al.*, 1996) and hemangioma (Kuzu *et al.*, 1993). Constitutive expression of VCAM-1 in man is confined to a few vessels in the thyroid, thymus and kidney (Kuzu *et al.*, 1993; Bruijn and Dinklo, 1993), and in the mouse to vessels in the heart and lung (Fries *et al.*, 1993).

25

Certain of the data presented herein even further supplement those provided in U.S.

A

30

Patents 5,855,855 and 5,877,289 and ~~5,____~~ ^{6,004,555} (corresponding to U.S. Serial No. ~~08/487,427;~~

A ~~Issue Fee paid~~; each incorporated herein by reference) and show the selective induction of thrombosis and tumor infarction resulting from administration of an anti-VCAM-1-tTF coaguligand. The results presented were generated using mice bearing L540 human Hodgkin lymphoma. When grown as a xenograft in SCID mice, this tumor shows close similarity to the human disease with respect to expression of inflammatory cytokines (Diehl *et al.*, 1985) and the presence of VCAM-1 and other endothelial cell activation molecules on its vasculature.

Using a covalently-linked anti-VCAM-1-tTF coaguligand, in which tTF was directly linked to the anti-VCAM-1 antibody, it is shown herein that the coaguligand localizes selectively to tumor vessels, induces thrombosis of those vessels, causes necrosis to develop throughout the tumor and retards tumor growth in mice bearing solid L540 Hodgkin tumors. Tumors generally needed to be at least about 0.3 cm in diameter to respond to the coaguligand, because VCAM-1 was absent from smaller tumors. Presumably, in small tumors, the levels of cytokines secreted by tumor cells or host cells that infiltrate the tumor are too low for VCAM-1 induction. This is in accordance with the studies in U.S. Patents 5,855,866, 5,877,289, 5,776,427, ~~5,_____ and 6,004,888 and 6,036,983~~ (Serial Nos. 08/487,427 and 08/479,727, ~~Issue Fees paid~~), where the inventions were shown to be most useful in larger solid tumors.

Although VCAM-1 staining was initially observed more in the periphery of the tumor, the coaguligand evidently bound to and occluded blood transporting vessels - - as it was capable of curtailing blood flow in all tumor regions. Furthermore, one of the inventors contemplates that the thrombin generation caused by the initial administration of the coaguligand likely leads to further VCAM-1 induction on central vessels (Sluiter *et al.*, 1993), resulting in an amplified signal and evident destruction of the intratumoral region. This type of coagulant-induced expression of further targetable markers, and hence signal amplification, is also disclosed in U.S. ~~Serial No. 08/481,904 and U.S. Patent No. 6,036,983~~ ~~Serial No. 08/479,727 and 08/481,904 (U.S. Patent 5,_____, Issue Fee paid)~~.

B. Mechanism of VCAM-1-Targeted Tumor Destruction

As shown herein, although localization to VCAM-1-expressing vessels in the heart and lungs of mice was observed upon administration of an anti-VCAM-1 coaguligand, this construct

did not induce thrombosis in such non-tumor sites. Furthermore, the anti-VCAM-1 coaguligand was no more toxic to mice than was a control coaguligand of irrelevant specificity, again indicating that the constitutive expression of VCAM-1 on heart and lung vessels did not lead to toxicity. This data is important to the immediate clinical progress of coaguligand therapy, given
5 that VCAM-1 is a naturally occurring marker of tumor vascular endothelium in humans. However, this phenomenon also provided the inventors with a unique insight, leading to a totally different approach to tumor vasculature destruction.

The inventors sought to understand the mechanism behind the ability of the anti-
10 VCAM-1 coaguligand to bind to the VCAM-1 constitutively expressed on blood vessels in the heart and lungs, and yet not to cause thrombosis in those vessels. There are numerous scientific possibilities for this empirical observation, generally connected with the prothrombotic nature of the tumor environment any fibrinolytic predisposition in the heart and lungs.

15 Generally, there is a biological equilibrium between the coagulation system (fibrin deposition) and the fibrinolytic system (degradation of fibrin by enzymes). However, in malignant disease, particularly carcinomas, this equilibrium is disrupted, resulting in the abnormal activation of coagulation (hypercoagulability or the "prothrombotic state"). Evidence also indicates that various components of these pathways may contribute to the disorderly
20 characteristics of malignancy, such as proliferation, invasion, and metastasis (Zacharski *et al.*, 1993).

Donati (1995) reviewed the complex interplay between the original clinical observations of thrombotic complications of malignant diseases, and the subsequent progress in the cell
25 biology and biochemistry of tumor cell activities. However, despite extensive research, a clear molecular explanation for the prothrombotic nature of the tumor environment could not be provided (Donati, 1995). Donati did emphasize, though, the role of tumor cells in this process. It was explained that tumor cells express procoagulant activities, such as tissue thromboplastin and cancer procoagulant (CP) (Donati, 1995). WO 91/07187 also reported a procoagulant
30 activity of tumor cells.

Numerous other studies have also identified the tumor cells themselves as being responsible for the prothrombotic state within a tumor. For example, Nawroth *et al.* (1988) reported that factor(s) elaborated by sarcoma cells enhance the procoagulant response of nearby endothelium to TNF. These authors reported that fibrin formation occurred throughout the tumor vascular bed 30 minutes after TNF infusion, but that fibrin deposition and platelet aggregates were not observed in normal vasculature (Nawroth *et al.*, 1988). TNF was later shown to enhance the expression of tissue factor on the surface of endothelial cells (Murray *et al.*, 1991). This was proposed to explain earlier studies showing that cultured endothelial cells incubated with recombinant TNF have enhanced procoagulant activity, tissue factor, and concomitant suppression of the protein C pathway, an anti-thrombotic mechanism that functions on the surface of quiescent endothelial cells (Nawroth *et al.*, 1985; Nawroth and Stern, 1986).

Data from Sugimura *et al.* (1994) also implicated tumor cells as the key components of the procoagulant activity of the tumor. It was reported that four tumor cell lines were able to support different stages of the extrinsic pathway of coagulation (Sugimura *et al.*, 1994). Another study reported that a human ovarian carcinoma cell line, OC-2008, constitutively expressed surface membrane Tissue Factor activity and exhibited cell surface-dependent prothrombinase complex activity (Rao *et al.*, 1992). Connor *et al.* (1989) further suggested that it is the pathologic cells that control coagulation. Their results indicated that tumorigenic, undifferentiated murine erythroleukemic cells exhibit a 7- to 8-fold increase in the potency of their procoagulant activity (Connor *et al.*, 1989).

Zacharski *et al.* (1993) also focused on tumor cells and sought to define the mode of interaction of ovarian carcinoma cells with the coagulation (procoagulant-initiated) and fibrinolysis (urokinase-type plasminogen activator-initiated, u-PA) pathways. They reported that tumor cells expressed Tissue Factor and coagulation pathway intermediates that resulted in local thrombin generation - - as evidenced by the conversion of fibrinogen, present in tumor connective tissue, to fibrin that was found to hug the surfaces of tumor nodules and individual tumor cells. Detected fibrin could not be accounted for on the basis of necrosis or a local

inflammatory cell infiltrate (Zacharski *et al.*, 1993). These authors concluded that there exists a dominant tumor cell-associated procoagulant pathway that leads to thrombin generation and hypercoagulability.

5 Other hypotheses have proposed that it is changes in the tumor blood vessels that render these vessels better able to support the formation of thrombi and/or less able to dissolve fibrin. For example, tumor vessels have been reported to exhibit upregulation of Tissue Factor, down-regulation of plasminogen activators and/or upregulation of the inhibitor of plasminogen activators, PAI-1 (Nawroth and Stern, 1986; Nawroth *et al.*, 1988). Such effects are believed to
10 be magnified by tumor derived factors (Murray *et al.*, 1991; Ogawa *et al.*, 1990), possibly VEGF.

For example, Ogawa *et al.* (1990) reported that hypoxia caused endothelial cell surface coagulant properties to be shifted to promote activation of coagulation. This was accompanied
15 by suppression of the anticoagulant cofactor, thrombomodulin, and induction of an activator of factor X, distinct from the classical extrinsic and intrinsic systems (Ogawa *et al.*, 1990). Also, there could be an increase in the local concentration of Factors VIIa, IXa, Xa, or other molecules that interact with TF, within the tumor vessels, thus encouraging thrombosis.

20 Additionally, platelets are a major component of any procoagulant state. Recently, the procoagulant potential of platelets has been linked to their ability to shed procoagulant microparticles from the plasma membrane (Zwaal *et al.*, 1989; 1992; Dachary-Prigent *et al.*, 1996). It has been proposed that an increased proportion of circulating microparticles, vesicles or membrane fragments from platelets contributes to 'prethrombotic' (prothrombotic) states in
25 various pathological conditions (Zwaal *et al.*, 1989; 1992; Dachary-Prigent *et al.*, 1996, pp. 159 and references cited therein). McNeil *et al.* (1990) also reported that β_2 -GPI exerts multiple inhibitory effects on coagulation and platelet aggregation. Tumor platelet biology could thus explain the effectiveness of the anti-VCAM-1 coaguligand.

Further tenable explanations included the simple possibility that VCAM-1 is expressed at higher levels in tumor vessels than on blood vessels in the heart and lungs, probably due to induction by tumor-derived cytokines, and that binding to the healthy vessels cannot tip the balance into sustained thrombosis. Also the fibrinolytic mechanisms could be upregulated in the heart, as exemplified by increased Tissue Factor pathway inhibitor (TFPI), increased plasminogen activators, and/or decreased plasminogen activator inhibitors. Should the fibrinolytic physiology of the heart and lung vessels prove to be the major reason underlying the tumor-specific effects of the anti-VCAM-1 coaguligand, this would generally preclude the development of additional anti-tumor therapies targeted to unique aspects of tumor biology.

10

Despite all the possible options, the inventors reasoned that the failure of the anti-VCAM-1 coaguligand to cause thrombosis in vessels of normal tissues was due to the absence of the aminophospholipid, phosphatidylserine (PS), from the luminal surface of such vessels. To complete the theory, therefore, not only would phosphatidylserine have to be shown to be absent from these normal vessels, but its presence on the luminal side of tumor-associated vessels would have to be conclusively demonstrated.

15

The inventors therefore used immunohistochemical staining to evaluate the distribution of a monoclonal anti-phosphatidylserine (anti-PS) antibody injected intravenously into tumor-bearing mice. These studies revealed that the VCAM-1 expressing vessels in the heart and lungs lacked PS, whereas the VCAM-1 expressing vessels in the tumor expressed PS. The need for surface PS expression in coaguligand action is further indicated by the inventors' finding that annexin V, which binds to PS, blocks anti-VCAM-1•tTF coaguligand action, both *in vitro* and *in vivo*.

20

25

The lack of thrombotic effect of the anti-VCAM-1 coaguligand on normal heart and lung vessels can thus be explained, at least in part: the absence of the aminophospholipid, phosphatidylserine, means that the normal vessels lack a procoagulant surface upon which coagulation complexes can assemble. In the absence of surface PS, anti-VCAM-1•tTF binds to VCAM-1 expressing heart and lung vessels, but cannot induce thrombosis. In contrast,

30

VCAM-1 expressing vessels in the tumor show coincident expression of surface PS. The coaguligand thus binds to tumor vessels and activates coagulation factors locally to form an occlusive thrombus.

5 In addition to delineating the tumor-specific thrombotic effects of anti-VCAM-1 coaguligands, the specific expression of the aminophospholipid, phosphatidylserine, on the luminal surface of tumor blood vessels also allowed the inventors to explain the prothrombotic phenotype observed, but not understood, in earlier studies (Zacharski *et al.*, 1993; Donati, 1995). Rather than being predominantly due to tumor cells or elaborated factors; platelets, procoagulant
10 microparticles or membrane fragments; or due to imbalances in thromboplastin, thrombomodulin, cancer procoagulant, Tissue Factor, protein C pathway, plasminogen activators or plasminogen activator inhibitors (e.g., PAI-1), the inventors' studies indicate that it is PS expression that plays a significant role in the prothrombotic state of tumor vasculature.

5 C. Aminophospholipids as Markers of Tumor Vasculature

Following their discovery that the representative aminophospholipid, phosphatidylserine, was specifically expressed on the luminal surface of tumor blood vessels, but not in normal blood vessels, the inventors reasoned that aminophospholipids had potential as targets for therapeutic intervention. The present invention therefore encompasses targeting the
15 aminophospholipid constituents, particularly phosphatidylserine (PS) and phosphatidylethanolamine (PE), in tumor treatment. Although anti-tumor effects from aminophospholipid targeting are demonstrated herein, using art-accepted animal models, the ability of aminophospholipids to act as safe and effective targetable markers of tumor vasculature could not have been predicted from previous studies.

25

For example, although tumor vessels are generally prothrombotic in nature, as opposed to other blood vessels, it is an inherent property of the tumor to maintain a network of blood vessels in order to deliver oxygen and nutrients to the tumor cells. Evidently, tumor-associated blood vessels cannot be so predisposed towards thrombosis that they spontaneously and readily
30 support coagulation, as such coagulation would necessarily cause the tumor to self-destruct. It

is thus unexpected that any thrombosis-associated tumor vessel marker, such as the presently identified phosphatidylserine, could be discovered that is expressed in quantities sufficient to allow effective therapeutic intervention by targeting, and yet is expressed at levels low enough to ordinarily maintain blood flow through the tumor.

5

The present identification of aminophospholipids as safe and effective tumor vasculature targets is even more surprising given (1) the previous speculations regarding the role of other cell types and/or various factors, activators and inhibitors underlying the complex, prothrombotic state of the tumor (as discussed above); and (2) the confusing and contradictory state of the art concerning aminophospholipid biology, in terms of both expression and function in various cell types.

10

Phosphatidylserine and phosphatidylethanolamine are normally segregated to the inner surface of the plasma membrane bilayer in different cells (Gaffet *et al.*, 1995; Julien *et al.*, 1995). In contrast, the outer leaflet of the bilayer membrane is rich in phosphatidylcholine analogs (Zwaal *et al.*, 1989; Gaffet *et al.*, 1995). This lipid segregation creates an asymmetric transbilayer. Although the existence of membrane asymmetry has been discussed for some time, the reason for its existence and the mechanisms for its generation and control are poorly understood (Williamson and Schlegel, 1994), particularly in cells other than platelets.

15

00351862-071209
603120-238366

20

There are even numerous conflicting reports regarding the presence or absence of PS and PE in different cells and tissues, let alone concerning the likely role that these aminophospholipids may play. For example, the many PS studies conducted with platelets, key components in blood coagulation (Dachary-Prigent *et al.*, 1996), have yielded highly variable results. Bevers *et al.* (1982) measured the platelet prothrombin-converting activity of non-activated platelets after treatment with various phospholipases or proteolytic enzymes. They concluded that negatively charged phosphatidylserine, and possibly phosphatidylinositol, were involved in the prothrombin-converting activity of non-activated platelets (Bevers *et al.*, 1982).

25

Bervers *et al.* (1983) then reported an increased exposure of phosphatidylserine, and a decreased exposure of sphingomyelinase, in activated platelets. However, these alterations were much less apparent in platelets activated either by thrombin or by collagen alone, in contrast to collagen plus thrombin, diamide, or a calcium ionophore (Bervers *et al.*, 1983). The surface expression of PS in response to diamide was contradicted by studies in erythrocytes, which showed no diamide-stimulated PS exposure (de Jong *et al.*, 1997). While echoing their earlier results, Bervers and colleagues then later reported that changes in the plasma membrane-cytoskeleton interaction, particularly increased degradation of cytoskeletal actin-binding protein, was important to platelet surface changes (Bervers *et al.*, 1985; pages 368-369).

10

Maneta-Peyret *et al.* (1989) also reported the detection of PS on human platelets. These authors noted that the platelet procoagulant surface could be formed by negatively charged phospholipids, such as phosphatidylserine and phosphatidylethanolamine (generally neutral or zwitterionic), or both. The role of phosphatidylserine in the process of coagulation has been questioned in favor of phosphatidylethanolamine (Maneta-Peyret *et al.*, 1989; Schick *et al.*, 1976; 1978). For example, studies have reported that 18% of phosphatidylethanolamine becomes surface-accessible after 2 hours, in contrast to zero phosphatidylserine (Schick *et al.*, 1976).

Ongoing studies with platelets were also reported as showing a further 16% increase in phosphatidylethanolamine exposure after thrombin treatment, with no increase in the phosphatidylserine levels (Schick *et al.*, 1976). Therefore, PS was said not to be a component of the functional surface of the platelet plasma membrane (Schick *et al.*, 1976; 1978). Nonetheless, current evidence does seem to indicate that both PS and PE are involved in the phospholipid asymmetry observed in the outer membrane of platelets and erythrocytes, and that PS is involved in the procoagulant activity of platelets (Gaffet *et al.*, 1995; de Jong *et al.*, 1997; U.S. Patent No. 5,627,036).

The mechanisms for achieving and maintaining differential aminophospholipid distribution, let alone the functional significance of doing so, have long been the subject of

controversial speculations. In reviewing the regulation of transbilayer phospholipid movement, Williamson and Schlegel (1994) indicated that elevating intracellular Ca^{2+} allows the major classes of phospholipids to move freely across the bilayer, scrambling lipids and dissipating asymmetry. de Jong *et al.* (1997) also reported that an increase of intracellular calcium leads to a rapid scrambling of the lipid bilayer and the exposure of PS, which could be partially inhibited by cellular oxidation. The interaction of aminophospholipids with cytoskeletal proteins has also been proposed as a mechanism for regulating membrane phospholipid asymmetry (Zwaal *et al.*, 1989).

Gaffet *et al.* (1995) stated that the transverse redistribution of phospholipids during human platelet activation is achieved by a vectorial outflux of aminophospholipids, not counterbalanced by a rapid reciprocal influx of choline head phospholipids, *i.e.* not scrambling. They suggested that the specific vectorial outflux of aminophospholipids could be catalyzed by a "reverse aminophospholipid translocase" activity (Gaffet *et al.*, 1995). An alternative hypothesis would be that the activity of an inward translocase was inhibited. Zwaal *et al.* (1989) proposed the involvement of a phospholipid-translocase that catalyzed both the outward and inward movement of aminophospholipids.

The presence of an energy- and protein-dependent aminophospholipid translocase activity that transports phosphatidylethanolamine from the outer to the inner leaflet of the lipid bilayer was reported by Julien *et al.* (1993). They then showed that the aminophospholipid translocase activity could also transfer phosphatidylserine, and that the activity could be maintained, suppressed and restored depending on the conditions of cell incubation (Julien *et al.*, 1993), and inhibited by the tumor promoter, 12-*O*-tetradecanoylphorbol-13-acetate (TPA) (Julien *et al.*, 1997).

A 35 kD phospholipid scramblase that promotes the Ca^{2+} -dependent bidirectional movement of phosphatidylserine and other phospholipids was recently cloned from a cDNA library (Zhou *et al.*, 1997). This "PL scramblase" protein is a proline-rich, type II plasma membrane protein with a single transmembrane segment near the C terminus. Subsequent

studies confirmed that this protein was responsible for the rapid movement of phospholipids from the inner to the outer plasma membrane leaflets in cells exposed to elevated cytosolic calcium concentrations (Zhao *et al.*, 1998).

5 The aminophospholipid translocase activity reported by Julien *et al.* (1993; 1997), which transports PS and PE from the outer to the inner leaflet, is different to the bidirectional Ca^{2+} -dependent scramblase (Zhou *et al.*, 1997; Zhao *et al.*, 1998). The scramblase is activated by Ca^{2+} , and mostly functions to move PS from the inner to the outer leaflet in response to increased Ca^{2+} levels. It is now generally believed that the aminophospholipid translocase maintains
10 membrane asymmetry during normal conditions, but that the scramblase is activated by Ca^{2+} influx, over-riding the translocase and randomizing aminophospholipid distribution.

15 The normal segregation of PS and PE to the inner surface of the plasma membrane is thus now generally accepted, and certain membrane components involved in the asymmetric processes have even been identified. However, doubts remain about the conditions, mechanisms and cell types that are capable of re-locating aminophospholipids to the outer leaflet of the membrane, and the biological implications of such events.

20 Contradictory reports concerning aminophospholipid expression are not limited to studies of platelets. Phosphatidylserine and phosphatidylethanolamine are generally about 7% and about 10%, respectively, of the phospholipid composition of cultured human endothelial cells from human artery, saphenous and umbilical vein (7.1% and 10.2%, respectively; Murphy *et al.*, 1992). However, an important example of the contradictions in the literature concerns the ability of anti-PS antibodies to bind to endothelial cells (Lin *et al.*, 1995).

25

 The anti-PS antibodies present in recurrent pregnancy loss (Rote *et al.*, 1995; Rote, 1996; Vogt *et al.*, 1996; Vogt *et al.*, 1997) were believed to modulate endothelial cell function, without evidence of binding to endothelial cells. In an attempt to explain this discrepancy, Lin *et al.* (1995) tried but failed to demonstrate anti-PS antibody binding to resting endothelial cells. They
30 concluded that PS antigenic determinants are not expressed on the surface of resting endothelial

cells, although a PS-dependent antigenic determinant was associated with cytoskeletal-like components in acetone-fixed cells (Lin *et al.*, 1995).

Van Heerde *et al.* (1994) reported that vascular endothelial cells *in vitro* can catalyze the formation of thrombin by the expression of binding sites at which procoagulant complexes can assemble. In contrast to other studies with activated platelets (Beyers *et al.*, 1982; 1983; 1985; Maneta-Peyret *et al.*, 1989; Schick *et al.*, 1976; 1978), stimulated HUVEC endothelial cells did not exhibit an increase in PS binding sites as compared to quiescent cells (Van Heerde *et al.*, 1994). Phosphatidylserine was reported to be necessary for Factor Xa formation via the extrinsic as well as the intrinsic route (Van Heerde *et al.*, 1994). Nonetheless, Brinkman *et al.* (1994) published contradictory results, indicating that other membrane constituents besides negatively charged phospholipids are involved in endothelial cell mediated, intrinsic activation of factor X.

Ravanat *et al.* (1992) also studied the catalytic potential of phospholipids in pro- and anti-coagulant reactions in purified systems and at the surface of endothelial cells in culture after stimulation. Their seemingly contradictory results were proposed to confirm a role for phospholipid-dependent mechanisms in both procoagulant Tissue-Factor activity and anticoagulant activities (activation of protein C by the thrombin-thrombomodulin complex and by Factor Xa) (Ravanat *et al.*, 1992). The Ravanat *et al.* (1992) results were also said to provide evidence of phospholipid exposure during activation of human endothelial cells, which was not observed by Van Heerde *et al.* (1994) or Brinkman *et al.* (1994). However, they did note that anionic phospholipids are of restricted accessibility in the vicinity of cellular Tissue Factor. The situation is further complicated as, even after Tissue Factor induction, other events are likely necessary for coagulation, as the Tissue Factor remains inaccessible, being under the cell.

25

Ravanat *et al.* (1992) went on to suggest that the different extent of inhibition of Tissue Factor and thrombomodulin activities on stimulated endothelial cells means that the cofactor environments differ for the optimal expression of these opposite cellular activities. However, the acknowledged difficulties in trying to reproduce exact cellular phospholipid environments (Ravanat *et al.*, 1992), raise the possibility of artifactual data from these *in vitro* studies. Indeed,

30

PS-dependent phagocytosis marks virally infected cells for phagocytic uptake (WO 97/17084). The surface expression of aminophospholipids could also confer "fusion competence" to a cell (Williamson and Schlegel, 1994).

5 Williamson and Schlegel (1994) also speculated that there is a more general *raison d'être* for lipid asymmetry. For example, although the different head groups have received most attention, it could well be that fatty acid asymmetry is the important factor (Williamson and Schlegel, 1994). A further hypothesis is that the asymmetric distribution of transbilayer phospholipids has no function in itself, but that it is the dynamic process of lipid movement that
10 is important to biological systems (Williamson and Schlegel, 1994).

Many groups have reported that tumor cells are responsible for the prothrombinase activity of the tumor (Connor *et al.*, 1989; Rao *et al.*, 1992; Zacharski *et al.*, 1993; Sugimura *et al.*, 1994; Donati, 1995). This could have been reasoned to be due to PS (WO 91/07187).
15 However, the results of Sugimura *et al.* (1994) argue against this: they reported that although both the prothrombinase activity and total procoagulant activity of the tumorigenic cells, HepG2 and MKN-28, fell on reaching confluency, the PS levels remained constant.

Rather than supporting a role for tumor cell PS in prothrombinase activity, Connor *et al.* (1989) suggested that the increased expression of PS in tumorigenic cells is relevant to their ability to be recognized and bound by macrophages. Utsugi *et al.* (1991) similarly proposed that the presence of PS in the outer membrane of human tumor cells explains their recognition by monocytes.
20

25 Jamasbi *et al.* (1994) suggested a totally different role for lipid components in tumorigenic cells, proposing that the lipids interfere with tumor antigen accessibility. Thus, tumor cell lipids would act to modify the tumor cell surface antigen(s), thus protecting the tumor cells from host immune destruction (Jamasbi *et al.*, 1994). This hypothesis is not unlike that proposed by Qu *et al.* (1996), in terms of endothelial cells. These authors showed that T cells

adhered to thrombin-treated human umbilical endothelial cells by virtue of binding to PS (Qu *et al.*, 1996).

It has thus been proposed that PS-mediated T cell adhesion to endothelial cells *in vivo* is
5 important to both immune surveillance, and also to the disease processes of atherosclerosis (Qu
et al., 1996; Moldovan *et al.*, 1994). Bombeli *et al.* (1997) and Flynn *et al.* (1997) also
suggested that cells within atherosclerotic plaques may contribute to disease progression by
exposing PS, although this was based solely on *in vitro* studies. Qu *et al.* (1996) and Moldovan
et al. (1994) even hinted at an approach opposite to that of the present invention, *i.e.*, the
10 manipulation of phosphatidylserine interactions as an anticoagulant approach. U.S. Patent No.
5,658,877 and No. 5,296,467 have proposed annexin (or "annexine") for use as anti-endotoxins
and anti-coagulants. U.S. Patent No. 5,632,986 (incorporated herein by reference) suggests the
use of the phosphatidylserine-binding ligand, annexin V, as a conjugate with a component, such
as urokinase, that lyses thrombi.

5
Toti *et al.* (1996) suggested that Scott syndrome, an inherited bleeding disorder, may
reflect the deletion or mutation of a putative outward phosphatidylserine translocase or
"scramblase". Although an interesting notion, Stout *et al.* (1997) later isolated a membrane
protein from Scott erythrocytes that exhibited normal PL scramblase activity when reconstituted
20 in vesicles with exogenous PLs. It was suggested that the defect in Scott syndrome is related to
an altered interaction of Ca^{2+} with PL scramblase on the endofacial surface of the cell
membrane, due either to an intrinsic constraint upon the protein, preventing interaction with Ca^{2+}
in situ, or due to an unidentified inhibitor or cofactor in the Scott cell that is dissociated
by detergent (Stout *et al.*, 1997).

25

More variable results have been reported in connection with the possible role of PS in
apoptosis. Williamson and Schlegel (1994) discussed the theme of PS as a marker of
programmed cell death (PCD or apoptosis). It is generally accepted that programmed cell death,
at least in the hematopoietic system, requires the phagocytic sequestration of the apoptotic cells
30 before the loss of membrane integrity or "rupture". The loss of membrane asymmetry in

apoptotic cells, and particularly the appearance of PS in the external leaflet, was proposed to be the trigger for their recognition by phagocytic macrophages (Williamson and Schlegel, 1994).

Martin *et al.* (1995) further reported PS externalization to be an early and widespread event during apoptosis of a variety of murine and human cell types, regardless of the initiating stimulus. They also indicated that, under conditions in which the morphological features of apoptosis were prevented (macromolecular synthesis inhibition, overexpression of Bcl-2 or Abl), the appearance of PS on the external leaflet of the plasma membrane was similarly prevented (Martin *et al.*, 1995).

However, other analyses argue against the Williamson and Schlegel (1994) and Martin *et al.* (1995) proposals to some extent (Vermes *et al.*, 1995). Although these authors indicate that the translocation of PS to the outer membrane surface is a marker of apoptosis, they reason that this is not unique to apoptosis, but also occurs during cell necrosis. The difference between these two forms of cell death is that during the initial stages of apoptosis the cell membrane remains intact, while at the very moment that necrosis occurs the cell membrane loses its integrity and becomes leaky. Therefore, according to this reasoning, PS expression at the cell surface does not indicate apoptosis unless a dye exclusion assay has been conducted to establish cell membrane integrity (Vermes *et al.*, 1995).

Nonetheless, the body of literature prior to the present invention does seem to indicate that the appearance of PS on the outer surface of a cell identifies an apoptotic cell and signals that cell's ingestion (Hampton *et al.*, 1996; WO 95/27903). Hampton *et al.* (1996) concluded that while an elevation of intracellular Ca^{2+} was an ineffective trigger of apoptosis in the cells investigated, extracellular Ca^{2+} was required for efficient PS exposure during apoptosis. In contrast, the proposal of Martin *et al.* (1995) that activation of an inside-outside PS translocase is an early widespread event during apoptosis would seem to require at least some intracellular Ca^{2+} (Zhou *et al.*, 1997; Zhao *et al.*, 1998).

Blankenberg *et al.* (1998) very recently reported that annexin V, an endogenous human protein with a high affinity for PS, can be used to concentrate at sites of apoptotic cell death *in vivo*. Radiolabeled annexin V localized to sites of apoptosis in three models, including acute cardiac allograft rejection (Blankenberg *et al.*, 1998). Staining of cardiac allografts for
5 exogenously administered annexin V revealed myocytes at the periphery of mononuclear infiltrates, of which only a few demonstrated positive apoptotic nuclei.

Finally, the transbilayer movement of phospholipids in the plasma membrane has even been analyzed in ram sperm cells, where the existence of a transverse segregation of
10 phospholipids has been implicated in the fertilization process (Müller *et al.*, 1994). Phospholipid asymmetry has thus been receiving increasing attention, although a clear understanding of this phenomenon, or its relationship to health or disease, has not been realized.

Irrespective of the confusing state of the art regarding aminophospholipid biology, the
15 present inventors discovered, in controlled *in vivo* studies, that aminophospholipids, such as PS and PE, were specific markers of tumor blood vessels. This is surprising in light of the earlier studies of aminophospholipid function, particularly those indicating that the cell surface expression of PS is accompanied by binding of circulating cells, such as T cells (Qu *et al.*, 1996), macrophages (Connor *et al.*, 1989), monocytes (Utsugi *et al.*, 1991) or phagocytes (Zwaal
20 *et al.*, 1989; Williamson and Schlegel, 1994) and is a marker of apoptotic cells (Hampton *et al.*, 1996; Martin *et al.*, 1995; Zhou *et al.*, 1997; Zhao *et al.*, 1998).

Thus, prior to this invention, the possibility of using aminophospholipids as targetable markers of any disease, let alone of tumor vasculature, would be unlikely to be contemplated,
25 due to the perceived masking of these molecules by the binding of one or more cell types, or their transient expression before apoptotic death. In fact, speculative suggestions have concerned the disruption of PS-cellular interactions, such as in preventing leukocyte binding, an initial event in atherosclerosis (Qu *et al.*, 1996).

Other surprising aspects of this discovery are evident in a comparison to earlier work concerning the shedding of procoagulant microparticles from plasma membranes and the demarcation of cells for phagocytosis (WO 97/17084). Zwaal *et al.* (1989; 1992) and Dachary-Prigent *et al.* (1996) explained that PS translocation to the plasma membrane is followed by
5 release of microparticles, microvesicles or microspheres from the cells. Zwaal *et al.* (1989) and Williamson and Schlegel (1994) indicated that PS surface expression prompts clearance by the reticuloendothelial system. In light of these fates of PS-expressing cells, and the various documented bilayer translocase activities (Julien *et al.*, 1995; Zhou *et al.*, 1997; Zhao *et al.*, 1998), it is surprising that cell surface aminophospholipids such as PS and PE can form static and
10 stable enough markers to allow antibody localization and binding.

Prior to the present invention, there was mounting evidence that surface PS appears as part of the apoptotic process, marking cells for rapid destruction (Hampton *et al.*, 1996; Martin *et al.*, 1995). Therefore, although reasonable for use as a diagnostic marker for certain disease
15 states, such as graft rejection (Blankenberg *et al.*, 1998), the apparently limited life time of surface PS would also advise against its use as a viable marker for targeting in therapeutic intervention.

Nonetheless, the present study did indeed discover aminophospholipids to be markers of
20 tumor vascular endothelial cells suitable for targeting. After postulating that PS expression was necessary for VCAM coaguligand action, the presence of PS on tumor blood vessels, but normal vessels, was demonstrated *in vivo*. The *in vivo* observations allowed the inventors to explain the safety and effectiveness of the anti-VCAM coaguligands. This is due to the requirement for coincident expression of a targeted marker (e.g., VCAM) and PS on tumor endothelium. Even if
25 the target molecule is present on endothelium in normal or pathological conditions, thrombosis will not result if surface PS expression is lacking.

The value of the present invention is not limited to explaining coaguligand action, nor to the surprising development of naked antibody therapies. These discoveries have allowed the
30 inventors to show, for the first time, that PS translocation in endothelial cells can occur without

significant cell damage or cell death (Example XIV). In the inventors' new model of tumor biology, the translocation of PS to the surface of tumor blood vessel endothelial cells occurs, at least in a significant part, independently of apoptotic or other cell-death mechanisms. Thus, PS surface expression in the tumor environment is not a consequence of cell death, nor does it trigger immediate cell destruction. This is of fundamental importance and represents a breakthrough in the scientific understanding of PS biology, membrane translocation, cell signaling and apoptosis pathways.

The separation of endothelial cell PS translocation from apoptosis (Example XIV) is also integral to methods of therapeutic intervention based upon PS surface expression. Should PS translocation to the outer membrane in tumor vascular endothelial cells occur only in dying cells, or should it inevitably trigger cell death, then the PS marker would not likely be sufficiently available to serve as a targetable entity for successful therapy (using either naked antibodies or therapeutic conjugates). That is not to say that PS expression on certain tumor vascular endothelial cells is not transient, and that turnover and cell death do not occur in this endothelial cell population, but the finding that significant stable PS expression can be achieved without cell death is a landmark discovery important to various fields of biology and to new therapies.

D. Naked Antibodies Against Aminophospholipids for Tumor Treatment

The present aminophospholipid tumor vasculature expression studies further support the use of coaguligands directed against known tumor vasculature markers as selective thrombotic agents for the treatment of solid tumors. The present observations also led the inventors to develop further tumor treatment methods. For example, the use of anti-aminophospholipid immunotoxins and anti-aminophospholipid coaguligands in tumor treatment is disclosed and claimed in first and second provisional applications Serial Nos. 60/092,589 (filed July 13, 1998) and 60/110,600 (filed December 02, 1998) and in co-filed U.S. and PCT patent applications (Attorney Docket Nos. 4001.002300, 4001.002382, 4001.002383 and 4001.002210), each specifically incorporated herein by reference. The surprising discovery of stable PS expression on intact tumor-associated endothelial cells, which are not undergoing cell death, renders such

methods both practicable and surprising (given that PS expression was thought to be associated only with cell destruction).

However, yet even more unexpected methods of tumor treatment were then discovered.

- 5 In investigating the potential use of aminophospholipid targeting, in the context of later delivering a toxin or coagulant to the tumor vasculature, the inventors serendipitously discovered that naked anti-PS antibodies have a destructive effect on tumor vasculature *in vivo* - - in the complete absence of any additional effector moiety.

- 10 One of the present inventors has been developing anti-tumor vasculature immunotoxins and coaguligands for therapeutic use for some time (for example, see U.S. Patent Nos. 5,585,866 and 5,877,289; and U.S. Applications Serial Nos. 08/350,212, 08/487,427 and 08/482,369; Issue Fees paid; each incorporated herein by reference). In the normal course of these studies, various antibodies, including anti Class II, anti endoglin, anti-VCAM-1 and
15 anti-VEGF, have been administered to tumor-bearing animals and shown to specifically localize to the intratumoral vasculature. Following such confirmation, the antibodies are linked to the toxic or coagulative effector portion to form an immunotoxin or coaguligand, which is then administered to exert an anti-tumor effect.

- 20 During such studies, no naked antibodies have been found to exert an anti-tumor effect in themselves. The ability of anti-aminophospholipid antibodies to both specifically localize to tumor vasculature and to exert a concomitant destructive effect, leading to tumor necrosis, is therefore most unexpected.

- 25 Although a precise molecular understanding of exactly how the naked antibodies function is not necessary in order to practice the present invention, the inventors have contemplated several mechanisms that may account for the observed endothelial cell killing. The favored mechanisms include cell-mediated cytotoxicity, complement-mediated lysis and/or apoptosis, although antibody-induced cell signaling and/or disturbances to the cytoskeleton may
30 also be involved.

As the naked or unconjugated anti-aminophospholipid antibodies or antibody fragments bind to aminophospholipids at the surface of the tumor vascular endothelial cells, they will form an antibody coating on the luminal surface. This may function to attract immune effector cells, such as cytotoxic T cells and/or NK cells, which will then exert a cell-mediated cytotoxic effect on the vascular endothelial cells.

Binding of intact anti-aminophospholipid antibodies to the vascular endothelial cell surface will also mean that the Fc portions of the antibodies will protrude into the vessel lumen. As antibody Fc fragments activate the complement pathway, the observed cellular destruction may be a result of complement-directed lysis. Antibody binding thus activates the complement-dependent coagulation cascade, causing multi-component complexes to assemble and, ultimately, to generate a lytic complex that permeabilizes the target cell. "Complement-activated ADCC" may also be operating in the destruction, in which complement binds to the antibody coated target cell, and in which cells, such as neutrophils, having receptors for complement, lyse the target cell.

Anti-aminophospholipid antibody binding may also induce apoptosis in the tumor vascular endothelial cells. Other groups have identified PS as a possible marker of apoptosis (Williamson and Schlegel, 1994). However, these previous studies were connected with the appearance of externalized PS after other stimuli had initiated the apoptotic event (Martin *et al.*, 1995), the inverse of the present apoptosis induction proposal. There are no known reports of antibody binding to PS actually inducing apoptosis. Still, the inventors consider this to be about as likely a mechanism as the cell-mediated cytotoxicity or complement-mediated lysis, despite the fact that tangential evidence to the contrary has very recently been published by others (Nakamura *et al.*, 1998).

Nakamura *et al.* (1998) analyzed antibody fractions from patients with lupus anticoagulant (LAC), a disorder associated with arterial and venous thrombosis, thrombocytopenia, and recurrent fetal loss. Plasma with LAC activity was initially reported to

induce apoptosis in endothelial cells (Nakamura *et al.*, 1994). The apoptotic activities of LAC antisera were then reported to be localized in an annexin V-binding antibody fraction in 10/10 patients studied (Nakamura *et al.*, 1998). As annexin binds to PS, the apparent ability of anti-annexin antibodies to induce apoptosis would be the opposite of one of the destructive mechanisms proposed by the present inventors, *i.e.*, the ability of anti-PS antibodies to induce apoptosis.

The ability of LAC antibody fractions to induce apoptosis was further reported to be inhibited by preincubation with annexin V (Nakamura *et al.*, 1998). In contrast, removal of anti-phospholipid antibodies from the patients' IgG fractions with phospholipid liposomes did not abolish the apoptosis-inducing activities or annexin V binding (Nakamura *et al.*, 1998). These results reasonably implied that patients with LAC often have antibodies that do not bind phospholipids and yet are responsible for the induction of apoptosis in endothelial cells (Nakamura *et al.*, 1998).

Without needing to equate the Nakamura *et al.* (1998) LAC data with the present observations from *in vivo* studies of tumors and tumor vasculature, due to the evidently disparate nature of these clinical conditions, the present inventors nonetheless have certain unifying theories. Nakamura *et al.* (1998) attempted to remove anti-phospholipid antibodies from patients' antisera using phospholipid liposomes, and observed that this did not abolish the apoptosis-inducing activity. These results led Nakamura *et al.* (1998) to conclude that the anti-phospholipids antibodies cannot be responsible for apoptotic activity. However, the present inventors now have the insight to suggest that the incubation with phospholipid liposomes may not have removed the anti-phospholipids antibodies from the antisera, as phospholipids are antigenically neutral in bilayer and liposomal form, and largely only bind antibodies in hexagonal form (Rauch *et al.*, 1986; Rauch and Janoff, 1990; Berard *et al.*, 1993; each incorporated herein by reference) or in association with membrane proteins. Thus, anti-phospholipids antibodies may remain in the LAC antisera and may cause, or contribute to, the observed apoptotic activity.

It is also conceivable that anti-aminophospholipid binding to the surface of tumor vascular endothelial cells may cause disturbances in the cytoskeletal organization of the cell. As the cytoskeleton plays a role in the organization of surface membranes, and as anti-aminophospholipid binding may disturb (or further disturb) the membrane, antibody binding
5 may transmit changes to cytoskeletal proteins that interact with the bilayer. It is already known that the spatial organization of cytoskeletal proteins controls membrane stability and cell shape, and it is possible that perturbation of some cytoskeletal equilibrium may have far-reaching consequences on cell integrity.

10 A further mechanism of operation of the invention may be that anti-aminophospholipid antibody binding to the endothelial cell surface may initiate signal transduction by, as yet, undefined pathways. Anti-aminophospholipid antibody binding may also disturb known signal transduction pathways, *e.g.*, by altering the conformation and/or interactions of membrane receptors, signal transduction proteins, membrane channels, and the like. Signals for cell
15 destruction (apoptosis) may be initiated or mimicked, and/or preservation/homeostatic signals may be inhibited.

Although of scientific interest, determining the exact nature of the vascular destruction achieved by the naked anti-aminophospholipid antibodies is not necessary to the practice of the present invention. Given that the administration of anti-aminophospholipid antibodies is herein
20 shown to advantageously result in specific anti-tumor effects *in vivo*, the invention can be utilized irrespective of the molecular mechanism that underlies this phenomenon.

The naked antibody use of the present invention thus represent a significant advance in
25 tumor therapy. Although coaguligands are advantageous for tumor therapy, the targeting antibody or ligand still needs to be conjugated to, or functionally associated with, the effector coagulant, such as Tissue Factor. Therefore, the practice of the coaguligand targeting and tumor destruction methodology is somewhat laborious in that it requires the preparation of suitable conjugates, or co-ordinated molecular complexes (including bispecific antibodies). For example,
30 one must prepare a targeting antibody or ligand that binds to the desired target antigen; choose

obtained, the immunized animal can be bled and the serum isolated and stored. The animal can also be used to generate monoclonal antibodies.

As is well known in the art, the immunogenicity of a particular composition can be enhanced by the use of non-specific stimulators of the immune response, known as adjuvants. Exemplary adjuvants include complete Freund's adjuvant, a non-specific stimulator of the immune response containing killed *Mycobacterium tuberculosis*; incomplete Freund's adjuvant; and aluminum hydroxide adjuvant.

It may also be desired to boost the host immune system, as may be achieved by associating aminophospholipids with, or coupling aminophospholipids to, a carrier. Exemplary carriers are keyhole limpet hemocyanin (KLH) and bovine serum albumin (BSA). Other albumins such as ovalbumin, mouse serum albumin or rabbit serum albumin can also be used as carriers.

As is also known in the art, a given composition may vary in its immunogenicity. However, the generation of antibodies against aminophospholipids is not particularly difficult. For example, highly specific anti-phosphatidylserine antibodies were raised in rabbits immunized by intramuscular injections of phosphatidylserine-containing polyacrylamide gels and with phosphatidylserine-cytochrome *c* vesicles (Maneta-Peyret *et al.*, 1988; 1989; each incorporated herein by reference). The use of acrylamide implants enhanced the production of antibodies (Maneta-Peyret *et al.*, 1988; 1989). The anti-phosphatidylserine antibodies raised in this manner are able to detect phosphatidylserine *in situ* on human platelets (Maneta-Peyret *et al.*, 1988). The groups of Inoue, Rote and Rauch have also developed anti-PS and anti-PE antibodies (see below).

E2. Monoclonal Anti-Aminophospholipid Antibodies

Various methods for generating monoclonal antibodies (MAbs) are also now very well known in the art. The most standard monoclonal antibody generation techniques generally begin along the same lines as those for preparing polyclonal antibodies (Antibodies: A Laboratory

Manual, Cold Spring Harbor Laboratory, 1988; incorporated herein by reference). A polyclonal antibody response is initiated by immunizing an animal with an immunogenic aminophospholipid composition and, when a desired titer level is obtained, the immunized animal can be used to generate MAbs.

5

MAbs may be readily prepared through use of well-known techniques, such as those exemplified in US. Patent 4,196,265, incorporated herein by reference. Typically, this technique involves immunizing a suitable animal with the selected aminophospholipid immunogen composition. The immunizing composition is administered in a manner effective to stimulate antibody producing cells. Rodents such as mice and rats are preferred animals, however, the use of rabbit, sheep and frog cells is also possible. The use of rats may provide certain advantages (Goding, 1986, pp. 60-61; incorporated herein by reference), but mice are preferred, with the BALB/c mouse being most preferred as this is most routinely used and generally gives a higher percentage of stable fusions.

15

Following immunization, somatic cells with the potential for producing aminophospholipid antibodies, specifically B lymphocytes (B cells), are selected for use in the MAb generating protocol. These cells may be obtained from biopsied spleens, tonsils or lymph nodes, or from a peripheral blood sample. Spleen cells and peripheral blood cells are preferred, the former because they are a rich source of antibody-producing cells that are in the dividing plasmablast stage, and the latter because peripheral blood is easily accessible. Often, a panel of animals will have been immunized and the spleen of animal with the highest antibody titer will be removed and the spleen lymphocytes obtained by homogenizing the spleen with a syringe. Typically, a spleen from an immunized mouse contains approximately 5×10^7 to 2×10^8 lymphocytes.

25

The anti-aminophospholipid antibody-producing B lymphocytes from the immunized animal are then fused with cells of an immortal myeloma cell, generally one of the same species as the animal that was immunized. Myeloma cell lines suited for use in hybridoma-producing fusion procedures preferably are non-antibody-producing, have high fusion efficiency, and

30

enzyme deficiencies that render them incapable of growing in certain selective media which support the growth of only the desired fused cells (hybridomas).

Any one of a number of myeloma cells may be used, as are known to those of skill in the art (Goding, pp. 65-66, 1986; Campbell, pp. 75-83, 1984; each incorporated herein by reference). For example, where the immunized animal is a mouse, one may use P3-X63/Ag8, X63-Ag8.653, NS1/1.Ag 4 1, Sp210-Ag14, FO, NSO/U, MPC-11, MPC11-X45-GTG 1.7 and S194/5XX0 Bul; for rats, one may use R210.RCY3, Y3-Ag 1.2.3, IR983F, 4B210 or one of the above listed mouse cell lines; and U-266, GM1500-GRG2, LICR-LON-HMy2 and UC729-6, are all useful in connection with human cell fusions.

Methods for generating hybrids of antibody-producing spleen or lymph node cells and myeloma cells usually comprise mixing somatic cells with myeloma cells in a 4:1 proportion, though the proportion may vary from about 20:1 to about 1:1, respectively, in the presence of an agent or agents (chemical or electrical) that promote the fusion of cell membranes. Fusion methods using Sendai virus have been described by Kohler and Milstein (1975; 1976; each incorporated herein by reference), and those using polyethylene glycol (PEG), such as 37% (v/v) PEG, by Gefter *et al.* (1977; incorporated herein by reference). The use of electrically induced fusion methods is also appropriate (Goding pp. 71-74, 1986; incorporated herein by reference).

Fusion procedures usually produce viable hybrids at low frequencies, about 1×10^{-6} to 1×10^{-8} . However, this does not pose a problem, as the viable, fused hybrids are differentiated from the parental, unfused cells (particularly the unfused myeloma cells that would normally continue to divide indefinitely) by culturing in a selective medium. The selective medium is generally one that contains an agent that blocks the *de novo* synthesis of nucleotides in the tissue culture media. Exemplary and preferred agents are aminopterin, methotrexate, and azaserine. Aminopterin and methotrexate block *de novo* synthesis of both purines and pyrimidines, whereas azaserine blocks only purine synthesis. Where aminopterin or methotrexate is used, the media is supplemented with hypoxanthine and thymidine as a source of nucleotides (HAT medium). Where azaserine is used, the media is supplemented with hypoxanthine.

The preferred selection medium is HAT. Only cells capable of operating nucleotide salvage pathways are able to survive in HAT medium. The myeloma cells are defective in key enzymes of the salvage pathway, *e.g.*, hypoxanthine phosphoribosyl transferase (HPRT), and they cannot survive. The B cells can operate this pathway, but they have a limited life span in culture and generally die within about two weeks. Therefore, the only cells that can survive in the selective media are those hybrids formed from myeloma and B cells.

This culturing provides a population of hybridomas from which specific hybridomas are selected. Typically, selection of hybridomas is performed by culturing the cells by single-clone dilution in microtiter plates, followed by testing the individual clonal supernatants (after about two to three weeks) for the desired anti-aminophospholipid reactivity. The assay should be sensitive, simple and rapid, such as radioimmunoassays, enzyme immunoassays, cytotoxicity assays, plaque assays, dot immunobinding assays, and the like.

The selected hybridomas would then be serially diluted and cloned into individual anti-aminophospholipid antibody-producing cell lines, which clones can then be propagated indefinitely to provide MAbs. The cell lines may be exploited for MAb production in two basic ways. A sample of the hybridoma can be injected (often into the peritoneal cavity) into a histocompatible animal of the type that was used to provide the somatic and myeloma cells for the original fusion. The injected animal develops tumors secreting the specific monoclonal antibody produced by the fused cell hybrid. The body fluids of the animal, such as serum or ascites fluid, can then be tapped to provide MAbs in high concentration. The individual cell lines could also be cultured *in vitro*, where the MAbs are naturally secreted into the culture medium from which they can be readily obtained in high concentrations.

MAbs produced by either means will generally be further purified, *e.g.*, using filtration, centrifugation and various chromatographic methods, such as HPLC or affinity chromatography, all of which purification techniques are well known to those of skill in the art. These purification techniques each involve fractionation to separate the desired antibody from other components of

a mixture. Analytical methods particularly suited to the preparation of antibodies include, for example, protein A-Sepharose and/or protein G-Sepharose chromatography.

Umeda *et al.* (1989; incorporated herein by reference) reported the effective production of monoclonal antibodies recognizing stereo-specific epitopes of phosphatidylserine. The Umeda system is based on the direct immunization of phosphatidylserine into mouse spleen using a *Salmonella*-coated aminophospholipid sample (Umeda *et al.*, 1989; incorporated herein by reference). The Umeda protocol gives a high frequency of anti-PS MAbs, which exhibit three distinct reactivity profiles ranging from highly specific to broadly cross-reactive. Umeda is therefore also incorporated herein by reference for purposes of further describing screening assays to identify MAbs that bind specifically to PS, *e.g.*, and do not bind to phosphatidylcholine.

Any of the 61 hybridomas generated by Umeda could potentially be employed in the present invention. Examples are PSC8, PSF11, PSG3, PSD11, PSF10, PS1B, PS3D12, PS2C11; PS3A, PSF6, PSF7, PSB4, PS3H1; PS4A7 and PS1G3. More preferred are PS3A, PSF6, PSF7, PSB4 and PS3H1 as they bind only to phosphatidylserine and phosphatidylethanolamine. Preferred anti-PS antibodies are PS4A7 (IgM) and PS1G3 (IgG₃), as they are highly specific for PS and exhibit no cross-reaction with other phospholipids. PS4A7 recognizes the stereo-specific configuration of the serine residue in PS (Figure 1 Umeda *et al.*, 1989; incorporated herein by reference).

Igarashi *et al.* (1991; incorporated herein by reference) also reported the effective induction of anti-PS antibodies of the IgG isotype by intrasplenic immunization. Only a slight increase of the titer was observed when the antigen was again injected intravenously. A high frequency of anti-PS MAbs of the IgG isotype was also observed even when MAbs were produced 10 days after the intrasplenic injection of the antigen. These antibodies were also employed by Schuurmans Stekhoven *et al.* (1994).

The other significant anti-PS antibody production has been by Rote and colleagues. Rote *et al.* (1993; incorporated herein by reference) particularly employed PS micelles in combination with Freund's complete adjuvant to generate specific anti-PS antibodies. Rote *et al.* (1993) also generated monoclonal antibodies that differentiate between cardiolipin (CL) and PS. Rote *et al.* (1993) is therefore also incorporated herein by reference for purposes of further describing screening assays to identify MAb that bind specifically to PS by testing against resting and thrombin-activated platelets using flow cytometry.

The 3SB9b antibody produced by Rote *et al.* (1993) reacted with only with PS, and is a preferred antibody for use in the present invention. BA3B5C4 may also be used as it reacts with both PS and CL. These antibodies are also described in Lin *et al.* (1995), Obringer *et al.* (1995) and Katsuragawa *et al.* (1997).

E3. Anti-Aminophospholipid Antibodies from Phagemid Libraries

Recombinant technology now allows the preparation of antibodies having the desired specificity from recombinant genes encoding a range of antibodies (Van Dijk *et al.*, 1989; incorporated herein by reference). Certain recombinant techniques involve the isolation of the antibody genes by immunological screening of combinatorial immunoglobulin phage expression libraries prepared from RNA isolated from the spleen of an immunized animal (Morrison *et al.*, 1986; Winter and Milstein, 1991; each incorporated herein by reference).

For such methods, combinatorial immunoglobulin phagemid libraries are prepared from RNA isolated from the spleen of the immunized animal, and phagemids expressing appropriate antibodies are selected by panning using cells expressing the antigen and control cells. The advantages of this approach over conventional hybridoma techniques are that approximately 10^4 times as many antibodies can be produced and screened in a single round, and that new specificities are generated by H and L chain combination, which further increases the percentage of appropriate antibodies generated.

One method for the generation of a large repertoire of diverse antibody molecules in bacteria utilizes the bacteriophage lambda as the vector (Huse *et al.*, 1989; incorporated herein by reference). Production of antibodies using the lambda vector involves the cloning of heavy and light chain populations of DNA sequences into separate starting vectors. The vectors are subsequently combined randomly to form a single vector that directs the co-expression of heavy and light chains to form antibody fragments. The heavy and light chain DNA sequences are obtained by amplification, preferably by PCR™ or a related amplification technique, of mRNA isolated from spleen cells (or hybridomas thereof) from an animal that has been immunized with a selected antigen. The heavy and light chain sequences are typically amplified using primers that incorporate restriction sites into the ends of the amplified DNA segment to facilitate cloning of the heavy and light chain segments into the starting vectors.

09351662-0712-90

Another method for the generation and screening of large libraries of wholly or partially synthetic antibody combining sites, or paratopes, utilizes display vectors derived from filamentous phage such as M13, f1 or fd. These filamentous phage display vectors, referred to as "phagemids", yield large libraries of monoclonal antibodies having diverse and novel immunospecificities. The technology uses a filamentous phage coat protein membrane anchor domain as a means for linking gene-product and gene during the assembly stage of filamentous phage replication, and has been used for the cloning and expression of antibodies from combinatorial libraries (Kang *et al.*, 1991; Barbas *et al.*, 1991; each incorporated herein by reference).

This general technique for filamentous phage display is described in U.S. Patent 5,658,727, incorporated herein by reference. In a most general sense, the method provides a system for the simultaneous cloning and screening of pre-selected ligand-binding specificities from antibody gene repertoires using a single vector system. Screening of isolated members of the library for a pre-selected ligand-binding capacity allows the correlation of the binding capacity of an expressed antibody molecule with a convenient means to isolate the gene that encodes the member from the library.

Linkage of expression and screening is accomplished by the combination of targeting of a fusion polypeptide into the periplasm of a bacterial cell to allow assembly of a functional antibody, and the targeting of a fusion polypeptide onto the coat of a filamentous phage particle during phage assembly to allow for convenient screening of the library member of interest. Periplasmic targeting is provided by the presence of a secretion signal domain in a fusion polypeptide. Targeting to a phage particle is provided by the presence of a filamentous phage coat protein membrane anchor domain (*i.e.*, a cpIII- or cpVIII-derived membrane anchor domain) in a fusion polypeptide.

10

The diversity of a filamentous phage-based combinatorial antibody library can be increased by shuffling of the heavy and light chain genes, by altering one or more of the complementarity determining regions of the cloned heavy chain genes of the library, or by introducing random mutations into the library by error-prone polymerase chain reactions. Additional methods for screening phagemid libraries are described in U.S. Patent No. 5,580,717; 5,427,908; 5,403,484; and 5,223,409, each incorporated herein by reference.

Another method for the screening of large combinatorial antibody libraries has been developed, utilizing expression of populations of diverse heavy and light chain sequences on the surface of a filamentous bacteriophage, such as M13, fl or fd (U.S. Patent No. 5,698,426; incorporated herein by reference). Two populations of diverse heavy (Hc) and light (Lc) chain sequences are synthesized by polymerase chain reaction (PCR™). These populations are cloned into separate M13-based vector containing elements necessary for expression. The heavy chain vector contains a gene VIII (gVIII) coat protein sequence so that translation of the heavy chain sequences produces gVIII-Hc fusion proteins. The populations of two vectors are randomly combined such that only the vector portions containing the Hc and Lc sequences are joined into a single circular vector.

The combined vector directs the co-expression of both Hc and Lc sequences for assembly of the two polypeptides and surface expression on M13 (U.S. Patent No. 5,698,426; incorporated

herein by reference). The combining step randomly brings together different Hc and Lc encoding sequences within two diverse populations into a single vector. The vector sequences donated from each independent vector are necessary for production of viable phage. Also, since the pseudo gVIII sequences are contained in only one of the two starting vectors, co-expression of functional antibody fragments as Lc associated gVIII-Hc fusion proteins cannot be accomplished on the phage surface until the vector sequences are linked in the single vector.

Surface expression of the antibody library is performed in an amber suppressor strain. An amber stop codon between the Hc sequence and the gVIII sequence unlinks the two components in a non-suppressor strain. Isolating the phage produced from the non-suppressor strain and infecting a suppressor strain will link the Hc sequences to the gVIII sequence during expression. Culturing the suppressor strain after infection allows the coexpression on the surface of M13 of all antibody species within the library as gVIII fusion proteins (gVIII-Fab fusion proteins). Alternatively, the DNA can be isolated from the non-suppressor strain and then introduced into a suppressor strain to accomplish the same effect.

The surface expression library is screened for specific Fab fragments that bind preselected molecules by standard affinity isolation procedures. Such methods include, for example, panning (Parmley and Smith, 1988; incorporated herein by reference), affinity chromatography and solid phase blotting procedures. Panning is preferred, because high titers of phage can be screened easily, quickly and in small volumes. Furthermore, this procedure can select minor Fab fragments species within the population, which otherwise would have been undetectable, and amplified to substantially homogenous populations. The selected Fab fragments can be characterized by sequencing the nucleic acids encoding the polypeptides after amplification of the phage population.

Another method for producing diverse libraries of antibodies and screening for desirable binding specificities is described in U.S. Patent No. 5,667,988 and U.S. Patent No. 5,759,817, each incorporated herein by reference. The method involves the preparation of libraries of heterodimeric immunoglobulin molecules in the form of phagemid libraries using degenerate

oligonucleotides and primer extension reactions to incorporate the degeneracies into the CDR regions of the immunoglobulin variable heavy and light chain variable domains, and display of the mutagenized polypeptides on the surface of the phagemid. Thereafter, the display protein is screened for the ability to bind to a preselected antigen.

5

The method for producing a heterodimeric immunoglobulin molecule generally involves (1) introducing a heavy or light chain V region-coding gene of interest into the phagemid display vector; (2) introducing a randomized binding site into the phagemid display protein vector by primer extension with an oligonucleotide containing regions of homology to a CDR of the antibody V region gene and containing regions of degeneracy for producing randomized coding sequences to form a large population of display vectors each capable of expressing different putative binding sites displayed on a phagemid surface display protein; (3) expressing the display protein and binding site on the surface of a filamentous phage particle; and (4) isolating (screening) the surface-expressed phage particle using affinity techniques such as panning of phage particles against a preselected antigen, thereby isolating one or more species of phagemid containing a display protein containing a binding site that binds a preselected antigen.

10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995

A further variation of this method for producing diverse libraries of antibodies and screening for desirable binding specificities is described in U.S. Patent No. 5,702,892, incorporated herein by reference. In this method, only heavy chain sequences are employed, the heavy chain sequences are randomized at all nucleotide positions which encode either the CDRI or CDRIII hypervariable region, and the genetic variability in the CDRs is generated independent of any biological process.

25 In the method, two libraries are engineered to genetically shuffle oligonucleotide motifs within the framework of the heavy chain gene structure. Through random mutation of either CDRI or CDRIII, the hypervariable regions of the heavy chain gene were reconstructed to result in a collection of highly diverse sequences. The heavy chain proteins encoded by the collection of mutated gene sequences possessed the potential to have all of the binding characteristics of an immunoglobulin while requiring only one of the two immunoglobulin chains.

30

Specifically, the method is practiced in the absence of the immunoglobulin light chain protein. A library of phage displaying modified heavy chain proteins is incubated with an immobilized ligand to select clones encoding recombinant proteins that specifically bind the immobilized ligand. The bound phage are then dissociated from the immobilized ligand and amplified by growth in bacterial host cells. Individual viral plaques, each expressing a different recombinant protein, are expanded, and individual clones can then be assayed for binding activity.

10 **E4. Anti-Aminophospholipid Antibodies from Human Patients**

Antibodies against aminophospholipids, particularly phosphatidylserine and phosphatidylethanolamine, occur in the human population, where they are correlated with certain disease states. Anti-aminophospholipid antibodies are part of the heterogeneous anti-phospholipid antibodies (aPL), observed to have families of different specificities and classes. Primary anti-phospholipid syndrome (APS) has even been separated from other forms of autoimmune disease associated with anti-phospholipid antibody production.

Anti-PS antibodies are particularly associated with recurrent pregnancy loss (Rote *et al.*, 1995; Rote, 1996; Vogt *et al.*, 1996; Vogt *et al.*, 1997; incorporated herein by reference) and with the autoimmune disease, systemic lupus erythematosus (SLE or "lupus") (Branch *et al.*, 1987; incorporated herein by reference). Anti-PE antibodies have also been reported in human patients, particularly those with autoimmune diseases (Staub *et al.*, 1989). Branch *et al.* (1987) reported that 80% of patients with lupus anticoagulant (LA or LAC) had autoantibodies that recognized PE; with Drouvalakis and Buchanan (1998) increasing this number to 95% PE-positives from autoimmune LAC sera.

Anti-phospholipid antibodies are not to be confused with anti-endothelial cell antibodies (AECA), although they can be found in the same patient. The existence of AECA has been documented in a variety of clinical settings associated with vasculitis, such as systemic sclerosis

(SS). To study AECA, antibodies are obtained from patients that do not have anti-phospholipid antibodies (aPL-negative sera).

The pathogenic role of AECA remains unclear, although Bordron *et al.* (1998) very recently suggested that AECA may initiate apoptosis in endothelial cells, which would be followed by PS transfer to the outer face of the membrane. They proposed that this would account for the subsequent generation of the anti-phospholipid antibodies that are sometimes seen in conjunction with AECA in patients with skin lesions or connective tissue disease (Bordron *et al.*, 1998). However, although AECA binding to an apoptosis-inducing antigen was postulated, these studies did not lead to the further characterization of AECA, still said to represent an extremely heterogeneous family of antibodies reacting with different (non-lipid) structures on endothelial cells (Bordron *et al.*, 1998).

Anti-phosphatidylserine antibodies are closely associated with pregnancy loss, pregnancy-induced hypertension and intrauterine growth retardation. A phosphatidylserine-dependent antigen has been shown to be expressed on the surface of a choriocarcinoma model (BeWo) of differentiating cytotrophoblastic cells, indicating that it should be accessible *in vivo* to circulating anti-phosphatidylserine antibodies (Rote *et al.*, 1995). Indeed, Vogt *et al.* (1996) showed that the monoclonal antibody 3SB9b, which reacts with phosphatidylserine but not cardiolipin, induced a significant reduction in both fetal and placental weights in a mouse model for the anti-phospholipid antibody syndrome

These authors developed a model for explaining miscarriages associated with anti-phospholipid antibodies: anti-phosphatidylserine antibody reveals sites for prothrombin binding on the surface of the trophoblast, most likely by removing Annexin V (Vogt *et al.*, 1997). Trophoblast differentiation is associated with externalization of phosphatidylserine from the inner to the outer surface of the plasma membrane. Normally, externalization of phosphatidylserine is concurrent with binding of Annexin V, which prevents the phosphatidylserine-rich surface from acting as a site for activation of coagulation. Thus, when

characteristic of autoantibodies from patients with autoimmune disease (Rauch and Janoff, 1990).

The hexagonal II phase form of aminophospholipids should thus be advantageously used
5 to generate antibodies for use in the present invention. Indeed, Trudell reported that antibodies
raised against TFA- (trifluoroacetyl-) protein adducts bind to TFA-phosphatidylethanolamine in
hexagonal phase phospholipid micelles, but not in lamellar liposomes (Trudell *et al.*, 1991a;
incorporated herein by reference). The authors suggested that TFA-phosphatidylethanolamine
adducts that reside in non-lamellar domains on the hepatocyte surface could be recognition sites
10 for anti-TFA-adduct antibodies and potentially participate in immune-mediated halothane
hepatotoxicity (Trudell *et al.*, 1991a). It was later shown that these same antibodies cross-react
with TFA-dioleoylphosphatidylethanolamine when this adduct is incorporated into the surface of
hepatocytes (Trudell *et al.*, 1991b; incorporated herein by reference), thus supporting this
hypothesis.

15 Berard further explained the hexagonal II phase form of aminophospholipids, such as PE
(Berard *et al.*, 1993; incorporated herein by reference). In bilayers, phospholipids generally
adopt a gel structure, crystalline lattice or lamellar phase (Berard *et al.*, 1993). However,
depending on the cholesterol content, protein and ionic environments, phospholipids can easily
20 change phases, adopting a hexagonal II phase (Berard *et al.*, 1993; incorporated herein by
reference). It is this hexagonal II phase of aminophospholipids that is believed to be
immunogenic, as initially proposed for autoantibody generation in disease situations (Berard *et al.*,
1993; incorporated herein by reference).

25 Qamar *et al.* (1990; incorporated herein by reference) have developed a variation on the
hexagonal aminophospholipid recognition theme. Using phosphatidylethanolamine as a model,
these authors reported that anti-PE antibodies from aPL-positive SLE sera do not bind to PE, but
in fact are directed to lysophosphatidylethanolamine (1PE), a natural PE degradation product and
a likely contaminant of most PE preparations (Qamar *et al.*, 1990; incorporated herein by
30 reference).

Other recent data indicate that most anti-phospholipid antibodies recognize phospholipid in the context of nearby proteins (Rote, 1996; Chamley *et al.*, 1991). In plasma membranes, the majority of the phospholipid appears to be naturally in non-antigenic bilaminar form (Rote, 1996). Accessory molecules may help facilitate the transition to hexagonal antigenic forms and stabilize their expression (Galli *et al.*, 1993). For example, naturally occurring anti-phospholipid antibodies were first reported to recognize complexes of cardiolipin or phosphatidylserine with β_2 -glycoprotein I (β_2 -GPI or apolipoprotein H, apoH) (Galli *et al.*, 1990; 1993). β_2 -GPI is believed to stabilize phospholipids in antigenic conformations that do not exist in pure phospholipids (McNeil *et al.*, 1990; U.S. Patent No. 5,344,758; Chamley *et al.*, 1991; Matsuura *et al.*, 1994). Prothrombin has also been implicated in the phospholipid stabilization process (Bervers *et al.*, 1991).

CONFIDENTIAL - 071200

Phospholipid-binding plasma proteins are also generally necessary for antibody recognition of the electrically neutral or zwitterionic phospholipid, phosphatidylethanolamine. Sugi and McIntyre (1995; incorporated herein by reference) identified two prominent PE-binding plasma proteins as high molecular weight kininogen (HMWK or HK) and low molecular weight kininogen (LMWK or LK). Anti-PE antibodies from patients with SLE and/or recurrent spontaneous abortions were shown not to recognize PE, HMWK or LMWK when they were presented independently as sole antigens on ELISA plates (Sugi and McIntyre, 1995). Other anti-PE-positive sera that did not react with PE-HMWK or PE-LMWK were suggested to recognize factor XI or prekallikrein, which normally bind to HMWK (Sugi and McIntyre, 1995; incorporated herein by reference).

The validity of these results was confirmed by showing that intact HMWK binds to various phospholipids, such as cardiolipin, phosphatidylserine, phosphatidylcholine and phosphatidylethanolamine; but that anti-PE antibodies recognize only a kininogen-PE complex, and do not recognize kininogens presented with other phospholipid substrates (Sugi and McIntyre, 1996a; incorporated herein by reference). This indicates that PE induces unique

antigenic conformational changes in the kininogens that are not induced when the kininogens bind to other phospholipids (Sugi and McIntyre, 1996a).

It has further been suggested that kininogens can bind to platelets by virtue of exposed
5 PE in the platelet membrane (Sugi and McIntyre, 1996b; incorporated herein by reference).
Exogenously added kininogen-dependent anti-PE was shown to increase thrombin-induced
platelet aggregation *in vitro*, but not to alter ADP-induced aggregation (Sugi and McIntyre,
1996b; incorporated herein by reference). In contrast, kininogen independent anti-PE, which
recognized PE *per se*, was reported not augment thrombin-induced platelet aggregation. It was
10 thus proposed that kininogen dependent anti-PE may disrupt the normal anti-thrombotic effects
of kininogen (Sugi and McIntyre, 1996b; incorporated herein by reference).

Anti-aminophospholipid antibodies from human patients are therefore a mixture of
antibodies that generally recognize aminophospholipids stabilized by protein interactions (Rote,
5 1996). The antibodies may bind to stabilized phospholipid epitopes, or may bind to an epitope
formed from the interaction of the phospholipid and amino acids on the stabilizing protein
(Rote, 1996). Either way, such antibodies clearly recognize aminophospholipids in natural
membranes in the human body, probably associated with plasma proteins (McNeil *et al.*, 1990;
Bever *et al.*, 1991). These antibodies would thus be appropriate as starting materials for
15 generating an antibody for use in the present invention.

To prepare an anti-aminophospholipid antibody from a human patient, one would simply
obtain human lymphocytes from an individual having anti-aminophospholipid antibodies, for
example from human peripheral blood, spleen, lymph nodes, tonsils or the like, utilizing
25 techniques that are well known to those of skill in the art. The use of peripheral blood
lymphocytes will often be preferred.

Human monoclonal antibodies may be obtained from the human lymphocytes producing
the desired anti-aminophospholipid antibodies by immortalizing the human lymphocytes,
30 generally in the same manner as described above for generating any monoclonal antibody. The

reactivities of the antibodies in the culture supernatants are generally first checked, employing one or more selected aminophospholipid antigen(s), and the lymphocytes that exhibit high reactivity are grown. The resulting lymphocytes are then fused with a parent line of human or mouse origin, and further selection gives the optimal clones.

5

The recovery of monoclonal antibodies from the immortalized cells may be achieved by any method generally employed in the production of monoclonal antibodies. For instance, the desired monoclonal antibody may be obtained by cloning the immortalized lymphocyte by the limiting dilution method or the like, selecting the cell producing the desired antibody, growing
10 the selected cells in a medium or the abdominal cavity of an animal, and recovering the desired monoclonal antibody from the culture supernatant or ascites.

Such techniques have been used, for example, to isolate human monoclonal antibodies to *Pseudomonas aeruginosa* epitopes (U.S. Patent No. 5,196,337 and 5,252,480, each incorporated
15 herein by reference); polyribosylribitol phosphate capsular polysaccharides (U.S. Patent No. 4,954,449, incorporated herein by reference); the Rh(D) antigen (U.S. Patent No. 5,665,356, incorporated herein by reference); and viruses, such as human immunodeficiency virus, respiratory syncytial virus, herpes simplex virus, varicella zoster virus and cytomegalovirus (U.S. Patent No. 5,652,138; 5,762,905; and 4,950,595, each incorporated herein by reference).

The applicability of the foregoing techniques to the generation of human anti-aminophospholipid antibodies is clear. Rauch *et al.* (1986; incorporated herein by reference) generally used such methods to produce hybridomas by fusing lymphocytes from SLE patients with a lymphoblastoid line. This produced human antibodies that bound to hexagonal phase
20 phospholipids, including natural and synthetic forms of phosphatidylethanolamine (Rauch *et al.*, 1986; incorporated herein by reference).

Additionally, the methods described in U.S. Patent 5,648,077 (incorporated herein by reference) can be used to form a trioma or a quadroma that produces a human antibody against a
30 selected aminophospholipid. In a general sense, a hybridoma cell line comprising a parent

rodent immortalizing cell, such as a murine myeloma cell, *e.g.* SP-2, is fused to a human partner cell, resulting in an immortalizing xenogeneic hybridoma cell. This xenogeneic hybridoma cell is fused to a cell capable of producing an anti-aminophospholipid human antibody, resulting in a trioma cell line capable of generating human antibody effective against such antigen in a human.

- 5 Alternately, when greater stability is desired, a trioma cell line which preferably no longer has the capability of producing its own antibody is made, and this trioma is then fused with a further cell capable of producing an antibody useful against the aminophospholipid antigen to obtain a still more stable hybridoma (quadroma) that produces antibody against the antigen.

10 E5. Anti-Aminophospholipid Antibodies from Human Lymphocytes

In vitro immunization, or antigen stimulation, may also be used to generate a human anti-aminophospholipid antibody. Such techniques can be used to stimulate peripheral blood lymphocytes from both anti-aminophospholipid antibody-producing human patients, and also from normal, healthy subjects. Indeed, Vlachoyiannopoulos *et al.* (1992; incorporated herein by
15 reference) reported that low titer anti-aminophospholipid antibodies occur in normal subjects. Even if this were not the case, anti-aminophospholipid antibodies can be prepared from healthy human subjects, simply by stimulating antibody-producing cells with aminophospholipids *in vitro*.

20 Such "*in vitro* immunization" involves antigen-specific activation of non-immunized B lymphocytes, generally within a mixed population of lymphocytes (mixed lymphocyte cultures, MLC). *In vitro* immunizations may also be supported by B cell growth and differentiation factors and lymphokines. The antibodies produced by these methods are often IgM antibodies (Borrebaeck *et al.*, 1986; incorporated herein by reference).

25

Another method has been described (U.S. Patent No. 5,681,729, incorporated herein by reference) wherein human lymphocytes that mainly produce IgG (or IgA) antibodies can be obtained. The method involves, in a general sense, transplanting human lymphocytes to an immunodeficient animal so that the human lymphocytes "take" in the animal body; immunizing
30 the animal with a desired antigen, so as to generate human lymphocytes producing an antibody

specific to the antigen; and recovering the human lymphocytes producing the antibody from the animal. The human lymphocytes thus produced can be used to produce a monoclonal antibody by immortalizing the human lymphocytes producing the antibody, cloning the obtained immortalized human-originated lymphocytes producing the antibody, and recovering a
5 monoclonal antibody specific to the desired antigen from the cloned immortalized human-originated lymphocytes.

The immunodeficient animals that may be employed in this technique are those that do not exhibit rejection when human lymphocytes are transplanted to the animals. Such animals
10 may be artificially prepared by physical, chemical or biological treatments. Any immunodeficient animal may be employed. The human lymphocytes may be obtained from human peripheral blood, spleen, lymph nodes, tonsils or the like.

The "taking" of the transplanted human lymphocytes in the animals can be attained by
15 merely administering the human lymphocytes to the animals. The administration route is not restricted and may be, for example, subcutaneous, intravenous or intraperitoneal. The dose of the human lymphocytes is not restricted, and can usually be 10^6 to 10^8 lymphocytes per animal. The immunodeficient animal is then immunized with the desired aminophospholipid antigen.

After the immunization, human lymphocytes are recovered from the blood, spleen, lymph
20 nodes or other lymphatic tissues by any conventional method. For example, mononuclear cells can be separated by the Ficoll-Hypaque (specific gravity: 1.077) centrifugation method, and the monocytes removed by the plastic dish adsorption method. The contaminating cells originating from the immunodeficient animal may be removed by using an antiserum specific to the animal
25 cells. The antiserum may be obtained by, for example, immunizing a second, distinct animal with the spleen cells of the immunodeficient animal, and recovering serum from the distinct immunized animal. The treatment with the antiserum may be carried out at any stage. The human lymphocytes may also be recovered by an immunological method employing a human immunoglobulin expressed on the cell surface as a marker.

genetic material may be introduced to the host in conventional manner, for example by injection or other procedures into fertilized eggs or embryonic stem cells.

In preferred aspects, a host animal that initially does not carry genetic material encoding immunoglobulin constant regions is utilized, so that the resulting transgenic animal will use only the inserted human genetic material when producing immunoglobulins. This can be achieved either by using a naturally occurring mutant host lacking the relevant genetic material, or by artificially making mutants *e.g.*, in cell lines ultimately to create a host from which the relevant genetic material has been removed.

10

Where the host animal carries genetic material encoding immunoglobulin constant regions, the transgenic animal will carry the naturally occurring genetic material and the inserted genetic material and will produce immunoglobulins derived from the naturally occurring genetic material, the inserted genetic material, and mixtures of both types of genetic material. In this case the desired immunoglobulin can be obtained by screening hybridomas derived from the transgenic animal, *e.g.*, by exploiting the phenomenon of allelic exclusion of antibody gene expression or differential chromosome loss.

15

20

Once a suitable transgenic animal has been prepared, the animal is simply immunized with the desired immunogen. Depending on the nature of the inserted material, the animal may produce a chimeric immunoglobulin, *e.g.* of mixed mouse/human origin, where the genetic material of foreign origin encodes only part of the immunoglobulin; or the animal may produce an entirely foreign immunoglobulin, *e.g.* of wholly human origin, where the genetic material of foreign origin encodes an entire immunoglobulin.

25

Polyclonal antisera may be produced from the transgenic animal following immunization. Immunoglobulin-producing cells may be removed from the animal to produce the immunoglobulin of interest. Preferably, monoclonal antibodies are produced from the transgenic animal, *e.g.*, by fusing spleen cells from the animal with myeloma cells and screening the

resulting hybridomas to select those producing the desired antibody. Suitable techniques for such processes are described herein.

In an alternative approach, the genetic material may be incorporated in the animal in such a way that the desired antibody is produced in body fluids such as serum or external secretions of the animal, such as milk, colostrum or saliva. For example, by inserting *in vitro* genetic material encoding for at least part of a human immunoglobulin into a gene of a mammal coding for a milk protein and then introducing the gene to a fertilized egg of the mammal, *e.g.*, by injection, the egg may develop into an adult female mammal producing milk containing immunoglobulin derived at least in part from the inserted human immunoglobulin genetic material. The desired antibody can then be harvested from the milk. Suitable techniques for carrying out such processes are known to those skilled in the art.

5
10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995

The foregoing transgenic animals are usually employed to produce human antibodies of a single isotype, more specifically an isotype that is essential for B cell maturation, such as IgM and possibly IgD. Another preferred method for producing human anti-aminophospholipid antibodies is described in U.S. Patent No. 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016; and 5,770,429; each incorporated by reference, wherein transgenic animals are described that are capable of switching from an isotype needed for B cell development to other isotypes.

In the development of a B lymphocyte, the cell initially produces IgM with a binding specificity determined by the productively rearranged V_H and V_L regions. Subsequently, each B cell and its progeny cells synthesize antibodies with the same L and H chain V regions, but they may switch the isotype of the H chain. The use of mu or delta constant regions is largely determined by alternate splicing, permitting IgM and IgD to be coexpressed in a single cell. The other heavy chain isotypes (gamma, alpha, and epsilon) are only expressed natively after a gene rearrangement event deletes the C mu and C delta exons. This gene rearrangement process, termed isotype switching, typically occurs by recombination between so called switch segments

located immediately upstream of each heavy chain gene (except delta). The individual switch segments are between 2 and 10 kb in length, and consist primarily of short repeated sequences.

For these reasons, it is preferable that transgenes incorporate transcriptional regulatory sequences within about 1-2 kb upstream of each switch region that is to be utilized for isotype switching. These transcriptional regulatory sequences preferably include a promoter and an enhancer element, and more preferably include the 5' flanking (*i.e.*, upstream) region that is naturally associated (*i.e.*, occurs in germline configuration) with a switch region. Although a 5' flanking sequence from one switch region can be operably linked to a different switch region for transgene construction, in some embodiments it is preferred that each switch region incorporated in the transgene construct have the 5' flanking region that occurs immediately upstream in the naturally occurring germline configuration. Sequence information relating to immunoglobulin switch region sequences is known (Mills *et al.*, 1990; Sideras *et al.*, 1989; each incorporated herein by reference).

In the method described in U.S. Patent No. 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016; and 5,770,429, the human immunoglobulin transgenes contained within the transgenic animal function correctly throughout the pathway of B-cell development, leading to isotype switching. Accordingly, in this method, these transgenes are constructed so as to produce isotype switching and one or more of the following: (1) high level and cell-type specific expression, (2) functional gene rearrangement, (3) activation of and response to allelic exclusion, (4) expression of a sufficient primary repertoire, (5) signal transduction, (6) somatic hypermutation, and (7) domination of the transgene antibody locus during the immune response.

An important requirement for transgene function is the generation of a primary antibody repertoire that is diverse enough to trigger a secondary immune response for a wide range of antigens. The rearranged heavy chain gene consists of a signal peptide exon, a variable region exon and a tandem array of multi-domain constant region regions, each of which is encoded by several exons. Each of the constant region genes encode the constant portion of a different class of immunoglobulins. During B-cell development, V region proximal constant regions are

deleted leading to the expression of new heavy chain classes. For each heavy chain class, alternative patterns of RNA splicing give rise to both transmembrane and secreted immunoglobulins.

5 The human heavy chain locus consists of approximately 200 V gene segments spanning 2 Mb, approximately 30 D gene segments spanning about 40 kb, six J segments clustered within a 3 kb span, and nine constant region gene segments spread out over approximately 300 kb. The entire locus spans approximately 2.5 Mb of the distal portion of the long arm of chromosome 14. Heavy chain transgene fragments containing members of all six of the known V_H families, the D and J gene segments, as well as the mu, delta, gamma 3, gamma 1 and alpha 1 constant regions are known (Berman *et al.*, 1988; incorporated herein by reference). Genomic fragments containing all of the necessary gene segments and regulatory sequences from a human light chain locus is similarly constructed.

10 The expression of successfully rearranged immunoglobulin heavy and light transgenes usually has a dominant effect by suppressing the rearrangement of the endogenous immunoglobulin genes in the transgenic nonhuman animal. However, in certain embodiments, it is desirable to effect complete inactivation of the endogenous Ig loci so that hybrid immunoglobulin chains comprising a human variable region and a non-human (*e.g.*, murine) constant region cannot be formed, for example by trans-switching between the transgene and endogenous Ig sequences. Using embryonic stem cell technology and homologous recombination, the endogenous immunoglobulin repertoire can be readily eliminated. In addition, suppression of endogenous Ig genes may be accomplished using a variety of techniques, such as antisense technology.

25

In other aspects of the invention, it may be desirable to produce a trans-switched immunoglobulin. Antibodies comprising such chimeric trans-switched immunoglobulins can be used for a variety of applications where it is desirable to have a non-human (*e.g.*, murine) constant region, *e.g.*, for retention of effector functions in the host. The presence of a murine constant region can afford advantages over a human constant region, for example, to provide

30

murine effector functions (*e.g.*, ADCC, murine complement fixation) so that such a chimeric antibody may be tested in a mouse disease model. Subsequent to the animal testing, the human variable region encoding sequence may be isolated, *e.g.*, by PCR amplification or cDNA cloning from the source (hybridoma clone), and spliced to a sequence encoding a desired human constant region to encode a human sequence antibody more suitable for human therapeutic use.

E7. Humanized Anti-Aminophospholipid Antibodies

Human antibodies generally have at least three potential advantages for use in human therapy. First, because the effector portion is human, it may interact better with the other parts of the human immune system, *e.g.*, to destroy target cells more efficiently by complement-dependent cytotoxicity (CDC) or antibody-dependent cellular cytotoxicity (ADCC). Second, the human immune system should not recognize the antibody as foreign. Third, the half-life in the human circulation will be similar to naturally occurring human antibodies, allowing smaller and less frequent doses to be given.

Various methods for preparing human anti-aminophospholipids are provided herein. In addition to human antibodies, "humanized" antibodies have many advantages. "Humanized" antibodies are generally chimeric or mutant monoclonal antibodies from mouse, rat, hamster, rabbit or other species, bearing human constant and/or variable region domains or specific changes. Techniques for generating a so-called "humanized" anti-aminophospholipid antibody are well known to those of skill in the art.

Humanized antibodies also share the foregoing advantages. First, the effector portion is still human. Second, the human immune system should not recognize the framework or constant region as foreign, and therefore the antibody response against such an injected antibody should be less than against a totally foreign mouse antibody. Third, injected humanized antibodies, as opposed to injected mouse antibodies, will presumably have a half-life more similar to naturally occurring human antibodies, also allowing smaller and less frequent doses.

A number of methods have been described to produce humanized antibodies. Controlled rearrangement of antibody domains joined through protein disulfide bonds to form new, artificial protein molecules or "chimeric" antibodies can be utilized (Konieczny *et al.*, 1981; incorporated herein by reference). Recombinant DNA technology can also be used to construct gene fusions between DNA sequences encoding mouse antibody variable light and heavy chain domains and human antibody light and heavy chain constant domains (Morrison *et al.*, 1984; incorporated herein by reference).

DNA sequences encoding the antigen binding portions or complementarity determining regions (CDR's) of murine monoclonal antibodies can be grafted by molecular means into the DNA sequences encoding the frameworks of human antibody heavy and light chains (Jones *et al.*, 1986; Riechmann *et al.*, 1988; each incorporated herein by reference). The expressed recombinant products are called "reshaped" or humanized antibodies, and comprise the framework of a human antibody light or heavy chain and the antigen recognition portions, CDR's, of a murine monoclonal antibody.

Another method for producing humanized antibodies is described in U.S. Patent No. 5,639,641, incorporated herein by reference. The method provides, via resurfacing, humanized rodent antibodies that have improved therapeutic efficacy due to the presentation of a human surface in the variable region. In the method: (1) position alignments of a pool of antibody heavy and light chain variable regions is generated to give a set of heavy and light chain variable region framework surface exposed positions, wherein the alignment positions for all variable regions are at least about 98% identical; (2) a set of heavy and light chain variable region framework surface exposed amino acid residues is defined for a rodent antibody (or fragment thereof); (3) a set of heavy and light chain variable region framework surface exposed amino acid residues that is most closely identical to the set of rodent surface exposed amino acid residues is identified; (4) the set of heavy and light chain variable region framework surface exposed amino acid residues defined in step (2) is substituted with the set of heavy and light chain variable region framework surface exposed amino acid residues identified in step (3), except for those amino

acid residues that are within 5Å of any atom of any residue of the complementarity determining regions of the rodent antibody; and (5) the humanized rodent antibody having binding specificity is produced.

5 A similar method for the production of humanized antibodies is described in U.S. Patent No. 5,693,762; 5,693,761; 5,585,089; and 5,530,101, each incorporated herein by reference. These methods involve producing humanized immunoglobulins having one or more complementarity determining regions (CDR's) and possible additional amino acids from a donor immunoglobulin and a framework region from an accepting human immunoglobulin. Each
10 humanized immunoglobulin chain usually comprises, in addition to the CDR's, amino acids from the donor immunoglobulin framework that are capable of interacting with the CDR's to effect binding affinity, such as one or more amino acids that are immediately adjacent to a CDR in the donor immunoglobulin or those within about 3Å as predicted by molecular modeling. The heavy and light chains may each be designed by using any one, any combination, or all of the various position criteria described in U.S. Patent No. 5,693,762; 5,693,761; 5,585,089; and 5,530,101,
15 each incorporated herein by reference. When combined into an intact antibody, the humanized immunoglobulins are substantially non-immunogenic in humans and retain substantially the same affinity as the donor immunoglobulin to the original antigen.

20 An additional method for producing humanized antibodies is described in U.S. Patent 5,565,332 and 5,733,743, each incorporated herein by reference. This method combines the concept of humanizing antibodies with the phagemid libraries also described in detail herein. In a general sense, the method utilizes sequences from the antigen binding site of an antibody or population of antibodies directed against an antigen of interest. Thus for a single rodent
25 antibody, sequences comprising part of the antigen binding site of the antibody may be combined with diverse repertoires of sequences of human antibodies that can, in combination, create a complete antigen binding site.

The antigen binding sites created by this process differ from those created by CDR
30 grafting, in that only the portion of sequence of the original rodent antibody is likely to make

contacts with antigen in a similar manner. The selected human sequences are likely to differ in sequence and make alternative contacts with the antigen from those of the original binding site. However, the constraints imposed by binding of the portion of original sequence to antigen and the shapes of the antigen and its antigen binding sites, are likely to drive the new contacts of the human sequences to the same region or epitope of the antigen. This process has therefore been termed "epitope imprinted selection" (EIS).

Starting with an animal antibody, one process results in the selection of antibodies that are partly human antibodies. Such antibodies may be sufficiently similar in sequence to human antibodies to be used directly in therapy or after alteration of a few key residues. Sequence differences between the rodent component of the selected antibody with human sequences could be minimized by replacing those residues that differ with the residues of human sequences, for example, by site directed mutagenesis of individual residues, or by CDR grafting of entire loops. However, antibodies with entirely human sequences can also be created. EIS therefore offers a method for making partly human or entirely human antibodies that bind to the same epitope as animal or partly human antibodies respectively. In EIS, repertoires of antibody fragments can be displayed on the surface of filamentous phage and the genes encoding fragments with antigen binding activities selected by binding of the phage to antigen.

Additional methods for humanizing antibodies contemplated for use in the present invention are described in U.S. Patent No. 5,750,078; 5,502,167; 5,705,154; 5,770,403; 5,698,417; 5,693,493; 5,558,864; 4,935,496; and 4,816,567, each incorporated herein by reference.

E8. Mutagenesis by PCR

Site-specific mutagenesis is a technique useful in the preparation of individual antibodies through specific mutagenesis of the underlying DNA. The technique further provides a ready
5 ability to prepare and test sequence variants, incorporating one or more of the foregoing considerations, whether humanizing or not, by introducing one or more nucleotide sequence changes into the DNA.

Although many methods are suitable for use in mutagenesis, the use of the polymerase
10 chain reaction (PCR™) is generally now preferred. This technology offers a quick and efficient method for introducing desired mutations into a given DNA sequence. The following text particularly describes the use of PCR™ to introduce point mutations into a sequence, as may be used to change the amino acid encoded by the given sequence. Adaptations of this method are also suitable for introducing restriction enzyme sites into a DNA molecule.

In this method, synthetic oligonucleotides are designed to incorporate a point mutation at one end of an amplified segment. Following PCR™, the amplified fragments are blunt-ended by treating with Klenow fragments, and the blunt-ended fragments are then ligated and subcloned into a vector to facilitate sequence analysis.

To prepare the template DNA that one desires to mutagenize, the DNA is subcloned into a high copy number vector, such as pUC19, using restriction sites flanking the area to be mutated. Template DNA is then prepared using a plasmid miniprep. Appropriate oligonucleotide primers that are based upon the parent sequence, but which contain the desired
25 point mutation and which are flanked at the 5' end by a restriction enzyme site, are synthesized using an automated synthesizer. It is generally required that the primer be homologous to the template DNA for about 15 bases or so. Primers may be purified by denaturing polyacrylamide gel electrophoresis, although this is not absolutely necessary for use in PCR™. The 5' end of the oligonucleotides should then be phosphorylated.

30

The template DNA should be amplified by PCR™, using the oligonucleotide primers that contain the desired point mutations. The concentration of MgCl₂ in the amplification buffer will generally be about 15 mM. Generally about 20-25 cycles of PCR™ should be carried out as follows: denaturation, 35 sec. at 95°C; hybridization, 2 min. at 50°C; and extension, 2 min. at 72°C. The PCR™ will generally include a last cycle extension of about 10 min. at 72°C. After the final extension step, about 5 units of Klenow fragments should be added to the reaction mixture and incubated for a further 15 min. at about 30°C. The exonuclease activity of the Klenow fragments is required to make the ends flush and suitable for blunt-end cloning.

10 The resultant reaction mixture should generally be analyzed by nondenaturing agarose or acrylamide gel electrophoresis to verify that the amplification has yielded the predicted product. One would then process the reaction mixture by removing most of the mineral oils, extracting with chloroform to remove the remaining oil, extracting with buffered phenol and then concentrating by precipitation with 100% ethanol. Next, one should digest about half of the amplified fragments with a restriction enzyme that cuts at the flanking sequences used in the oligonucleotides. The digested fragments are purified on a low gelling/melting agarose gel.

15
20 To subclone the fragments and to check the point mutation, one would subclone the two amplified fragments into an appropriately digested vector by blunt-end ligation. This would be used to transform *E. coli*, from which plasmid DNA could subsequently be prepared using a miniprep. The amplified portion of the plasmid DNA would then be analyzed by DNA sequencing to confirm that the correct point mutation was generated. This is important as Taq DNA polymerase can introduce additional mutations into DNA fragments.

25 The introduction of a point mutation can also be effected using sequential PCR™ steps. In this procedure, the two fragments encompassing the mutation are annealed with each other and extended by mutually primed synthesis. This fragment is then amplified by a second PCR™ step, thereby avoiding the blunt-end ligation required in the above protocol. In this method, the preparation of the template DNA, the generation of the oligonucleotide primers and the first PCR™ amplification are performed as described above. In this process, however, the chosen

30

oligonucleotides should be homologous to the template DNA for a stretch of between about 15 and about 20 bases and must also overlap with each other by about 10 bases or more.

5 In the second PCR™ amplification, one would use each amplified fragment and each flanking sequence primer and carry PCR™ for between about 20 and about 25 cycles, using the conditions as described above. One would again subclone the fragments and check that the point mutation was correct by using the steps outlined above.

10 In using either of the foregoing methods, it is generally preferred to introduce the mutation by amplifying as small a fragment as possible. Of course, parameters such as the melting temperature of the oligonucleotide, as will generally be influenced by the GC content and the length of the oligo, should also be carefully considered. The execution of these methods, and their optimization if necessary, will be known to those of skill in the art, and are further described in various publications, such as *Current Protocols in Molecular Biology*, 1995, incorporated herein by reference.

5 When performing site-specific mutagenesis, **Table A** can be employed as a reference.

THE UNIVERSITY OF CHICAGO

95

E9. Recombinant Expression and Delivery

Given that many methods are available for cloning antibodies, anti-aminophospholipid antibodies may be prepared by routine methods of recombinant expression. The term "expression construct" is meant to include any type of genetic construct containing a nucleic acid coding for a gene product in which part or all of the nucleic acid encoding sequence is capable of being transcribed. The transcript will generally be translated into a protein. Thus, as intended herein, expression preferably includes both transcription of an anti-aminophospholipid antibody gene or DNA and translation of the mRNA into an anti-aminophospholipid antibody protein product.

For the expression of an anti-aminophospholipid antibody, once a suitable clone or clones have been obtained, whether they be cDNA based or genomic, one may proceed to prepare an expression system or "construct" for recombinant antibody production. The engineering of DNA segment(s) for expression in a prokaryotic or eukaryotic system will be performed by techniques generally known to those of skill in recombinant antibody expression (Sambrook *et al.*, 1989; incorporated herein by reference). In order for the construct to effect expression of an anti-aminophospholipid antibody transcript, the polynucleotide encoding the antibody will preferably be under the transcriptional control of a promoter that promotes expression in the chosen host cell.

Recombinantly produced antibodies may be purified and formulated for human administration. Alternatively, nucleic acids encoding anti-aminophospholipid antibodies may be delivered via gene therapy. Although naked recombinant DNA or plasmids may be employed, the use of liposomes or vectors is preferred. The ability of certain viruses to enter cells via receptor-mediated endocytosis and to integrate into host cell genome and express viral genes stably and efficiently have made them attractive candidates for the transfer of foreign genes into mammalian cells. Preferred gene therapy vectors for use in the present invention will generally be viral vectors.

Retroviruses have promise as gene delivery vectors due to their ability to integrate their genes into the host genome, transferring a large amount of foreign genetic material, infecting a broad spectrum of species and cell types and of being packaged in special cell-lines. Other viruses, such as adenovirus, herpes simplex viruses (HSV), cytomegalovirus (CMV), and adeno-associated virus (AAV), such as those described by U.S. Patent 5,139,941 (incorporated herein by reference), may also be engineered to serve as vectors for gene transfer.

Although some viruses that can accept foreign genetic material are limited in the number of nucleotides they can accommodate and in the range of cells they infect, these viruses have been demonstrated to successfully effect gene expression. However, adenoviruses do not integrate their genetic material into the host genome and therefore do not require host replication for gene expression, making them ideally suited for rapid, efficient, heterologous gene expression. Techniques for preparing replication-defective infective viruses are well known in the art.

In certain further embodiments, the gene therapy vector will be HSV. A factor that makes HSV an attractive vector is the size and organization of the genome. Because HSV is large, incorporation of multiple genes or expression cassettes is less problematic than in other smaller viral systems. In addition, the availability of different viral control sequences with varying performance (*e.g.*, temporal, strength) makes it possible to control expression to a greater extent than in other systems. It also is an advantage that the virus has relatively few spliced messages, further easing genetic manipulations. HSV also is relatively easy to manipulate and can be grown to high titers.

Of course, in using viral delivery systems, one will desire to purify the virion sufficiently to render it essentially free of undesirable contaminants, such as defective interfering viral particles or endotoxins and other pyrogens such that it will not cause any untoward reactions in the cell, animal or individual receiving the vector construct. A preferred means of purifying the vector involves the use of buoyant density gradients, such as cesium chloride gradient centrifugation.

E10. Antibody Fragments

Irrespective of the source of the original anti-aminophospholipid antibody, either the intact antibody, antibody multimers, or any one of a variety of functional, antigen-binding regions of the antibody may be used in the present invention. Exemplary functional regions include scFv, Fv, Fab', Fab and F(ab')₂ fragments of the anti-aminophospholipid antibodies. Techniques for preparing such constructs are well known to those in the art and are further exemplified herein.

The choice of antibody construct may be influenced by various factors. For example, prolonged half-life can result from the active readsorption of intact antibodies within the kidney, a property of the Fc piece of immunoglobulin. IgG based antibodies, therefore, are expected to exhibit slower blood clearance than their Fab' counterparts. However, Fab' fragment-based compositions will generally exhibit better tissue penetrating capability.

Fab fragments can be obtained by proteolysis of the whole immunoglobulin by the non-specific thiol protease, papain. Papain must first be activated by reducing the sulphhydryl group in the active site with cysteine, 2-mercaptoethanol or dithiothreitol. Heavy metals in the stock enzyme should be removed by chelation with EDTA (2 mM) to ensure maximum enzyme activity. Enzyme and substrate are normally mixed together in the ratio of 1:100 by weight. After incubation, the reaction can be stopped by irreversible alkylation of the thiol group with iodoacetamide or simply by dialysis. The completeness of the digestion should be monitored by SDS-PAGE and the various fractions separated by protein A-Sepharose or ion exchange chromatography.

25

The usual procedure for preparation of F(ab')₂ fragments from IgG of rabbit and human origin is limited proteolysis by the enzyme pepsin. The conditions, 100x antibody excess w/w in acetate buffer at pH 4.5, 37°C, suggest that antibody is cleaved at the C-terminal side of the inter-heavy-chain disulfide bond. Rates of digestion of mouse IgG may vary with subclass and it may be difficult to obtain high yields of active F(ab')₂ fragments without some undigested or

completely degraded IgG. In particular, IgG_{2b} is highly susceptible to complete degradation. The other subclasses require different incubation conditions to produce optimal results, all of which is known in the art.

5 Digestion of rat IgG by pepsin requires conditions including dialysis in 0.1 M acetate buffer, pH 4.5, and then incubation for four hours with 1% w/w pepsin; IgG₁ and IgG_{2a} digestion is improved if first dialyzed against 0.1 M formate buffer, pH 2.8, at 4°C, for 16 hours followed by acetate buffer. IgG_{2b} gives more consistent results with incubation in staphylococcal V8 protease (3% w/w) in 0.1 M sodium phosphate buffer, pH 7.8, for four hours at 37°C.

10

The following patents and patent applications are specifically incorporated herein by reference for the purposes of even further supplementing the present teachings regarding the preparation and use of functional, antigen-binding regions of antibodies, including scFv, Fv, Fab', Fab and F(ab')₂ fragments of the anti aminophospholipid antibodies: U.S. Patent Nos. 5,855,866 and 5,877,289, ^{5,915,132; 6,004,555, and 6,048,319} and U.S. Applications Serial Nos. 08/350,212 (Patent No. 5, ~~_____~~; Issue Fee paid); 08/482,369 (U.S. Patent 5, ~~_____~~; Issue Fee paid); and 08/487,427 (U.S. Patent 5, ~~_____~~; Issue Fee paid).

F. Pharmaceutical Compositions

20 The most basic pharmaceutical compositions of the present invention will generally comprise an effective amount of at least a first naked anti-aminophospholipid antibody, or antigen-binding fragment thereof, dissolved or dispersed in a pharmaceutically acceptable carrier or aqueous medium. Combined therapeutics are also contemplated, and the same type of underlying pharmaceutical compositions may be employed for both single and combined
25 medicaments.

The phrases "pharmaceutically or pharmacologically acceptable" refer to molecular entities and compositions that do not produce an adverse, allergic or other untoward reaction when administered to an animal, or a human, as appropriate. As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and
30

antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the therapeutic compositions is contemplated. For human administration, preparations should meet sterility,
5 pyrogenicity, general safety and purity standards as required by FDA Office of Biologics standards. Supplementary active ingredients can also be incorporated into the compositions.

F1. Parenteral Formulations

The anti-aminophospholipid antibodies of the present invention will most often be
10 formulated for parenteral administration, *e.g.*, formulated for injection *via* the intravenous, intramuscular, sub-cutaneous, transdermal, or other such routes, including peristaltic administration and direct instillation into a tumor or disease site (intracavity administration). The preparation of an aqueous composition that contains an anti-aminophospholipid antibody as an active ingredient will be known to those of skill in the art in light of the present disclosure.
15 Typically, such compositions can be prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for using to prepare solutions or suspensions upon the addition of a liquid prior to injection can also be prepared; and the preparations can also be emulsified.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or
20 dispersions; formulations including sesame oil, peanut oil or aqueous propylene glycol; and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases, the form should be sterile and fluid to the extent that syringability exists. It should be stable under the conditions of manufacture and storage and should be preserved against the contaminating action of microorganisms, such as bacteria and fungi.

25

The anti-aminophospholipid antibody compositions can be formulated into a sterile aqueous composition in a neutral or salt form. Solutions of the anti-aminophospholipid antibodies as free base or pharmacologically acceptable salts can be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Pharmaceutically acceptable salts,
30 include the acid addition salts (formed with the free amino groups of the protein), and those that

are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, trifluoroacetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, histidine, procaine and the like.

Suitable carriers include solvents and dispersion media containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion and/or by the use of surfactants.

Under ordinary conditions of storage and use, all such preparations should contain a preservative to prevent the growth of microorganisms. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

Prior to or upon formulation, the anti-aminophospholipid antibodies should be extensively dialyzed to remove undesired small molecular weight molecules, and/or lyophilized for more ready formulation into a desired vehicle, where appropriate. Sterile injectable solutions are prepared by incorporating the active anti-aminophospholipid antibodies in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as desired, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle that contains the basic dispersion medium and the required other ingredients from those enumerated above.

In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques that yield a powder of the active anti-aminophospholipid antibody ingredient, plus any additional desired ingredient from a previously sterile-filtered solution thereof.

5

Suitable pharmaceutical compositions in accordance with the invention will generally include an amount of the anti-aminophospholipid antibody admixed with an acceptable pharmaceutical diluent or excipient, such as a sterile aqueous solution, to give a range of final concentrations, depending on the intended use. The techniques of preparation are generally well known in the art as exemplified by Remington's Pharmaceutical Sciences, 16th Ed. Mack Publishing Company, 1980, incorporated herein by reference. It should be appreciated that endotoxin contamination should be kept minimally at a safe level, for example, less than 0.5 ng/mg protein. Moreover, for human administration, preparations should meet sterility, pyrogenicity, general safety and purity standards as required by FDA Office of Biological Standards.

10
15
20
25

Upon formulation, anti-aminophospholipid antibody solutions will be administered in a manner compatible with the dosage formulation and in such amount as is therapeutically effective. Formulations are easily administered in a variety of dosage forms, such as the type of injectable solutions described above, but other pharmaceutically acceptable forms are also contemplated, *e.g.*, tablets, pills, capsules or other solids for oral administration, suppositories, pessaries, nasal solutions or sprays, aerosols, inhalants, liposomal forms and the like. Pharmaceutical "slow release" capsules or compositions may also be used. Slow release formulations are generally designed to give a constant drug level over an extended period and may be used to deliver anti-aminophospholipid antibodies in accordance with the present invention.

F2. Liposomes and Nanocapsules

In certain embodiments, liposomes and/or nanoparticles may also be employed with the anti-aminophospholipid antibodies. The formation and use of liposomes is generally known to those of skill in the art, as summarized below.

5

Liposomes are formed from phospholipids that are dispersed in an aqueous medium and spontaneously form multilamellar concentric bilayer vesicles (also termed multilamellar vesicles (MLVs)). MLVs generally have diameters of from 25 nm to 4 μ m. Sonication of MLVs results in the formation of small unilamellar vesicles (SUVs) with diameters in the range of 200 to 10 500 Å, containing an aqueous solution in the core.

Phospholipids can form a variety of structures other than liposomes when dispersed in water, depending on the molar ratio of lipid to water. At low ratios the liposome is the preferred structure. The physical characteristics of liposomes depend on pH, ionic strength and the presence of divalent cations. Liposomes can show low permeability to ionic and polar substances, but at elevated temperatures undergo a phase transition which markedly alters their permeability. The phase transition involves a change from a closely packed, ordered structure, known as the gel state, to a loosely packed, less-ordered structure, known as the fluid state. This occurs at a characteristic phase-transition temperature and results in an increase in permeability to ions, sugars and drugs.

Liposomes interact with cells via four different mechanisms: Endocytosis by phagocytic cells of the reticuloendothelial system such as macrophages and neutrophils; adsorption to the cell surface, either by nonspecific weak hydrophobic or electrostatic forces, or by specific 25 interactions with cell-surface components; fusion with the plasma cell membrane by insertion of the lipid bilayer of the liposome into the plasma membrane, with simultaneous release of liposomal contents into the cytoplasm; and by transfer of liposomal lipids to cellular or subcellular membranes, or *vice versa*, without any association of the liposome contents. Varying the liposome formulation can alter which mechanism is operative, although more than one may 30 operate at the same time.

Nanocapsules can generally entrap compounds in a stable and reproducible way. To avoid side effects due to intracellular polymeric overloading, such ultrafine particles (sized around 0.1 μm) should be designed using polymers able to be degraded *in vivo*. Biodegradable polyalkyl-cyanoacrylate nanoparticles that meet these requirements are contemplated for use in the present invention, and such particles may be easily made.

G. Therapeutic Kits

This invention also provides therapeutic kits comprising anti-aminophospholipid antibodies for use in the present treatment methods. Such kits will generally contain, in suitable container means, a pharmaceutically acceptable formulation of at least one anti-aminophospholipid antibody. The kits may also contain other pharmaceutically acceptable formulations, either for diagnosis/imaging or combined therapy. For example, such kits may contain any one or more of a range of chemotherapeutic or radiotherapeutic drugs; anti-angiogenic agents; anti-tumor cell antibodies; and/or anti-tumor vasculature or anti-tumor stroma immunotoxins or coaguligands.

The kits may have a single container (container means) that contains the anti-aminophospholipid antibody, with or without any additional components, or they may have distinct containers for each desired agent. Where combined therapeutics are provided, a single solution may be pre-mixed, either in a molar equivalent combination, or with one component in excess of the other. Alternatively, each of the anti-aminophospholipid antibody and other anti-cancer agent components of the kit may be maintained separately within distinct containers prior to administration to a patient.

25

When the components of the kit are provided in one or more liquid solutions, the liquid solution is preferably an aqueous solution, with a sterile aqueous solution being particularly preferred. However, the components of the kit may be provided as dried powder(s). When reagents or components are provided as a dry powder, the powder can be reconstituted by the

addition of a suitable solvent. It is envisioned that the solvent may also be provided in another container.

5 The containers of the kit will generally include at least one vial, test tube, flask, bottle, syringe or other container means, into which the anti-aminophospholipid antibody, and any other desired agent, may be placed and, preferably, suitably aliquoted. Where separate components are included, the kit will also generally contain a second vial or other container into which these are placed, enabling the administration of separated designed doses. The kits may also comprise a second/third container means for containing a sterile, pharmaceutically acceptable buffer or other
10 diluent.

15 The kits may also contain a means by which to administer the anti-aminophospholipid antibody to an animal or patient, *e.g.*, one or more needles or syringes, or even an eye dropper, pipette, or other such like apparatus, from which the formulation may be injected into the animal or applied to a diseased area of the body. The kits of the present invention will also typically include a means for containing the vials, or such like, and other component, in close confinement for commercial sale, such as, *e.g.*, injection or blow-molded plastic containers into which the desired vials and other apparatus are placed and retained.

20 H. Tumor Treatment

25 The most important use of the present invention is in the treatment of vascularized, malignant tumors; with the treatment of benign tumors, such as BPH, also being contemplated. The invention may also be used in the therapy of other diseases and disorders having, as a component of the disease, prothrombotic blood vessels. Such vasculature-associated diseases include diabetic retinopathy, macular degeneration, vascular restenosis, including restenosis following angioplasty, arteriovenous malformations (AVM), meningioma, hemangioma, neovascular glaucoma and psoriasis; and also angiofibroma, arthritis, rheumatoid arthritis, atherosclerotic plaques, corneal graft neovascularization, hemophilic joints, hypertrophic scars, osler-weber syndrome, pyogenic granuloma retrolental fibroplasia, scleroderma, trachoma,

vascular adhesions, synovitis, dermatitis, various other inflammatory diseases and disorders, and even endometriosis.

5 The anti-aminophospholipid antibody treatment of the invention is most preferably exploited for the treatment of solid tumors. Such uses may employ anti-aminophospholipid antibodies alone or in combination with chemotherapeutic, radiotherapeutic, apoptotic, anti-angiogenic agents and/or immunotoxins or coaguligands. The anti-aminophospholipid antibody methods provided by this invention are broadly applicable to the treatment of any malignant tumor having a vascular component. Typical vascularized tumors are the solid tumors, particularly carcinomas, which require a vascular component for the provision of oxygen and nutrients. Exemplary solid tumors that may be treated using the invention include, but are not limited to, carcinomas of the lung, breast, ovary, stomach, pancreas, larynx, esophagus, testes, liver, parotid, biliary tract, colon, rectum, cervix, uterus, endometrium, kidney, bladder, prostate, thyroid, squamous cell carcinomas, adenocarcinomas, small cell carcinomas, melanomas, gliomas, neuroblastomas, and the like.

20 The present invention is contemplated for use in the treatment of any patient that presents with a solid tumor. However, in that this invention is particularly successful in the treatment of solid tumors of moderate or large sizes, patients in these categories are likely to receive more significant benefits from treatment in accordance with the methods and compositions provided herein.

25 Therefore, in general, the invention can be used to treat tumors of about 0.3-0.5 cm and upwards, although it is a better use of the invention to treat tumors of greater than 0.5 cm in size. From the studies already conducted in acceptable animal models, it is believed that patients presenting with tumors of between about 1.0 and about 2.0 cm in size will be in the preferred treatment group of patients for anti-aminophospholipid antibody therapy, although tumors up to and including the largest tumors found in humans may also be treated.

Although the present invention is not generally intended as a preventative or prophylactic treatment, use of the invention is certainly not confined to the treatment of patients having tumors of only moderate or large sizes. There are many reasons underlying this aspect of the breadth of the invention. For example, a patient presenting with a primary tumor of moderate size or above may also have various other metastatic tumors that are considered to be small-sized or even in the earlier stages of metastatic tumor seeding. Given that the anti-aminophospholipid antibodies, or combinations, of the invention are generally administered into the systemic circulation of a patient, they will naturally have effects on the secondary, smaller and metastatic tumors, although this may not be the primary intent of the treatment. Furthermore, even in situations where the tumor mass as a whole is a single small tumor, certain beneficial anti-tumor effects will result from the use of the present anti-aminophospholipid antibody treatment.

The guidance provided herein regarding the most suitable patients for use in connection with the present invention is intended as teaching that certain patient's profiles may assist with the selection of patients for treatment by the present invention. The pre-selection of certain patients, or categories of patients, does not in any way negate the basic usefulness of the present invention in connection with the treatment of all patients having a vascularized tumor. A further consideration is the fact that the assault on the tumor provided by the anti-aminophospholipid antibody therapy of the invention may predispose the tumor to further therapeutic treatment, such that the subsequent treatment results in an overall synergistic effect or even leads to total remission or cure.

It is not believed that any particular type of tumor should be excluded from treatment using the present invention. However, the type of tumor cells may be relevant to the use of the invention in combination with secondary therapeutic agents, particularly chemotherapeutics and anti-tumor cell immunotoxins. As the effect of the present therapy is to destroy the tumor vasculature, and as the vasculature is substantially or entirely the same in all solid tumors, it will be understood that the present anti-aminophospholipid methodology is widely or entirely applicable to the treatment of all solid tumors, irrespective of the particular phenotype or

genotype of the tumor cells themselves. The data presented herein is compelling as it shows impressive results in a tumor model that is resistant to necrosis.

Therapeutically effective doses of anti-aminophospholipid antibodies are readily
5 determinable using data from an animal model, as shown in the studies detailed herein. Experimental animals bearing solid tumors are frequently used to optimize appropriate therapeutic doses prior to translating to a clinical environment. Such models are known to be very reliable in predicting effective anti-cancer strategies. For example, mice bearing solid tumors, such as used in the Examples, are widely used in pre-clinical testing. The inventors have
10 used such art-accepted mouse models to determine working ranges of naked anti-aminophospholipid antibodies that give beneficial anti-tumor effects with minimal toxicity.

As is known in the art, there are realistic objectives that may be used as a guideline in connection with pre-clinical testing before proceeding to clinical treatment. However, due to the
15 safety already demonstrated in accepted models, pre-clinical testing of the present invention will be more a matter of optimization, rather than to confirm effectiveness. Thus, pre-clinical testing may be employed to select the most advantageous anti-aminophospholipid antibodies, doses or combinations.

Any anti-aminophospholipid antibody dose, or combined medicament, that results in any
20 consistent detectable tumor vasculature destruction, thrombosis and anti-tumor effects will still define a useful invention. Destructive, thrombotic and necrotic effects should be observed in between about 10% and about 40-50% of the tumor blood vessels and tumor tissues, upwards to between about 50% and about 99% of such effects being observed. The present invention may
25 also be effective against vessels downstream of the tumor, *i.e.*, target at least a sub-set of the draining vessels, particularly as cytokines released from the tumor will be acting on these vessels, changing their antigenic profile.

It will also be understood that even in such circumstances where the anti-tumor effects of
30 the anti-aminophospholipid antibody dose, or combined therapy, are towards the low end of this

range, it may be that this therapy is still equally or even more effective than all other known therapies in the context of the particular tumor targets. It is unfortunately evident to a clinician that certain tumors cannot be effectively treated in the intermediate or long term, but that does not negate the usefulness of the present therapy, particularly where it is at least about as effective
5 as the other strategies generally proposed.

In designing appropriate doses of anti-aminophospholipid antibodies, or combined therapeutics, for the treatment of vascularized tumors, one may readily extrapolate from the animal studies described herein in order to arrive at appropriate doses for clinical administration.
10 To achieve this conversion, one would account for the mass of the agents administered per unit mass of the experimental animal and, preferably, account for the differences in the body surface area between the experimental animal and the human patient. All such calculations are well known and routine to those of ordinary skill in the art.

For example, in taking the successful dose of 20 μ g antibody per mouse (total body weight of about 20 g), and applying standard calculations based upon mass and surface area, effective doses for use in human patients would be between about 1 mg and about 500 mgs antibody per patient, and preferably, between about 10 mgs and about 100 mgs antibody per patient. In addition to all variable clinical and therapeutic parameters, this variation also accounts for the present tumor necrosis data being generated using a tumor model that is resistant to necrosis, and in which less tumor vessel staining was observed than in other models.
15
20

Accordingly, using this information, the inventors contemplate that useful low doses of naked anti-aminophospholipid antibodies for human administration will be about 1, 2, 3, 4, 5, 6,
25 7, 8, 9, 10, 15, 20, 25 or about 30 mgs or so per patient; and useful high doses of naked anti-aminophospholipid antibodies for human administration will be about 250, 275, 300, 325, 350, 375, 400, 425, 450, 475 or about 500 mgs or so per patient. Useful intermediate doses of naked anti-aminophospholipid antibodies for human administration are contemplated to be about 35, 40, 50, 60, 70, 80, 90, 100, 125, 150, 175, 200 or about 225 mgs or so per patient. Any particular

range using any of the foregoing recited exemplary doses or any value intermediate between the particular stated ranges is also contemplated.

5 In general, dosage ranges of between about 5-100 mgs, about 10-80 mgs, about 20-70 mgs, about 25-60 mgs, or about 30-50 mgs or so of antibody per patient will be preferred. Notwithstanding these stated ranges, it will be understood that, given the parameters and detailed guidance presented herein, further variations in the active or optimal ranges will be encompassed within the present invention. Although doses in and around about 5 or 10 to about 70, 80, 90 or 100 mgs per patient are currently preferred, it will be understood that lower doses may be more
10 appropriate in combination with other agents, and that high doses can still be tolerated, particularly given the enhanced safety of the unconjugated anti-aminophospholipid antibodies for use in the invention. The use of human or humanized naked anti-aminophospholipid antibodies renders the present invention even safer for clinical use, further reducing the chances of significant toxicity or side effects in healthy tissues.

09131862-071219
52120-29815

The intention of the therapeutic regimens of the present invention is generally to produce significant anti-tumor effects whilst still keeping the dose below the levels associated with unacceptable toxicity. In addition to varying the dose itself, the administration regimen can also be adapted to optimize the treatment strategy. A currently preferred treatment strategy is to administer between about 1-500 mgs, and preferably, between about 10-100 mgs of the anti-aminophospholipid antibody, or therapeutic cocktail containing such, about 3 times within about
20 a 7 day period. For example, doses would be given on about day 1, day 3 or 4 and day 6 or 7.

In administering the particular doses themselves, one would preferably provide a
25 pharmaceutically acceptable composition (according to FDA standards of sterility, pyrogenicity, purity and general safety) to the patient systemically. Intravenous injection is generally preferred, and the most preferred method is to employ a continuous infusion over a time period of about 1 or 2 hours or so. Although it is not required to determine such parameters prior to treatment using the present invention, it should be noted that the studies detailed herein result in
30 at least some thrombosis being observed specifically in the blood vessels of a solid tumor within

about 12-24 hours of injection, and that the tumor cells themselves begin to die within about 24 to 72 hours. Widespread tumor necrosis is generally observed in the next about 48-96 hours, up to and including greater than 60% necrosis being observed.

5 Naturally, before wide-spread use, clinical trials will be conducted. The various elements of conducting a clinical trial, including patient treatment and monitoring, will be known to those of skill in the art in light of the present disclosure. The following information is being presented as a general guideline for use in establishing such trials.

10 Patients chosen for the first anti-aminophospholipid antibody treatment studies will have failed to respond to at least one course of conventional therapy, and will have objectively measurable disease as determined by physical examination, laboratory techniques, and/or radiographic procedures. Any chemotherapy should be stopped at least 2 weeks before entry into the study. Where murine monoclonal antibodies or antibody portions are employed, the patients
15 should have no history of allergy to mouse immunoglobulin.

20 Certain advantages will be found in the use of an indwelling central venous catheter with a triple lumen port. The anti-aminophospholipid antibodies should be filtered, for example, using a 0.22 μ filter, and diluted appropriately, such as with saline, to a final volume of 100 ml. Before use, the test sample should also be filtered in a similar manner, and its concentration assessed before and after filtration by determining the A_{280} . The expected recovery should be within the range of 87% to 99%, and adjustments for protein loss can then be accounted for.

25 These anti-aminophospholipid antibodies may be administered over a period of approximately 4-24 hours, with each patient receiving 2-4 infusions at 2-7 day intervals. Administration can also be performed by a steady rate of infusion over a 7 day period. The infusion given at any dose level should be dependent upon any toxicity observed. Hence, if Grade II toxicity was reached after any single infusion, or at a particular period of time for a steady rate infusion, further doses should be withheld or the steady rate infusion stopped unless
30 toxicity improved. Increasing doses of anti-aminophospholipid antibodies should be

administered to groups of patients until approximately 60% of patients showed unacceptable Grade III or IV toxicity in any category. Doses that are 2/3 of this value are defined as the safe dose.

- 5 Physical examination, tumor measurements, and laboratory tests should, of course, be performed before treatment and at intervals up to 1 month later. Laboratory tests should include complete blood counts, serum creatinine, creatine kinase, electrolytes, urea, nitrogen, SGOT, bilirubin, albumin, and total serum protein. Serum samples taken up to 60 days after treatment should be evaluated by radioimmunoassay for the presence of the administered anti-aminophospholipid antibodies, and antibodies against any portions thereof. Immunological analyses of sera, using any standard assay such as, for example, an ELISA or RIA, will allow the pharmacokinetics and clearance of the anti-aminophospholipid therapeutic agent to be evaluated.
- 10

To evaluate the anti-tumor responses, the patients should be examined at 48 hours to 1 week and again at 30 days after the last infusion. When palpable disease was present, two perpendicular diameters of all masses should be measured daily during treatment, within 1 week after completion of therapy, and at 30 days. To measure nonpalpable disease, serial CT scans could be performed at 1-cm intervals throughout the chest, abdomen, and pelvis at 48 hours to 1 week and again at 30 days. Tissue samples should also be evaluated histologically, and/or by flow cytometry, using biopsies from the disease sites or even blood or fluid samples if appropriate.

15

20

Clinical responses may be defined by acceptable measure. For example, a complete response may be defined by the disappearance of all measurable tumor 1 month after treatment.

- 25 Whereas a partial response may be defined by a 50% or greater reduction of the sum of the products of perpendicular diameters of all evaluable tumor nodules 1 month after treatment, with no tumor sites showing enlargement. Similarly, a mixed response may be defined by a reduction of the product of perpendicular diameters of all measurable lesions by 50% or greater 1 month after treatment, with progression in one or more sites.

In light of results from clinical trials, such as those described above, an even more precise treatment regimen may be formulated. Even so, some variation in dosage may later be necessary depending on the condition of the subject being treated. The physician responsible for administration will, in light of the present disclosure, be able to determine the appropriate dose for the individual subject. Such optimization and adjustment is routinely carried out in the art and by no means reflects an undue amount of experimentation.

I. Tumor Imaging

The present invention further provides combined tumor treatment and imaging methods, based upon anti-aminophospholipid binding ligands. Anti-aminophospholipid binding proteins or antibodies that are linked to one or more detectable agents are envisioned for use in pre-imaging the tumor, forming a reliable image prior to the treatment, which itself targets the aminophospholipid markers. In addition to antibodies, the use of detectably labeled annexins and other aminophospholipid binding ligands is contemplated, as disclosed and claimed in first and second provisional applications Serial Nos. 60/092,589 (filed July 13, 1998) and 60/110,600 (filed December 02, 1998) and in co-filed U.S. and PCT patent applications (Attorney Docket Nos. 4001.002300, 4001.002382, 4001.002383 and 4001.002210), each specifically incorporated herein by reference.

The anti-aminophospholipid imaging ligands or antibodies, or conjugates thereof, will generally comprise an anti-aminophospholipid antibody or binding ligand operatively attached, or conjugated to, a detectable label. "Detectable labels" are compounds or elements that can be detected due to their specific functional properties, or chemical characteristics, the use of which allows the component to which they are attached to be detected, and further quantified if desired. Preferably, the detectable labels are those detectable *in vivo* using non-invasive methods.

Antibody and binding protein conjugates for use as diagnostic agents generally fall into two classes, those for use in *in vitro* diagnostics, such as in a variety of immunoassays, and those for use *in vivo* diagnostic protocols. It is the *in vivo* imaging methods that are particularly intended for use with this invention.

Many appropriate imaging agents are known in the art, as are methods for their attachment to antibodies and binding ligands (see, *e.g.*, U.S. patents 5,021,236 and 4,472,509, both incorporated herein by reference). Certain attachment methods involve the use of a metal
5 chelate complex employing, for example, an organic chelating agent such as DTPA attached to the antibody (U.S. Patent 4,472,509). Monoclonal antibodies may also be reacted with an enzyme in the presence of a coupling agent such as glutaraldehyde or periodate. Conjugates with fluorescein markers are prepared in the presence of these coupling agents or by reaction with an isothiocyanate.

10

An example of detectable labels are the paramagnetic ions. In this case, suitable ions include chromium (III), manganese (II), iron (III), iron (II), cobalt (II), nickel (II), copper (II), neodymium (III), samarium (III), ytterbium (III), gadolinium (III), vanadium (II), terbium (III), dysprosium (III), holmium (III) and erbium (III), with gadolinium being particularly preferred.

Ions useful in other contexts, such as X-ray imaging, include but are not limited to lanthanum (III), gold (III), lead (II), and especially bismuth (III). Fluorescent labels include rhodamine, fluorescein and renographin. Rhodamine and fluorescein are often linked via an isothiocyanate intermediate.

In the case of radioactive isotopes for diagnostic applications, suitable examples include ¹⁴carbon, ⁵¹chromium, ³⁶chlorine, ⁵⁷cobalt, ⁵⁸cobalt, copper⁶⁷, ¹⁵²Eu, gallium⁶⁷, ³hydrogen, iodine¹²³, iodine¹²⁵, iodine¹³¹, indium¹¹¹, ⁵⁹iron, ³²phosphorus, rhenium¹⁸⁶, rhenium¹⁸⁸, ⁷⁵selenium, ³⁵sulphur, technetium^{99m} and yttrium⁹⁰. ¹²⁵I is often being preferred for use in certain embodiments, and
25 technetium^{99m} and indium¹¹¹ are also often preferred due to their low energy and suitability for long range detection.

Radioactively labeled anti-aminophospholipid antibodies and binding ligands for use in the present invention may be produced according to well-known methods in the art. For
30 instance, intermediary functional groups that are often used to bind radioisotopic metallic ions to

antibodies are diethylenetriaminepentaacetic acid (DTPA) and ethylene diaminetetracetic acid (EDTA).

Monoclonal antibodies can also be iodinated by contact with sodium or potassium iodide
5 and a chemical oxidizing agent such as sodium hypochlorite, or an enzymatic oxidizing agent, such as lactoperoxidase. Anti-aminophospholipid antibodies according to the invention may be labeled with technetium-^{99m} by ligand exchange process, for example, by reducing pertechnetate with stannous solution, chelating the reduced technetium onto a Sephadex column and applying the antibody to this column; or by direct labeling techniques, *e.g.*, by incubating pertechnetate, a
10 reducing agent such as SNCl_2 , a buffer solution such as sodium-potassium phthalate solution, and the antibody.

Any of the foregoing type of detectably labeled anti-aminophospholipid antibodies and aminophospholipid binding ligands may be used in the imaging aspects of the present invention.
15 Although not previously proposed for use in combined tumor imaging and treatment, the detectably-labeled annexins of U.S. Patent No. 5,627,036; WO 95/19791; WO 95/27903; WO 95/34315; WO 96/17618; and WO 98/04294; each incorporated herein by reference; may also be employed.

WO 95/27903 (incorporated herein by reference) provides annexins for use in detecting apoptotic cells. Any of the annexin-detectable agent markers of WO 95/27903 may be used
20 herein, although it will be known that certain of these are more suitable for *in vitro* uses. WO 95/27903 is also specifically incorporated herein by reference for purposes of providing detectable kits that may be adapted for combined use with the therapeutics of the present
25 invention.

Each of WO 95/19791; WO 95/34315; WO 96/17618; and WO 98/04294; are also incorporated herein by reference for purposes of further describing radiolabelled annexin conjugates for diagnostic imaging. The intent of each of the foregoing documents is to provide
30 radiolabelled annexins for use in imaging vascular thromboses, particularly in or near the heart,

such as in deep vein thrombosis, pulmonary embolism, myocardial infarction, atrial fibrillation, problems with prosthetic cardiovascular materials, stroke, and the like. These radiolabelled annexins were also proposed for use in imaging activated platelets, *e.g.*, in conditions such as abscesses, restenosis, inflammation of joints, clots in cerebral arteries, *etc.*

5

U.S. Patent No. 5,627,036 (incorporated herein by reference) also generally concerns 'annexine' (annexin) binding ligands for use in analyzing platelet phosphatidylserine. It is explained in U.S. Patent No. 5,627,036 that hemostatic disorders, such as arterial, coronary and venous thrombosis, are usually idiopathic, which makes prediction and prevention difficult. To
10 recognize such hemostatic disorders earlier, the detection of activated platelets is proposed. The detectably labeled annexins compositions are thus disclosed in order to detect activated platelets in hemostatic disorders (U.S. Patent No. 5,627,036).

003370652-071270

Although proposing a wide range of diagnostic uses, none of WO 95/19791; WO 95/34315; WO 96/17618; or WO 98/04294 make reference to imaging the vasculature of solid tumors. Neither does U.S. Patent No. 5,627,036 make any such suggestions. Nonetheless, the disclosed detectable and radiolabelled annexin compositions *per se* may now be used to advantage in this regard, in light of the surprising discoveries disclosed herein.

20 In particular, U.S. Patent No. 5,627,036 (incorporated herein by reference) discloses annexins detectably labeled with fluorescein isothiocyanate; radioisotopes of halogens, technetium, lead, mercury, thallium or indium; and paramagnetic contrast agents.

25 WO 95/19791 (incorporated herein by reference) provides conjugates of annexin bonded to an N_2S_2 chelate that can be radiolabelled by complexing a radionuclide to the chelate. WO 95/34315 (incorporated herein by reference) provides annexin conjugates comprising one or more galactose residues with the N_2S_2 chelate. The galactose moiety is said to facilitate the rapid elimination of the radiolabelled conjugate from the circulation, reducing radiation damage to non-target tissues and background 'noise.'

30

WO 96/17618 (incorporated herein by reference) in turn provides annexin conjugates suitable for radiolabeling with diagnostic imaging agents that comprise an annexin with a cluster of galactose residues and an N₂S₂ chelate. These are reported to have a shorter circulating half-life and a higher binding affinity for target sites than the foregoing radiolabeled annexin-galactose conjugates.

Still further radiolabeled annexin conjugates are provided by WO 98/04294 (incorporated herein by reference). These conjugates comprise an annexin that is modified to provide an accessible sulphhydryl group conjugated to a hexose moiety that is recognized by a mammalian liver receptor. Annexin multimer conjugates and chelating compounds conjugated via esterase-sensitive bonds are also provided.

Each of WO 95/19791; WO 95/34315; WO 96/17618; and WO 98/04294; are also specifically incorporated herein by reference for purposes of providing annexin conjugate components for radiolabelling that are amenable to packaging in "cold kits", i.e., wherein the components are provided in separate vials. U.S. Patent No. 5,627,036 similarly provides kits comprising a carrier being compartmentalized to receive detectably labeled annexins that may be adapted for use herewith.

Although suitable for use in *in vitro* diagnostics, the present aminophospholipid detection methods are more intended for forming an image of the tumor vasculature of a patient prior to treatment with therapeutic agent-targeting agent constructs. The *in vivo* diagnostic or imaging methods generally comprise administering to a patient a diagnostically effective amount of an anti-aminophospholipid antibody or binding ligand that is conjugated to a marker that is detectable by non-invasive methods. The antibody- or binding ligand-marker conjugate is allowed sufficient time to localize and bind to the aminophospholipid expressed on the luminal surface of the tumor vasculature. The patient is then exposed to a detection device to identify the detectable marker, thus forming an image of the tumor vasculature.

5 The nuclear magnetic spin-resonance isotopes, such as gadolinium, are detected using a nuclear magnetic imaging device; and radioactive substances, such as technetium^{99m} or indium¹¹¹, are detected using a gamma scintillation camera or detector. U.S. Patent No. 5,627,036 is also specifically incorporated herein by reference for purposes of providing even further guidance regarding the safe and effective introduction of such detectably labeled constructs into the blood of an individual, and means for determining the distribution of the detectably labeled annexin extracorporally, *e.g.*, using a gamma scintillation camera or by magnetic resonance measurement.

10 Dosages for imaging embodiments are generally less than for therapy, but are also dependent upon the age and weight of a patient. A one time dose of between about 0.1, 0.5 or about 1 mg and about 9 or 10 mgs, and more preferably, of between about 1 mg and about 5-10 mgs of anti-aminophospholipid antibody- or aminophospholipid binding ligand-conjugate per patient is contemplated to be useful. U.S. Patent No. 5,627,036; and WO 95/19791, each incorporated herein by reference, are also instructive regarding doses of detectably-labeled annexins.

15 J. Combination Therapies

20 The anti-aminophospholipid antibody treatment methods of the present invention may be combined with any other methods generally employed in the treatment of the particular tumor, disease or disorder that the patient exhibits. So long as a particular therapeutic approach is not known to be detrimental to the patient's condition in itself, and does not significantly counteract the anti-aminophospholipid antibody treatment, its combination with the present invention is contemplated.

25 In connection solid tumor treatment, the present invention may be used in combination with classical approaches, such as surgery, radiotherapy, chemotherapy, and the like. The invention therefore provides combined therapies in which anti-aminophospholipid antibodies are used simultaneously with, before, or after surgery or radiation treatment; or are administered to patients with, before, or after conventional chemotherapeutic, radiotherapeutic or anti-angiogenic agents, or targeted immunotoxins or coaguligands.

30

Combination therapy for other vascular diseases is also contemplated. A particular example of such is benign prostatic hyperplasia (BPH), which may be treated with anti-aminophospholipid antibodies in combination other treatments currently practiced in the art. For
5 example, targeting of immunotoxins to markers localized within BPH, such as PSA.

When one or more agents are used in combination with the anti-aminophospholipid antibody therapy, there is no requirement for the combined results to be additive of the effects observed when each treatment is conducted separately. Although at least additive effects are
10 generally desirable, any increased anti-tumor effect above one of the single therapies would be of benefit. Also, there is no particular requirement for the combined treatment to exhibit synergistic effects, although this is certainly possible and advantageous. Agents particularly contemplated for use in achieving potentially synergistic effects are those that injure, or induce apoptosis in, the tumor endothelium, as such injury or apoptosis should amplify the overall therapeutic effect.

To practice combined anti-tumor therapy, one would simply administer to an animal an anti-aminophospholipid antibody in combination with another anti-cancer agent in a manner effective to result in their combined anti-tumor actions within the animal. The agents would
15 therefore be provided in amounts effective and for periods of time effective to result in their combined presence within the tumor vasculature and their combined actions in the tumor environment. To achieve this goal, the anti-aminophospholipid antibodies and anti-cancer agents may be administered to the animal simultaneously, either in a single composition, or as two distinct compositions using different administration routes.

25 Alternatively, the anti-aminophospholipid antibody treatment may precede, or follow, the anti-cancer agent treatment by, *e.g.*, intervals ranging from minutes to weeks. In certain embodiments where the anti-cancer agent and anti-aminophospholipid antibody are applied separately to the animal, one would ensure that a significant period of time did not expire between the time of each delivery, such that the anti-cancer agent and anti-aminophospholipid
30 antibody composition would still be able to exert an advantageously combined effect on the

tumor. In such instances, it is contemplated that one would contact the tumor with both agents within about 5 minutes to about one week of each other and, more preferably, within about 12-72 hours of each other, with a delay time of only about 12-48 hours being most preferred.

5 Exemplary anti-cancer agents that would be given prior to the anti-aminophospholipid antibody are agents that induce the expression of aminophospholipids within the tumor vasculature. For example, agents that stimulate localized calcium production and/or that induce apoptosis will generally result in increased PS expression, which can then be targeted using a subsequent anti-PS antibody. Anti-aminophospholipid antibodies would be first administered in
10 other situations to cause tumor destruction, followed by anti-angiogenic therapies or therapies directed to targeting necrotic tumor cells.

The general use of combinations of substances in cancer treatment is well know. For example, U.S. Patent No. 5,710,134 (incorporated herein by reference) discloses components that induce necrosis in tumors in combination with non-toxic substances or "prodrugs". The enzymes set free by necrotic processes cleave the non-toxic "prodrug" into the toxic "drug", which leads to tumor cell death. Also, U.S. Patent No. 5,747,469 (incorporated herein by reference) discloses the combined use of viral vectors encoding p53 and DNA damaging agents. Any such similar approaches can be used with the present invention.

In some situations, it may even be desirable to extend the time period for treatment significantly, where several days (2, 3, 4, 5, 6 or 7), several weeks (1, 2, 3, 4, 5, 6, 7 or 8) or even several months (1, 2, 3, 4, 5, 6, 7 or 8) lapse between the respective administrations. This would be advantageous in circumstances where one treatment was intended to substantially destroy the
25 tumor, such as the anti-aminophospholipid antibody treatment, and another treatment was intended to prevent micrometastasis or tumor re-growth, such as the administration of an anti-angiogenic agent. The EN 7/44 antibody of Hagemeyer *et al.* (1986) is not believed to be an effective anti-angiogenic agent, lacking binding to a surface accessible antigen, amongst other deficiencies.

It also is envisioned that more than one administration of either the anti-aminophospholipid antibody or the anti-cancer agent will be utilized. The anti-aminophospholipid antibodies and anti-cancer agents may be administered interchangeably, on alternate days or weeks; or a sequence of anti-aminophospholipid antibody treatment may be given, followed by a sequence of anti-cancer agent therapy. In any event, to achieve tumor regression using a combined therapy, all that is required is to deliver both agents in a combined amount effective to exert an anti-tumor effect, irrespective of the times for administration.

In terms of surgery, any surgical intervention may be practiced in combination with the present invention. In connection with radiotherapy, any mechanism for inducing DNA damage locally within tumor cells is contemplated, such as γ -irradiation, X-rays, UV-irradiation, microwaves and even electronic emissions and the like. The directed delivery of radioisotopes to tumor cells is also contemplated, and this may be used in connection with a targeting antibody or other targeting means.

Cytokine therapy also has proven to be an effective partner for combined therapeutic regimens. Various cytokines may be employed in such combined approaches. Examples of cytokines include IL-1 α , IL-1 β , IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, TGF- β , GM-CSF, M-CSF, G-CSF, TNF α , TNF β , LAF, TCGF, BCGF, TRF, BAF, BDG, MP, LIF, OSM, TMF, PDGF, IFN- α , IFN- β , IFN- γ . Cytokines are administered according to standard regimens, consistent with clinical indications such as the condition of the patient and relative toxicity of the cytokine. Uteroglobins may also be used to prevent or inhibit metastases (U.S. Patent No. 5,696,092; incorporated herein by reference).

J1. Chemotherapeutics

In certain embodiments, the anti-aminophospholipid antibodies of the present invention may be administered in combination with a chemotherapeutic agent. Chemotherapeutic drugs can kill proliferating tumor cells, enhancing the necrotic areas created by the overall treatment. The drugs can thus enhance the thrombotic action of the anti-aminophospholipid antibodies.

By inducing the formation of thrombi in tumor vessels, the anti-aminophospholipid antibodies can enhance the action of the chemotherapeutics by retaining or trapping the drugs within the tumor. The chemotherapeutics are thus retained within the tumor, while the rest of the drug is cleared from the body. Tumor cells are thus exposed to a higher concentration of drug for a longer period of time. This entrapment of drug within the tumor makes it possible to reduce the dose of drug, making the treatment safer as well as more effective.

Irrespective of the underlying mechanism(s), a variety of chemotherapeutic agents may be used in the combined treatment methods disclosed herein. Chemotherapeutic agents contemplated as exemplary include, *e.g.*, tamoxifen, taxol, vincristine, vinblastine, etoposide (VP-16), adriamycin, 5-fluorouracil (5FU), camptothecin, actinomycin-D, mitomycin C, cisplatin (CDDP), combretastatin(s) and derivatives and prodrugs thereof.

As will be understood by those of ordinary skill in the art, the appropriate doses of chemotherapeutic agents will be generally around those already employed in clinical therapies wherein the chemotherapeutics are administered alone or in combination with other chemotherapeutics. By way of example only, agents such as cisplatin, and other DNA alkylating may be used. Cisplatin has been widely used to treat cancer, with efficacious doses used in clinical applications of 20 mg/m² for 5 days every three weeks for a total of three courses. Cisplatin is not absorbed orally and must therefore be delivered *via* injection intravenously, subcutaneously, intratumorally or intraperitoneally.

Further useful agents include compounds that interfere with DNA replication, mitosis, chromosomal segregation and/or tubulin activity. Such chemotherapeutic compounds include adriamycin, also known as doxorubicin, etoposide, verapamil, podophyllotoxin(s), combretastatin(s) and the like. Widely used in a clinical setting for the treatment of neoplasms, these compounds are administered through bolus injections intravenously at doses ranging from 25-75 mg/m² at 21 day intervals for adriamycin, to 35-50 mg/m² for etoposide intravenously or double the intravenous dose orally.

TABLE B

CHEMOTHERAPEUTIC AGENTS USEFUL IN NEOPLASTIC DISEASE

| CLASS | TYPE OF AGENT | NONPROPRIETARY NAMES (OTHER NAMES) | DISEASE |
|--------------------------|-----------------------------------|---|---|
| <i>Alkylating Agents</i> | Nitrogen Mustards | Mechlorethamine (HN ₂) | Hodgkin's disease, non-Hodgkin's lymphomas |
| | | Cyclophosphamide Ifosfamide | Acute and chronic lymphocytic leukemias, Hodgkin's disease, non-Hodgkin's lymphomas, multiple myeloma, neuroblastoma, breast, ovary, lung, Wilms' tumor, cervix, testis, soft-tissue sarcomas |
| | | Melphalan (L-sarcolysin) | Multiple myeloma, breast, ovary |
| | | Chlorambucil | Chronic lymphocytic leukemia, primary macroglobulinemia, Hodgkin's disease, non-Hodgkin's lymphomas |
| | Ethylenimenes and Methylmelamines | Hexamethylmelamine | Ovary |
| | | Thiotepa | Bladder, breast, ovary |
| | Alkyl Sulfonates | Busulfan | Chronic granulocytic leukemia |
| | Nitrosoureas | Carmustine (BCNU) | Hodgkin's disease, non-Hodgkin's lymphomas, primary brain tumors, multiple myeloma, malignant melanoma |
| | | Lomustine (CCNU) | Hodgkin's disease, non-Hodgkin's lymphomas, primary brain tumors, small-cell lung |
| | | Semustine (methyl-CCNU) | Primary brain tumors, stomach, colon |
| | | Streptozocin (streptozotocin) | Malignant pancreatic insulinoma, malignant carcinoid |
| | Triazines | Dacarbazine (DTIC; dimethyltriazenoimidazolecarboxamide) | Malignant melanoma, Hodgkin's disease, soft-tissue sarcomas |
| <i>Antimetabolites</i> | Folic Acid Analogs | Methotrexate (amethopterin) | Acute lymphocytic leukemia, choriocarcinoma, mycosis fungoides, breast, head and neck, lung, osteogenic sarcoma |
| | Pyrimidine Analogs | Fluoruracil (5-fluorouracil; 5-FU) Flouxuridine (fluorodeoxyuridine; FUdR) | Breast, colon, stomach, pancreas, ovary, head and neck, urinary bladder, premalignant skin lesions (topical) |

| CLASS | TYPE OF AGENT | NONPROPRIETARY NAMES (OTHER NAMES) | DISEASE |
|--|---------------------------------------|---|---|
| <i>Antimetabolites, continued</i> | | Cytarabine (cytosine arabinoside) | Acute granulocytic and acute lymphocytic leukemias |
| | Purine Analogs and Related Inhibitors | Mercaptopurine (6-mercaptopurine; 6-MP) | Acute lymphocytic, acute granulocytic and chronic granulocytic leukemias |
| | | Thioguanine (6-thioguanine; TG) | Acute granulocytic, acute lymphocytic and chronic granulocytic leukemias |
| | | Pentostatin (2-deoxycofomycin) | Hairy cell leukemia, mycosis fungoides, chronic lymphocytic leukemia |
| <i>Natural Products</i> | Vinca Alkaloids | Vinblastine (VLB) | Hodgkin's disease, non-Hodgkin's lymphomas, breast, testis |
| | | Vincristine | Acute lymphocytic leukemia, neuroblastoma, Wilms' tumor, rhabdomyosarcoma, Hodgkin's disease, non-Hodgkin's lymphomas, small-cell lung |
| | Epipodophyllotoxins | Etoposide Tertiposide | Testis, small-cell lung and other lung, breast, Hodgkin's disease, non-Hodgkin's lymphomas, acute granulocytic leukemia, Kaposi's sarcoma |
| | Antibiotics | Dactinomycin (actinomycin D) | Choriocarcinoma, Wilms' tumor, rhabdomyosarcoma, testis, Kaposi's sarcoma |
| | | Daunorubicin (daunomycin; rubidomycin) | Acute granulocytic and acute lymphocytic leukemias |
| | | Doxorubicin | Soft-tissue, osteogenic and other sarcomas; Hodgkin's disease, non-Hodgkin's lymphomas, acute leukemias, breast, genitourinary, thyroid, lung, stomach, neuroblastoma |
| | | Bleomycin | Testis, head and neck, skin, esophagus, lung and genitourinary tract; Hodgkin's disease, non-Hodgkin's lymphomas |
| | Antibiotics, continued | Plicamycin (mithramycin) | Testis, malignant hypercalcemia |
| | | Mitomycin (mitomycin C) | Stomach, cervix, colon, breast, pancreas, bladder, head and neck |
| | Enzymes | L-Asparaginase | Acute lymphocytic leukemia |
| <i>Natural Products, continued</i> | | | |

| CLASS | TYPE OF AGENT | NONPROPRIETARY NAMES (OTHER NAMES) | DISEASE |
|---------------------------------|---------------------------------------|--|--|
| | Biological Response Modifiers | Interferon alfa | Hairy cell leukemia., Kaposi's sarcoma, melanoma, carcinoid, renal cell, ovary, bladder, non-Hodgkin's lymphomas, mycosis fungoides, multiple myeloma, chronic granulocytic leukemia |
| <i>Miscellaneous Agents</i> | Platinum Coordination Complexes | Cisplatin (<i>cis</i> -DDP) Carboplatin | Testis, ovary, bladder, head and neck, lung, thyroid, cervix, endometrium, neuroblastoma, osteogenic sarcoma |
| | Anthracenedione | Mitoxantrone | Acute granulocytic leukemia, breast |
| | Substituted Urea | Hydroxyurea | Chronic granulocytic leukemia, polycythemia vera, essential thrombocytosis, malignant melanoma |
| | Methyl Hydrazine Derivative | Procarbazine (N-methylhydrazine, MIH) | Hodgkin's disease |
| | Adrenocortical Suppressant | Mitotane (<i>o,p'</i> -DDD) | Adrenal cortex |
| | | Aminoglutethimide | Breast |
| <i>Hormones and Antagonists</i> | Adrenocorticosteroids | Prednisone (several other equivalent preparations available) | Acute and chronic lymphocytic leukemias, non-Hodgkin's lymphomas, Hodgkin's disease, breast |
| | Progestins | Hydroxyprogesterone caproate Medroxyprogesterone acetate Megestrol acetate | Endometrium, breast |
| | Estrogens | Diethylstilbestrol Ethinyl estradiol (other preparations available) | Breast, prostate |
| | Antiestrogen | Tamoxifen | Breast |
| | Androgens | Testosterone propionate Fluoxymesterone (other preparations available) | Breast |
| | Antiandrogen | Flutamide | Prostate |
| | Gonadotropin-releasing hormone analog | Leuprolide | Prostate |

J2. Anti-Angi genics

The term "angiogenesis" refers to the generation of new blood vessels, generally into a tissue or organ. Under normal physiological conditions, humans or animals undergo angiogenesis only in very specific restricted situations. For example, angiogenesis is normally
5 observed in wound healing, fetal and embryonic development and formation of the corpus luteum, endometrium and placenta. Uncontrolled (persistent and/or unregulated) angiogenesis is related to various disease states, and occurs during tumor development and metastasis.

Both controlled and uncontrolled angiogenesis are thought to proceed in a similar
10 manner. Endothelial cells and pericytes, surrounded by a basement membrane, form capillary blood vessels. Angiogenesis begins with the erosion of the basement membrane by enzymes released by endothelial cells and leukocytes. The endothelial cells, which line the lumen of blood vessels, then protrude through the basement membrane. Angiogenic stimulants induce the endothelial cells to migrate through the eroded basement membrane. The migrating cells form a
15 "sprout" off the parent blood vessel, where the endothelial cells undergo mitosis and proliferate. The endothelial sprouts merge with each other to form capillary loops, creating the new blood vessel.

As persistent, unregulated angiogenesis occurs during tumor development and metastasis,
20 the treatment methods of this invention may be used in combination with any one or more "anti-angiogenic" therapies. Exemplary anti-angiogenic agents that are useful in connection with combined therapy are listed in Table C. Each of the agents listed therein is exemplary and by no means limiting.

TABLE C
Inhibitors and Negative Regulators of Angiogenesis

| Substances | References |
|--|--|
| Angiostatin | O'Reilly <i>et al.</i> , 1994 |
| Endostatin | O'Reilly <i>et al.</i> , 1997 |
| 16kDa prolactin fragment | Ferrara <i>et al.</i> , 1991; Clapp <i>et al.</i> , 1993; D'Angelo <i>et al.</i> , 1995; Lee <i>et al.</i> , 1998 |
| Laminin peptides | Kleinman <i>et al.</i> , 1993; Yamamura <i>et al.</i> , 1993; Iwamoto <i>et al.</i> , 1996; Tryggvason, 1993 |
| Fibronectin peptides | Grant <i>et al.</i> , 1998; Sheu <i>et al.</i> , 1997 |
| Tissue metalloproteinase inhibitors (TIMP 1, 2, 3, 4) | Sang, 1998 |
| Plasminogen activator inhibitors (PAI-1, -2) | Soff <i>et al.</i> , 1995 |
| Tumor necrosis factor α (high dose, <i>in vitro</i>) | Frater-Schroder <i>et al.</i> , 1987 |
| TGF- β 1 | RayChadhury and D'Amore, 1991; Tada <i>et al.</i> , 1994 |
| Interferons (IFN- α , - β , γ) | Moore <i>et al.</i> , 1998; Lingen <i>et al.</i> , 1998 |
| ELR- CXC Chemokines: IL-12; SDF-1; MIG; Platelet factor 4 (PF-4); IP-10 | Moore <i>et al.</i> , 1998; Hiscox and Jiang, 1997; Coughlin <i>et al.</i> , 1998; Tanaka <i>et al.</i> , 1997 |
| Thrombospondin (TSP) | Good <i>et al.</i> , 1990; Frazier, 1991; Bornstein, 1992; Tolsma <i>et al.</i> , 1993; Sheibani and Frazier, 1995; Volpert <i>et al.</i> , 1998 |
| SPARC | Hasselaar and Sage, 1992; Lane <i>et al.</i> , 1992; Jendraschak and Sage, 1996 |
| 2-Methoxyoestradiol | Fotsis <i>et al.</i> , 1994 |
| Proliferin-related protein | Jackson <i>et al.</i> , 1994 |
| Suramin | Gagliardi <i>et al.</i> , 1992; Takano <i>et al.</i> , 1994; Waltenberger <i>et al.</i> , 1996; Gagliardi <i>et al.</i> , 1998; Manetti <i>et al.</i> , 1998 |
| Thalidomide | D'Amato <i>et al.</i> , 1994; Kenyon <i>et al.</i> , 1997 Wells, 1998 |
| Cortisone | Thorpe <i>et al.</i> , 1993 Folkman <i>et al.</i> , 1983 Sakamoto <i>et al.</i> , 1986 |

00361862-071219
15
A certain preferred component for use in inhibiting angiogenesis is a protein named "angiostatin". This component is disclosed in U.S. Patents 5,776,704; 5,639,725 and 5,733,876, each incorporated herein by reference. Angiostatin is a protein having a molecular weight of between about 38 kD and about 45 kD, as determined by reducing polyacrylamide gel electrophoresis, which contains approximately Kringle regions 1 through 4 of a plasminogen molecule. Angiostatin generally has an amino acid sequence substantially similar to that of a fragment of murine plasminogen beginning at amino acid number 98 of an intact murine plasminogen molecule.

10 The amino acid sequence of angiostatin varies slightly between species. For example, in human angiostatin, the amino acid sequence is substantially similar to the sequence of the above described murine plasminogen fragment, although an active human angiostatin sequence may start at either amino acid number 97 or 99 of an intact human plasminogen amino acid sequence. Further, human plasminogen may be used, as it has similar anti-angiogenic activity, as shown in a mouse tumor model.

Certain anti-angiogenic therapies have already been shown to cause tumor regressions, and angiostatin is one such agent. Endostatin, a 20 kDa COOH-terminal fragment of collagen XVIII, the bacterial polysaccharide CM101, and the antibody LM609 also have angiostatic activity. However, in light of their other properties, they are referred to as anti-vascular therapies or tumor vessel toxins, as they not only inhibit angiogenesis but also initiate the destruction of tumor vessels through mostly undefined mechanisms. Their combination with the present invention is clearly envisioned.

25 Angiostatin and endostatin have become the focus of intense study, as they are the first angiogenesis inhibitors that have demonstrated the ability to not only inhibit tumor growth but also cause tumor regressions in mice. There are multiple proteases that have been shown to produce angiostatin from plasminogen including elastase, macrophage metalloelastase (MME), matrilysin (MMP-7), and 92 kDa gelatinase B/type IV collagenase (MMP-9).

MME can produce angiostatin from plasminogen in tumors and granulocyte-macrophage colony-stimulating factor (GM-CSF) upregulates the expression of MME by macrophages inducing the production of angiostatin. The role of MME in angiostatin generation is supported by the finding that MME is in fact expressed in clinical samples of hepatocellular carcinomas from patients. Another protease thought to be capable of producing angiostatin is stromelysin-1 (MMP-3). MMP-3 has been shown to produce angiostatin-like fragments from plasminogen *in vitro*.

The mechanism of action for angiostatin is currently unclear, it is hypothesized that it binds to an unidentified cell surface receptor on endothelial cells inducing endothelial cell to undergo programmed cell death or mitotic arrest. Endostatin appears to be an even more powerful anti-angiogenesis and anti-tumor agent although its biology is much less clear. Endostatin is effective at causing regressions in a number of tumor models in mice. Tumors do not develop resistance to endostatin and, after multiple cycles of treatment, tumors enter a dormant state during which they do not increase in volume. In this dormant state, the percentage of tumor cells undergoing apoptosis was increased, yielding a population that essentially stays the same size. Endostatin is also thought to bind an unidentified endothelial cell surface receptor that mediates its effect.

CM101 is a bacterial polysaccharide that has been well characterized in its ability to induce neovascular inflammation in tumors. CM101 binds to and cross-links receptors expressed on dedifferentiated endothelium that stimulates the activation of the complement system. It also initiates a cytokine-driven inflammatory response that selectively targets the tumor. It is a uniquely antipathoangiogenic agent that downregulates the expression VEGF and its receptors. CM101 is currently in clinical trials as an anti-cancer drug, and can be used in combination herewith.

Thrombospondin (TSP-1) and platelet factor 4 (PF4) may also be used in combination with the present invention. These are both angiogenesis inhibitors that associate with heparin and are found in platelet α -granules. TSP-1 is a large 450kDa multi-domain glycoprotein that is

constituent of the extracellular matrix. TSP-1 binds to many of the proteoglycan molecules found in the extracellular matrix including, HSPGs, fibronectin, laminin, and different types of collagen. TSP-1 inhibits endothelial cell migration and proliferation *in vitro* and angiogenesis *in vivo*. TSP-1 can also suppress the malignant phenotype and tumorigenesis of transformed endothelial cells. The tumor suppressor gene p53 has been shown to directly regulate the expression of TSP-1 such that, loss of p53 activity causes a dramatic reduction in TSP-1 production and a concomitant increase in tumor initiated angiogenesis.

PF4 is a 70aa protein that is member of the CXC ELR- family of chemokines that is able to potentially inhibit endothelial cell proliferation *in vitro* and angiogenesis *in vivo*. PF4 administered intratumorally or delivered by an adenoviral vector is able to cause an inhibition of tumor growth.

Interferons and metalloproteinase inhibitors are two other classes of naturally occurring angiogenic inhibitors that can be combined with the present invention. The anti-endothelial activity of the interferons has been known since the early 1980s, however, the mechanism of inhibition is still unclear. It is known that they can inhibit endothelial cell migration and that they do have some anti-angiogenic activity *in vivo* that is possibly mediated by an ability to inhibit the production of angiogenic promoters by tumor cells. Vascular tumors in particular are sensitive to interferon, for example, proliferating hemangiomas can be successfully treated with IFN α .

Tissue inhibitors of metalloproteinases (TIMPs) are a family of naturally occurring inhibitors of matrix metalloproteinases (MMPs) that can also inhibit angiogenesis and can be used in combined treatment protocols with the present invention. MMPs play a key role in the angiogenic process as they degrade the matrix through which endothelial cells and fibroblasts migrate when extending or remodeling the vascular network. In fact, one member of the MMPs, MMP-2, has been shown to associate with activated endothelium through the integrin $\alpha v \beta 3$ presumably for this purpose. If this interaction is disrupted by a fragment of MMP-2, then angiogenesis is downregulated and in tumors growth is inhibited.

There are a number of pharmacological agents that inhibit angiogenesis, any one or more of which may be used in combination with the present invention. These include AGM-1470/TNP-470, thalidomide, and carboxyamidotriazole (CAI). Fumagillin was found to be a potent inhibitor of angiogenesis in 1990, and since then the synthetic analogues of fumagillin, AGM-1470 and TNP-470 have been developed. Both of these drugs inhibit endothelial cell proliferation *in vitro* and angiogenesis *in vivo*. TNP-470 has been studied extensively in human clinical trials with data suggesting that long-term administration is optimal.

Thalidomide was originally used as a sedative but was found to be a potent teratogen and was discontinued. In 1994 it was found that thalidomide is an angiogenesis inhibitor. Thalidomide is currently in clinical trials as an anti-cancer agent as well as a treatment of vascular eye diseases.

CAI is a small molecular weight synthetic inhibitor of angiogenesis that acts as a calcium channel blocker that prevents actin reorganization, endothelial cell migration and spreading on collagen IV. CAI inhibits neovascularization at physiological attainable concentrations and is well tolerated orally by cancer patients. Clinical trials with CAI have yielded disease stabilization in 49 % of cancer patients having progressive disease before treatment.

Cortisone in the presence of heparin or heparin fragments was shown to inhibit tumor growth in mice by blocking endothelial cell proliferation. The mechanism involved in the additive inhibitory effect of the steroid and heparin is unclear although it is thought that the heparin may increase the uptake of the steroid by endothelial cells. The mixture has been shown to increase the dissolution of the basement membrane underneath newly formed capillaries and this is also a possible explanation for the additive angiostatic effect. Heparin-cortisol conjugates also have potent angiostatic and anti-tumor effects activity *in vivo*.

Further specific angiogenesis inhibitors, including, but not limited to, Anti-Invasive Factor, retinoic acids and paclitaxel (U.S. Patent No. 5,716,981; incorporated herein by

reference); AGM-1470 (Ingber *et al.*, 1990; incorporated herein by reference); shark cartilage extract (U.S. Patent No. 5,618,925; incorporated herein by reference); anionic polyamide or polyurea oligomers (U.S. Patent No. 5,593,664; incorporated herein by reference); oxindole derivatives (U.S. Patent No. 5,576,330; incorporated herein by reference); estradiol derivatives
5 (U.S. Patent No. 5,504,074; incorporated herein by reference); and thiazolopyrimidine derivatives (U.S. Patent No. 5,599,813; incorporated herein by reference) are also contemplated for use as anti-angiogenic compositions for the combined uses of the present invention.

Compositions comprising an antagonist of an $\alpha_v\beta_3$ integrin may also be used to inhibit
10 angiogenesis in combination with the present invention. As disclosed in U.S. Patent No. 5,766,591 (incorporated herein by reference), RGD-containing polypeptides and salts thereof, including cyclic polypeptides, are suitable examples of $\alpha_v\beta_3$ integrin antagonists.

The antibody LM609 against the $\alpha_v\beta_3$ integrin also induces tumor regressions. Integrin
15 $\alpha_v\beta_3$ antagonists, such as LM609, induce apoptosis of angiogenic endothelial cells leaving the quiescent blood vessels unaffected. LM609 or other $\alpha_v\beta_3$ antagonists may also work by inhibiting the interaction of $\alpha_v\beta_3$ and MMP-2, a proteolytic enzyme thought to play an important role in migration of endothelial cells and fibroblasts.

Apoptosis of the angiogenic endothelium in this case may have a cascade effect on the
20 rest of the vascular network. Inhibiting the tumor vascular network from completely responding to the tumor's signal to expand may, in fact, initiate the partial or full collapse of the network resulting in tumor cell death and loss of tumor volume. It is possible that endostatin and angiostatin function in a similar fashion. The fact that LM609 does not affect quiescent vessels
25 but is able to cause tumor regressions suggests strongly that not all blood vessels in a tumor need to be targeted for treatment in order to obtain an anti-tumor effect.

Targeted or non-targeted angiopoietins, preferably angiopoietin-2, may also be used in combination with the present invention (see discussion below for targeted angiopoietins).
30 Angiopoietin-2 (SEQ ID NO:3 and SEQ ID NO:4) is a ligand for Tie2 and generally counteracts

blood vessel maturation/stability mediated by angiopoietin-1. It is thus an antagonist of angiopoietin-1, and acts to disturb capillary structure. In the absence of another angiogenic signal (such as VEGF), angiopoietin-2 causes vessels to destabilize and become immature (Holash *et al.*, 1999; incorporated herein by reference). Provision of targeted or non-targeted angiopoietin-2 in connection with the methods of the present invention is thus contemplated, particularly in tumors with low VEGF levels and/or in combination with VEGF inhibition. Manipulation of angiopoietin-1 and two new angiopoietins, angiopoietin-3 (mouse) and angiopoietin-4 (human), could also be used in conjunction with this invention.

Other methods of therapeutic intervention based upon altering signaling through the Tie2 receptor can also be used in combination with the present invention, such as using a soluble Tie2 receptor capable of blocking Tie2 activation (Lin *et al.*, 1998). Delivery of such a construct using recombinant adenoviral gene therapy has been shown to be effective in treating cancer and reducing metastases (Lin *et al.*, 1998).

J3. Apoptosis-Inducing Agents

Therapeutic agent-targeting agent treatment may also be combined with treatment methods that induce apoptosis in any cells within the tumor, including tumor cells and tumor vascular endothelial cells. Although many anti-cancer agents may have, as part of their mechanism of action, an apoptosis-inducing effect, certain agents have been discovered, designed or selected with this as a primary mechanism, as described below.

All anti-cancer agents have this activity

A number of oncogenes have been described that inhibit apoptosis, or programmed cell death. Exemplary oncogenes in this category include, but are not limited to, bcr-abl, bcl-2 (distinct from bcl-1, cyclin D1; GenBank accession numbers M14745, X06487; U.S. Patent No. 5,650,491; and 5,539,094; each incorporated herein by reference) and family members including Bcl-xl, Mcl-1, Bak, A1, A20. Overexpression of bcl-2 was first discovered in T cell lymphomas. bcl-2 functions as an oncogene by binding and inactivating Bax, a protein in the apoptotic pathway. Inhibition of bcl-2 function prevents inactivation of Bax, and allows the apoptotic pathway to proceed. Thus, inhibition of this class of oncogenes, *e.g.*, using antisense nucleotide

sequences, is contemplated for use in the present invention in aspects wherein enhancement of apoptosis is desired (U.S. Patent No. 5,650,491; 5,539,094; and 5,583,034; each incorporated herein by reference).

5 Many forms of cancer have reports of mutations in tumor suppressor genes, such as p53. Inactivation of p53 results in a failure to promote apoptosis. With this failure, cancer cells progress in tumorigenesis, rather than become destined for cell death. Thus, provision of tumor suppressors is also contemplated for use in the present invention to stimulate cell death. Exemplary tumor suppressors include, but are not limited to, p53, Retinoblastoma gene (Rb),
10 Wilm's tumor (WT1), bax alpha, interleukin-1 β -converting enzyme and family, MEN-1 gene, neurofibromatosis, type 1 (NF1), cdk inhibitor p16, colorectal cancer gene (DCC), familial adenomatosis polyposis gene (FAP), multiple tumor suppressor gene (MTS-1), BRCA1 and BRCA2.

15 Preferred for use are the p53 (U.S. Patent No. 5,747,469; 5,677,178; and 5,756,455; each incorporated herein by reference), Retinoblastoma, BRCA1 (U.S. Patent No. 5,750,400; 5,654,155; 5,710,001; 5,756,294; 5,709,999; 5,693,473; 5,753,441; 5,622,829; and 5,747,282; each incorporated herein by reference), MEN-1 (GenBank accession number U93236) and adenovirus E1A (U.S. Patent No. 5,776,743; incorporated herein by reference) genes.

20 Other compositions that may be used include genes encoding the tumor necrosis factor related apoptosis inducing ligand termed TRAIL, and the TRAIL polypeptide (U.S. Patent No. 5,763,223; incorporated herein by reference); the 24 kD apoptosis-associated protease of U.S. Patent No. 5,605,826 (incorporated herein by reference); Fas-associated factor 1, FAF1 (U.S.
25 Patent No. 5,750,653; incorporated herein by reference). Also contemplated for use in these aspects of the present invention is the provision of interleukin-1 β -converting enzyme and family members, which are also reported to stimulate apoptosis.

30 Compounds such as carbostyryl derivatives (U.S. Patent No. 5,672,603; and 5,464,833; each incorporated herein by reference); branched apogenic peptides (U.S. Patent No. 5,591,717;

incorporated herein by reference); phosphotyrosine inhibitors and non-hydrolyzable phosphotyrosine analogs (U.S. Patent No. 5,565,491; and 5,693,627; each incorporated herein by reference); agonists of RXR retinoid receptors (U.S. Patent No. 5,399,586; incorporated herein by reference); and even antioxidants (U.S. Patent No. 5,571,523; incorporated herein by reference) may also be used. Tyrosine kinase inhibitors, such as genistein, may also be linked to ligands that target a cell surface receptor (U.S. Patent No. 5,587,459; incorporated herein by reference).

J4. Immunotoxins and Coaguligands

The naked anti-aminophospholipid antibody treatment methods of the invention may be used in combination with immunotoxins (ITs) and/or coaguligands in which the targeting portion thereof, *e.g.*, antibody or ligand, is directed to a relatively specific marker of the tumor cells, tumor vasculature or tumor stroma. In common with the chemotherapeutic and anti-angiogenic agents discussed above, the use of an anti-aminophospholipid antibody in combination with a targeted toxin or coagulant will generally result in the distinct agents being directed against different targets within the tumor environment. This should result in additive, markedly greater than additive or even synergistic anti-tumor results.

Generally speaking, antibodies or ligands for use in these additional aspects of the invention will preferably recognize accessible tumor antigens that are preferentially, or specifically, expressed in the tumor site. The antibodies or ligands will also preferably exhibit properties of high affinity; and the antibodies, ligands or conjugates thereof, will not exert significant *in vivo* side effects against life-sustaining normal tissues, such as one or more tissues selected from heart, kidney, brain, liver, bone marrow, colon, breast, prostate, thyroid, gall bladder, lung, adrenals, muscle, nerve fibers, pancreas, skin, or other life-sustaining organ or tissue in the human body. The term "significant side effects", as used herein, refers to an antibody, ligand or antibody conjugate, that, when administered *in vivo*, will produce only negligible or clinically manageable side effects, such as those normally encountered during chemotherapy.

At least one binding region of these second anti-cancer agents employed in combination with the anti-aminophospholipid antibodies of the invention will be a component that is capable of delivering a toxin or coagulation factor to the tumor region, *i.e.*, capable of localizing within a tumor site. Such targeting agents may be directed against a component of a tumor cell, tumor vasculature or tumor stroma. The targeting agents will generally bind to a surface-expressed, surface-accessible or surface-localized component of a tumor cell, tumor vasculature or tumor stroma. However, once tumor vasculature and tumor cell destruction begins, internal components will be released, allowing additional targeting of virtually any tumor component.

Many tumor cell antigens have been described, any one which could be employed as a target in connection with the combined aspects of the present invention. Appropriate tumor cell antigens for additional immunotoxin and coaguligand targeting include those recognized by the antibodies B3 (U.S. Patent 5,242,813; incorporated herein by reference; ATCC HB 10573); KSI/4 (U.S. Patent 4,975,369; incorporated herein by reference, obtained from a cell comprising the vectors NRRL B-18356 and/or NRRL B-18357); 260F9 (ATCC HB 8488); and D612 (U.S. Patent 5,183,756; incorporated herein by reference; ATCC HB 9796). One may also consult the ATCC Catalogue of any subsequent year to identify other appropriate cell lines producing anti-tumor cell antibodies.

For tumor vasculature targeting, the targeting antibody or ligand will often bind to a marker expressed by, adsorbed to, induced on or otherwise localized to the intratumoral blood vessels of a vascularized tumor. Appropriate expressed target molecules include, for example, endoglin, E-selectin, P-selectin, VCAM-1, ICAM-1, PSMA (Liu *et al.*, 1997), a TIE, a ligand reactive with LAM-1, a VEGF/VPF receptor, an FGF receptor, $\alpha_v\beta_3$ integrin, pleiotropin and endosialin. Suitable adsorbed targets are those such as VEGF, FGF, TGF β , HGF, PF4, PDGF, TIMP, a ligand that binds to a TIE and tumor-associated fibronectin isoforms. Antigens naturally and artificially inducible by cytokines and coagulants may also be targeted, such as ELAM-1, VCAM-1, ICAM-1, a ligand reactive with LAM-1, endoglin, and even MHC Class II (cytokine-inducible, *e.g.*, by IL-1, TNF- α , IFN- γ , IL-4 and/or TNF- β); and E-selectin, P-selectin, PDGF

and ICAM-1 (coagulant-inducible *e.g.*, by thrombin, Factor IX/IXa, Factor X/Xa and/or plasmin).

5 The following patents and patent applications are specifically incorporated herein by reference for the purposes of even further supplementing the present teachings regarding the preparation and use of immunotoxins directed against expressed, adsorbed, induced or localized markers of tumor vasculature: U.S. Patents Nos. 5,855,866; 5,776,427; 5,863,538; 5,660,827; 5,855,866 and 5,877,289; ^{6,004,555; 6,036,955; and U.S. Application Serial No. 07/846,349} and ~~U.S. Applications Serial No. 07/846,349; 08/295,868 (Patent No. 5,____; Issue Fee paid); 08/350,212 (Patent No. 5,____; Issue Fee paid); 08/457,869 (Notice of Allowance Received); 08/482,369 (U.S. Patent 5,____; Issue Fee paid); 08/487,427 (U.S. Patent 5,____; Issue Fee paid); and 08/479,727 (U.S. Patent 5,____; Issue Fee paid).~~

Suitable tumor stromal targets include components of the tumor extracellular matrix or stroma, or components thereof bound therein; including basement membrane markers, type IV collagen, laminin, heparan sulfate, proteoglycan, fibronectins, activated platelets, LIBS and tenascin. A preferred target for such uses is RIBS.

15 The following patents and patent applications are specifically incorporated herein by reference for the purposes of even further supplementing the present teachings regarding the preparation and use of tumor stromal targeting agents: U.S. Patent No. 5,877,289; ^{6,004,555; 6,036,955; and 6,093,389} and ~~U.S. Applications Serial Nos. 08/482,369 (U.S. Patent 5,____; Issue Fee paid); 08/487,427 (U.S. Patent 5,____; Issue Fee paid); and 08/479,727 (U.S. Patent 5,____; Issue Fee paid).~~

25 In certain applications, the second anti-cancer therapeutics will be antibodies or ligands operatively attached to cytotoxic or otherwise anti-cellular agents having the ability to kill or suppress the growth or cell division of endothelial cells. Suitable anti-cellular agents include chemotherapeutic agents and radioisotopes, as well as cytotoxins. Exemplary chemotherapeutic agents include: steroids; cytokines; anti-metabolites, such as cytosine arabinoside, fluorouracil, methotrexate or aminopterin; anthracyclines; mitomycin C; vinca alkaloids; antibiotics;

demecolcine; etoposide; mithramycin; and anti-tumor alkylating agents, such as chlorambucil or melphalan.

In most combined therapeutic applications, toxin moieties will be preferred, due to the much greater ability of most toxins to deliver a cell killing effect, as compared to other potential agents. Therefore, preferred anti-cellular agents for second therapeutics are plant-, fungus- or bacteria-derived toxins. Exemplary toxins include epipodophyllotoxins; bacterial endotoxin or the lipid A moiety of bacterial endotoxin; ribosome inactivating proteins, such as saporin or gelonin; α -sarcin; aspergillin; restrictocin; ribonucleases, such as placental ribonuclease; diphtheria toxin and *pseudomonas* exotoxin.

Certain preferred toxins are gelonin and the A chain toxins, such as ricin A chain. The most preferred toxin moiety is often ricin A chain that has been treated to modify or remove carbohydrate residues, so called "deglycosylated A chain" (dgA). Deglycosylated ricin A chain is preferred because of its extreme potency, longer half-life, and because it is economically feasible to manufacture it a clinical grade and scale. Recombinant and/or truncated ricin A chain may also be used.

Other agents for use with immunoconjugates for targeting tumor vasculature or tumor stroma are the angiopoietins. The angiopoietins, like the members of the VEGF family, are growth factors largely specific for vascular endothelium (Davis and Yancopoulos, 1999; Holash *et al.*, 1999; incorporated herein by reference). The angiopoietins first described were a naturally occurring agonist, angiopoietin-1 (Ang-1; SEQ ID NO:1 and SEQ ID NO:2), and a naturally occurring antagonist, angiopoietin-2 (Ang-2; SEQ ID NO:3 and SEQ ID NO:4), both of which act by means of the endothelial cell tyrosine kinase receptor, Tie2.

Two new angiopoietins, angiopoietin-3 (mouse) and angiopoietin-4 (human) have also been identified (Valenzuela *et al.*, 1999). Angiopoietin-3 appears to act as an antagonist, whereas angiopoietin-4 appears to function as an agonist (Valenzuela *et al.*, 1999). A protein termed angiopoietin-3 was also cloned from human heart and reported not to have mitogenic

effects on endothelial cells (Kim *et al.*, 1999). Fusion proteins of angiopoietin-1 and angiopoietin-2 have also been created, as exemplified by the stable Ang-1-Ang-2 fusion protein included herein as SEQ ID NO:5.

5 Whereas VEGF is necessary for the early stages of vascular development, angiopoietin-1 is generally required for the later stages of vascular remodeling. Angiopoietin-1 is thus a maturation or stabilization factor, which converts immature vessels to mature vessels.

10 Angiopoietin-1 has been shown to augment revascularization in ischemic tissue (Shyu *et al.*, 1998) and to increase the survival of vascular networks exposed to either VEGF or a form of aFGF (Papapetropoulos *et al.*, 1999). These authors also showed that angiopoietin-1 prevents apoptotic death in HUVEC triggered by withdrawal of the same form of aFGF (Papapetropoulos *et al.*, 1999). Such data are consistent with the direct role of angiopoietin-1 on human endothelial cells and its interaction with other angiogenic molecules to stabilize vascular structures by promoting the survival of differentiated endothelial cells.

15 Angiopoietin-2 is a preferred agent for use in targeted combination therapy, particularly in tumors with low VEGF levels and/or in combination with VEGF inhibition. Angiopoietin-2 is also a ligand for Tie2, but generally counteracts blood vessel maturation/stability mediated by angiopoietin-1. It is thus an antagonist of angiopoietin-1, and acts to disturb capillary structure. However, as angiopoietin-2 renders endothelial cells responsive to angiogenic stimuli, it can initiate neovascularization in combination with other appropriate signals, particularly VEGF (Asahara *et al.*, 1998; Holash *et al.*, 1999; incorporated herein by reference).

25 In the absence of another angiogenic signal, angiopoietin-2 causes vessels to destabilize and become immature. In the presence of a stimulus, such as VEGF, angiopoietin-2 promotes angiogenesis. Indeed, the angiogenic effects of a number of regulators are believed to be achieved, at least in part, through the regulation of an autocrine loop of angiopoietin-2 activity in microvascular endothelial cells (Mandriota and Pepper, 1998).

30

Angiopoietin-2 expression in tumor tissue has been reported (Tanaka *et al.*, 1999), where it presumably acts in combination with VEGF to promote angiogenesis (Stratmann *et al.*, 1998; Holash *et al.*, 1999). However, as angiopoietin-2 provides a negative signal when VEGF is low or absent, provision of angiopoietin-2 can be a useful therapeutic approach. Angiopoietin-2 can be administered as a protein therapeutic or via gene therapy (see above), or in a tumor-targeted form. Although all types of targeted angiopoietin-2 constructs are envisioned for use in the combined therapy aspects of the invention, currently preferred agents for targeting angiopoietin-2 to the tumor are those that bind to aminophospholipids, including anti-PS antibodies and annexins.

10

For tumor targeting and treatment with immunotoxins, the following patents and patent applications are specifically incorporated herein by reference for the purposes of even further supplementing the present teachings regarding anticellular and cytotoxic agents: U.S. Patents Nos. 5,855,866; 5,776,427; 5,863,538; and 5,660,827; and ~~U.S. Applications Serial Nos. 07/846,349; 08/295,868 (Patent No. 5,___,___; Issue Fee paid); 08/330,212 (Patent No. 5,___,___; Issue Fee paid); and 08/457,869 (Notice of Allowance Received).~~ *6,004,557; 5,965,132 and 6,051,230; and U.S. Application Serial No. 107,846,349*

The second, targeted agent for optional use with the anti-aminophospholipid antibodies of the invention may also comprise a targeted component that is capable of promoting coagulation, *i.e.*, a "coaguligand". Here, the targeting antibody or ligand may be directly or indirectly, *e.g.*, via another antibody, linked to a factor that directly or indirectly stimulates coagulation.

Preferred coagulation factors for such uses are Tissue Factor (TF) and TF derivatives, such as truncated TF (tTF), dimeric and multimeric TF, and mutant TF deficient in the ability to activate Factor VII. Other suitable coagulation factors include vitamin K-dependent coagulants, such as Factor II/IIa, Factor VII/VIIa, Factor IX/IXa and Factor X/Xa; vitamin K-dependent coagulation factors that lack the Gla modification; Russell's viper venom Factor X activator; platelet-activating compounds, such as thromboxane A₂ and thromboxane A₂ synthase; and inhibitors of fibrinolysis, such as α 2-antiplasmin.

30

Tumor targeting and treatment with coaguligands is described in the following patents and patent applications, each of which are specifically incorporated herein by reference for the purposes of even further supplementing the present teachings regarding coaguligands and coagulation factors: U.S. Patent Nos. 5,855,866 and 5,877,289; ~~U.S. Applications Serial Nos. 07/846,349; 08/350,212 (Patent No. 5, , , ; Issue Fee paid); 08/482,369 (U.S. Patent 5, , , ; Issue Fee paid); 08/487,427 (U.S. Patent 5, , , ; Issue Fee paid); and 08/479,727 (U.S. Patent 5, , , ; Issue Fee paid).~~

As somewhat wider distribution of a coagulating agent will not be associated with severe side effects, there is a less stringent requirement imposed on the targeting element of coaguligands than with immunotoxins. Therefore, to achieve specific targeting means that coagulation is promoted in the tumor vasculature relative to the vasculature in non-tumor sites. Thus, specific targeting of a coaguligand is a functional term, rather than a purely physical term relating to the biodistribution properties of the targeting agent. It is not unlikely that useful targets may be not be entirely tumor-restricted, and that targeting ligands that are effective to promote tumor-specific coagulation may nevertheless be safely found at other sites of the body following administration, as occurs with VCAM-1.

The preparation of immunotoxins is generally well known in the art (see, e.g., U.S. Patent 4,340,535, incorporated herein by reference). Each of the following patents and patent applications are further incorporated herein by reference for the purposes of even further supplementing the present teachings regarding immunotoxin generation, purification and use: U.S. Patents Nos. 5,855,866; 5,776,427; 5,863,538; and 5,660,827; ~~and U.S. Applications Serial Nos. 07/846,349; 08/295,868 (Patent No. 5, , , ; Issue Fee paid); 08/350,212 (Patent No. 5, , , ; Issue Fee paid); and 08/457,869 (Notice of Allowance Received).~~

In the preparation of immunotoxins, advantages may be achieved through the use of certain linkers. For example, linkers that contain a disulfide bond that is sterically "hindered" are often preferred, due to their greater stability *in vivo*, thus preventing release of the toxin moiety prior to binding at the site of action. It is generally desired to have a conjugate that will remain

intact under conditions found everywhere in the body except the intended site of action, at which point it is desirable that the conjugate have good "release" characteristics.

Depending on the specific toxin compound used, it may be necessary to provide a peptide spacer operatively attaching the targeting agent and the toxin compound, wherein the peptide spacer is capable of folding into a disulfide-bonded loop structure. Proteolytic cleavage within the loop would then yield a heterodimeric polypeptide wherein the targeting agent and the toxin compound are linked by only a single disulfide bond.

When certain other toxin compounds are utilized, a non-cleavable peptide spacer may be provided to operatively attach the targeting agent and the toxin compound. Toxins that may be used in conjunction with non-cleavable peptide spacers are those that may, themselves, be converted by proteolytic cleavage, into a cytotoxic disulfide-bonded form. An example of such a toxin compound is a *Pseudomonas* exotoxin compound.

A variety of chemotherapeutic and other pharmacological agents can also be successfully conjugated to antibodies or targeting ligands. Exemplary antineoplastic agents that have been conjugated to antibodies include doxorubicin, daunomycin, methotrexate and vinblastine. Moreover, the attachment of other agents such as neocarzinostatin, macromycin, trenimon and α -amanitin has been described (see U.S. Patent No. 5,855,866; and Patent No. 5,965,132 (Application Serial No. 08/350,212, Issue Fee paid) and references incorporated therein.

In light of one of the present inventors earlier work, the preparation of coaguligands is now also easily practiced. The operable association of one or more coagulation factors with a targeting agent may be a direct linkage, such as those described above for the immunotoxins. Alternatively, the operative association may be an indirect attachment, such as where the targeting agent is operatively attached to a second binding region, preferably an antibody or antigen binding region of an antibody, that binds to the coagulation factor. The coagulation factor should be attached to the targeting agent at a site distinct from its functional coagulating site, particularly where a covalent linkage is used to join the molecules.

Indirectly linked coaguligands are most often based upon bispecific antibodies. The preparation of bispecific antibodies is also well known in the art. One preparative method involves the separate preparation of antibodies having specificity for the targeted tumor component, on the one hand, and the coagulating agent on the other. Peptic F(ab')₂ fragments from the two chosen antibodies are then generated, followed by reduction of each to provide separate Fab'γ_{SH} fragments. The SH groups on one of the two partners to be coupled are then alkylated with a cross-linking reagent, such as o-phenylenedimaleimide, to provide free maleimide groups on one partner. This partner may then be conjugated to the other by means of a thioether linkage, to give the desired F(ab'γ)₂ heteroconjugate (Glennie *et al.*, 1987; incorporated herein by reference). Other approaches, such as cross-linking with SPDP or protein A may also be carried out.

Another method for producing bispecific antibodies is by the fusion of two hybridomas to form a quadroma. As used herein, the term "quadroma" is used to describe the productive fusion of two B cell hybridomas. Using now standard techniques, two antibody producing hybridomas are fused to give daughter cells, and those cells that have maintained the expression of both sets of clonotype immunoglobulin genes are then selected.

A preferred method of generating a quadroma involves the selection of an enzyme deficient mutant of at least one of the parental hybridomas. This first mutant hybridoma cell line is then fused to cells of a second hybridoma that had been lethally exposed, *e.g.*, to iodoacetamide, precluding its continued survival. Cell fusion allows for the rescue of the first hybridoma by acquiring the gene for its enzyme deficiency from the lethally treated hybridoma, and the rescue of the second hybridoma through fusion to the first hybridoma. Preferred, but not required, is the fusion of immunoglobulins of the same isotype, but of a different subclass. A mixed subclass antibody permits the use if an alternative assay for the isolation of a preferred quadroma.

Microtiter identification embodiments, FACS, immunofluorescence staining, idiotype specific antibodies, antigen binding competition assays, and other methods common in the art of antibody characterization may be used to identify preferred quadromas. Following the isolation of the quadroma, the bispecific antibodies are purified away from other cell products. This may
5 be accomplished by a variety of antibody isolation procedures, known to those skilled in the art of immunoglobulin purification (see, *e.g.*, Antibodies: A Laboratory Manual, 1988; incorporated herein by reference). Protein A or protein G sepharose columns are preferred.

In the preparation of both immunotoxins and coaguligands, recombinant expression may
10 be employed. The nucleic acid sequences encoding the chosen targeting agent, and toxin or coagulant, are attached in-frame in an expression vector. Recombinant expression thus results in translation of the nucleic acid to yield the desired targeting agent-toxin/coagulant compound.

The following patents and patent applications are each incorporated herein by reference
15 for the purposes of even further supplementing the present teachings regarding coaguligand preparation, purification and use, including bispecific antibody coaguligands: U.S. Patent Nos. 5,855,866 and 5,877,289; ~~U.S. Applications Serial Nos. 07/846,349; 08/350,212 (Patent No. 5,855,866; Issue Fee paid); 08/273,567; 08/482,369; 08/485,482; and 08/472,631 (U.S. Patent 5,855,866; Issue Fee paid); 08/487,427 (U.S. Patent 5,855,866; Issue Fee paid); and 08/479,727 and 08/481,904 (U.S. Patent 5,855,866; Issue Fee paid).~~
5,965,132; 6,004,555; 6,036,985 and 6,093,395 and U.S. Applications Serial Nos. 07/846,349; 08/350,212; 08/273,567; 08/482,369; 08/485,482; and 08/472,631

Effective doses of immunotoxins and coaguligands for combined use with the naked anti-aminophospholipid antibodies in the treatment of cancer will be between about 0.1 mg/kg and about 2 mg/kg, and preferably, of between about 0.8 mg/kg and about 1.2 mg/kg, when
25 administered via the IV route at a frequency of about 1 time per week. Some variation in dosage will necessarily occur depending on the condition of the subject being treated. The physician responsible for administration will determine the appropriate dose for the individual subject.

* * *

The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

10

EXAMPLE I

VCAM-1 Expression on Tumor and Normal Blood Vessels

A. Materials and Methods

1. Materials

Na¹²⁵I was obtained from Amersham (Arlington Heights, IL). Dulbecco's modified Eagle's tissue culture medium (DMEM) and Dulbecco PBS containing Ca²⁺ and Mg²⁺ were obtained from Gibco (Grand Island, NY). Fetal calf serum was obtained from Hyclone (Logan, Utah). O-phenylenediamine, hydrogen peroxide, 3-aminopropyltriethoxy-silane and sterile, endotoxin-free saline (0.9% NaCl in 100 ml of water) were from Sigma (St. Louis, MO). SMPT was from Pierce (Rockford, IL). Proplex T containing factor VII (74 IU/ml), factor X and factor IX (17 IU/ml) was purchased from Baxter Diagnostics Inc. (McGraw Park, IL). Chromogenic substrate, S-2765, for measuring factor Xa proteolytic activity was obtained from Chromogenix (Franklin, OH). Purified factor Xa was purchased from American Diagnostica (Greenwich, CT). 96 and 48 flat bottom microtiter plates were obtained from Falcon (Becton Dickinson and Co., Lincoln Park, NJ). Sepharose-Protein G beads and S200 Superdex were purchased from Pharmacia (Piscataway, NJ). Recombinant murine IL-1 α was purchased from R&D Systems (Minneapolis, MN).

2. Antibodies

The MK2.7 hybridoma, secreting a rat IgG1 antibody against murine VCAM-1, was obtained from the American Type Culture Collection (ATCC, Rockville, MD; ATCC CRL

1909). The characterization of this anti-VCAM-1 antibody has been reported by Miyake *et al.* (1991, incorporated herein by reference). The R187 hybridoma, secreting a rat IgG1 antibody against murine viral protein p30 gag, was also obtained from the ATCC, and was used as an isotype matched control for the anti-VCAM-1 antibody.

5

Mouse monoclonal antibody, 10H10, against human tissue factor was prepared as described in Morrissey *et al.* (1988), and in U.S. application Serial No. 08/482,369, each incorporated herein by reference.

10

MECA 32, a pan anti-mouse vascular endothelial cell antibody, was prepared as described by Leppink *et al.* (1989, incorporated herein by reference). MJ 7/18 rat IgG, reactive with murine endoglin, was prepared as described by Ge and Butcher (1994, incorporated herein by reference). The MECA 32 and MJ 7/18 antibodies served as positive controls for immunohistochemical studies.

15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995

Rabbit anti-rat and rat anti-mouse secondary antibodies conjugated with horseradish peroxidase (HRP) were purchased from Dako (Carpinteria, CA).

3. Antibody Purification

Anti-VCAM-1 hybridoma, MK 2.7, and the irrelevant control hybridoma, R187, were grown in bioreactors (Heraeus, Inc., Germany) for 12 days. Supernatants were centrifuged, filtered through 0.22 μ m filters and loaded onto Sepharose-Protein G columns. IgG was eluted with citric acid buffer, pH 3.5, dialyzed into PBS and stored thereafter at 4°C in the same buffer. Purity was estimated by SDS-PAGE and was routinely > 90%. Binding capacity of the purified anti-VCAM-1 antibody was assessed immunohistochemically on frozen sections of L540 tumor and by cell-based ELISA using IL-1 α stimulated bEnd.3 cells, as described herein below.

4. Tumor-Bearing Mice and Immunohistochemistry

Male CB17 SCID mice (Charles River, Wilmington, MA) weighing approximately 25 g were injected with 1×10^7 L540 Hodgkin's lymphoma cells subcutaneously into the right flank.

Tumors were allowed to grow to a size of 0.4-0.7 cm³. Animals were anesthetized with metaferane and their blood circulation was perfused with heparinized saline as described by Burrows *et al.* (1992, incorporated herein by reference). The tumor and major organs were removed and snap-frozen in liquid nitrogen.

5

Cryostat sections of the tissues were cut, incubated with the anti-VCAM-1 antibody and stained immunohistochemically to detect VCAM-1. Rat IgG was detected using rabbit anti-rat IgG conjugated to HRP followed by development with carbazole (Fries *et al.*, 1993).

10 B. Results

The blood vessels of the major organs and a tumor from mice bearing subcutaneous L540 human Hodgkin's tumors were examined immunohistochemically for VCAM-1 expression using an anti-VCAM-1 antibody. VCAM-1 expression on tumor blood vessels was more peripheral than central. However, as demonstrated in Example VI and Example VII, the anti-VCAM-1 antibody and coaguligand were evidently binding to blood transporting vessels, as clearly shown by the ability of the coaguligand to arrest blood flow in all tumor regions and to cause destruction of the intratumoral region.

Overall, VCAM-1 expression was observed on 20-30% of total tumor blood vessels stained by the anti-endoglin antibody, MJ 7/18. VCAM-1 staining of the tumor vessels was largely observed on venules. VCAM-1 expression was similar in tumors up to 1500 mm³, but larger tumors appeared to have reduced staining, with 5-10% of MJ 7/18 positive vessels being positive for VCAM-1.

Constitutive vascular expression of VCAM-1 was found in heart and lungs in both tumor-bearing and normal animals (Table 1). In the heart, strong staining was observed on venules and veins. Approximately 10% of MECA 32 positive vessels were positive for VCAM-1. Staining in lung endothelium was weak in comparison to heart and tumor, and was confined to a few large blood vessels. Strong stromal staining was observed in testis where VCAM-1 expression was

strictly extravascular. Similar findings regarding constitutive VCAM-1 expression in rodent lung and testis were previously reported (Fries *et al.*, 1993).

5 **TABLE 1**
Expression of VCAM-1 on Endothelium in Tissues of L540 Tumor Bearing Mice and
Localization of Anti-VCAM-1 Antibody

| Tissue | VCAM-1 expression ^a | anti-VCAM-1 antibody localization ^b |
|----------------------|--------------------------------|--|
| Adrenal | - ^c | - |
| Brain Cerebellum | - | - |
| Brain Cortex | - | - |
| Heart | ++ | ++ |
| Kidney | - | - |
| Large Intestine | - | - |
| Liver | - | - |
| Lung | + | + |
| Pancreas | - | - |
| Small Intestine | - | - |
| Spleen | - | - |
| Testis | - ^d | - |
| L540 Hodgkin's tumor | +++ | +++ |

^a VCAM-1 was detected by anti-VCAM-1 antibody followed by anti-rat IgG-HRP.

^b Localization of anti-VCAM-1 antibody *in vivo* was determined by injecting the antibody, exsanguinating the mice and staining tissues staining with anti-rat IgG-HRP.

10 ^c Intensity of staining was compared to pan-endothelial markers MJ 7/18 and MECA 32; - no staining; + weak; ++ moderate; +++ strong.

^d No vascular expression was observed; however, extravascular stroma of testis was stained by anti-VCAM-1 antibody.

EXAMPLE II

Localization of Anti-VCAM-1 Antibody *In Vivo*

A. Methods

Male CB17 SCID mice (Charles River, Wilmington, MA) weighing approximately 25 g
5 were injected with 1×10^7 L540 Hodgkin's lymphoma cells subcutaneously into the right flank.
Tumors were allowed to grow to a size of 0.4-0.7 cm³.

Mice were injected intravenously with 30 µg/25 g body weight of anti-VCAM-1
antibody, R187 antibody or corresponding coaguligands in 200 µl of saline. Two hours later,
10 animals were anesthetized with metaferne and their blood circulation was perfused with
heparinized saline as described (Burrows *et al.*, 1992; incorporated herein by reference). The
tumor and major organs were removed and snap-frozen in liquid nitrogen.

Cryostat sections of the tissues were cut and were stained immunohistochemically for the
15 presence of rat IgG or TF. Rat IgG was detected using rabbit anti-rat IgG conjugated to HRP
followed by development with carbazole (Fries *et al.*, 1993). Coaguligand was detected using
the 10H10 antibody that recognizes human tissue factor, followed by HRP-labeled anti-mouse
IgG. 10H10 antibody does not cross-react detectably with murine tissue factor (Morrissey *et al.*,
1988, incorporated herein by reference) or other murine proteins.

B. Results

Mice bearing subcutaneous L540 tumors were injected intravenously with anti-VCAM-1
antibody and, two hours later, the mice were exsanguinated. The tumor and normal organs were
removed and frozen sections were prepared and examined immunohistochemically to determine
25 the location of the antibody. Serial sections of the tissues were examined. Localized rat IgG was
detected by HRP-labeled anti-rat Ig; and murine blood vessels were identified by pan-endothelial
antibody, MECA 32.

Anti-VCAM-1 antibody was detected on endothelium of tumor, heart and lung (Table 1).
30 The intensity and number of stained vessels was identical to that on serial sections of the same

tissues stained directly with anti-VCAM-1 antibody (Table 1). Staining was specific as no staining of endothelium was observed in the tumor and organs of mice injected with a species isotype matched antibody of irrelevant specificity, R187. No localization of anti-VCAM-1 antibody was found in testis or any normal organ except heart and lung.

5

EXAMPLE III

Preparation of Anti-VCAM-1•tTF Coaguligand

An anti-VCAM-1•tTF conjugate or "coaguligand" was prepared as follows. Truncated tissue factor (tTF), with an additional added cysteine introduced at N-terminus (U.S. application
10 Serial No. 08/482,369, incorporated herein by reference), was expressed in *E. coli* and purified as described by Stone *et al.* (1995, incorporated herein by reference). After purification, the sulfhydryl group of N' cysteine-tTF was protected by reaction with Ellman's reagent. The tTF derivative was stored in small volumes at -70°C.

0031362-07129

15 To prepare the anti-VCAM-1 coaguligand, 5 ml of anti-VCAM-1 antibody IgG (2 mg/ml) in PBS were mixed with 36 µl of SMPT (10 mM) dissolved in dry DMF and incubated at room temperature for 1 h. The mixture was filtered through a column of Sephadex G25 equilibrated in PBS containing 1 mM EDTA. The fractions containing the SMPT-derivatized antibody were concentrated to 4 ml by ultrafiltration in an Amicon cell equipped with a 10,000 Da cut-off filter.
20 Freshly thawed tTF derivative was incubated with 30 µl of DTT (10 mM) in H₂O for 10 min. at room temperature and was filtered through a column of Sephadex G25 equilibrated in PBS containing 1 mM EDTA. The eluted fractions containing reduced tTF were concentrated by ultrafiltration under nitrogen to a final volume of 3 ml.

25 The reduced tTF was mixed with the SMPT-derivatized antibody and the mixture was allowed to react for 24 h at room temperature. At the end of the incubation, the reaction mixture was resolved by gel filtration on a column of Superdex S200 equilibrated in PBS. Fractions containing anti-VCAM-1•tTF having a M_r of 180,000 and corresponding to one molecule of antibody linked to one molecule of tTF were collected.

30

EXAMPLE IV

Binding of Anti-VCAM-1 Coaguligand to Activated Endothelial Cells

A. Methods

1. Iodination of 10H10 antibody

5 Anti-human tissue factor antibody, 10H10, was radiolabeled with ^{125}I using Chloramine T as described by Bocci (1964, incorporated herein by reference). The specific activity was approximately 10,000 cpm/ μg , as calculated from protein determinations measured by a Bradford assay (Bradford, 1976).

10 2. Cells

L540 Hodgkin cells (L540 Cy), derived from a patient with end-stage disease, were prepared as described in Diehl *et al.* (1985, incorporated herein by reference), and were obtained from Prof. Volker Diehl (Klinik für Innere Medizin der Universität, Köln, Germany). bEnd.3 cells (murine brain endothelioma) were prepared as described in Bussolino *et al.* (1991) and
15 Montesano *et al.* (1990), each incorporated herein by reference, and were obtained from Prof. Werner Risau (Max Planck Institute, Bad Nauheim, Germany).

3. Tissue Culture

bEnd.3 cells and hybridomas were maintained in DMEM supplemented with 10% fetal calf serum, 2 mM L-glutamine, 2 units/ml penicillin G and 2 $\mu\text{g}/\text{ml}$ streptomycin. L540 cells were maintained in RPMI 1640 containing the same additives. All cells were subcultured once a week. bEnd.3 trypsinization was performed using 0.125% trypsin in PBS solution containing 0.2% EDTA. For binding studies, cells were seeded at a density of 5×10^4 cells/ml in 0.5 ml of medium in 48 well plates and incubated for 48-96 h. Medium was refreshed 24 h before each
25 study.

4. Binding of Coaguligand to Activated Endothelial Cells

Binding of the anti-VCAM-1 antibody and coaguligand to VCAM-1 on activated bEnd.3 cells was determined using a cell based ELISA, as described by Hahne (1993, incorporated
30 herein by reference). bEnd.3 cells were incubated with 10 units/ml of IL-1 α for 4 h at 37°C in

96-well microtiter plates. At the end of this incubation, medium was replaced by DPBS containing 2 mM Ca^{2+} and Mg^{2+} and 0.2% (w/v) gelatin as a carrier protein. The same buffer was used for dilution of antibodies and for washing of cell monolayers between steps.

- 5 Cells were incubated with 4 $\mu\text{g}/\text{ml}$ of anti-VCAM-1•tTF conjugate, anti-VCAM•1 antibody or control reagents for 2 h, and were then washed and incubated for 1 h with rabbit anti-rat IgG-HRP conjugate (1:500 dilution). All steps were performed at room temperature. HRP activity was measured by adding O-phenylenediamine (0.5 mg/ml) and hydrogen peroxide (0.03% w/v) in citrate-phosphate buffer, pH 5.5. After 30 min., 100 μl of supernatant were
10 transferred to 96 well plates, 100 μl of 0.18 M H_2SO_4 were added and the absorbance was measured at 492 nm. Each study was performed in duplicate and repeated at least twice.

5. Detection of Coaguligand Bound to Endothelial cells

Anti-VCAM-1 coaguligand and appropriate controls were incubated with IL-1 α stimulated bEnd.3 cells in 96-well microtiter plates, as described above. Bound coaguligands were detected by identifying both the tissue factor component and the rat IgG component bound to bEnd.3 cells.

After removing the excess of unbound antibody, cells were incubated with 100 $\mu\text{l}/\text{well}$ of ^{125}I -labeled 10H10 antibody (0.2 $\mu\text{g}/\text{ml}$) or ^{125}I -labeled rabbit anti-rat Ig (0.2 $\mu\text{g}/\text{ml}$) in binding buffer. After 2 h incubation at room temperature, cells were washed extensively and dissolved in 0.5 M of NaOH. The entire volume of 0.5 ml was transferred to plastic tubes and counted in a γ counter. Each study was performed in duplicate and repeated at least twice.

25 B. Results

The ability of an anti-VCAM-1•tTF coaguligand to bind to IL-1 α activated murine bEnd.3 cells was determined by measuring the binding of radioiodinated anti-TF antibody to coaguligand-treated cells *in vitro*. VCAM-1 expression by bEnd.3 cells is transiently inducible by IL-1 α with a peak of VCAM-1 expression being obtained 4-6 h after addition of the cytokine

Chromogenic substrate S2765 for measuring factor Xa proteolytic activity was added in 50 μ l, giving a final concentration of 300 μ M. The breakdown of the substrate was determined by reading the absorbance at 405 nm over a 2 h period in a microplate reader (Molecular Devices, Palo Alto, CA).

5

Production of the chromogenic product was completely dependent on the presence of Proplex T and bEnd.3 cells preincubated with the specific coaguligand. Background hydrolysis of the substrate by Proplex T in the absence of cells was approximately 10% of the maximal value and was subtracted from each determination. Free coaguligands diluted in Proplex T solution were unable to generate factor Xa. The amount of Xa generated was calculated by reference to a standard curve constructed with known concentrations of purified factor Xa.

10
15

At the end of the study, cells were detached with trypsin-EDTA and counted. The results are expressed as the amount of factor Xa generated per 10^4 cells. Each study was performed in duplicate and was repeated at least 3 times.

B. Results

1. Factor X Activation

Anti-VCAM-1•tTF coaguligand bound to IL-1 α -activated bEnd.3 cells was capable of specifically activating factor X. The rate of generation of factor Xa by anti-VCAM-1•tTF coated cells was 3.2 ng per 10^4 cells per hour, which is 7-10 fold higher than was observed with activated cells treated with a control coaguligand of irrelevant specificity or with tTF alone (FIG. 1B). Anti-VCAM-1•tTF in the absence of cells had undetectable factor X activating activity, confirming that cell binding is essential for coaguligand activity.

20
25

Anti-VCAM-1•tTF bound to unstimulated bEnd.3 cells activated factor X at a rate of 1.6 ng per 10^4 cells per hour. This rate is about half that observed with the IL-1 α -stimulated cells, in accordance with the 50% lower amount of coaguligand that binds to unstimulated as compared with stimulated cells. Similar results to those shown in FIG. 1B were obtained in three separate studies.

30

2. Effect of Endothelial Cell Permeabilization

Permeabilization of bEnd.3 monolayers with saponin after treating them with anti-VCAM-1•tTF coaguligand increased the ability of the bound coaguligand to activate factor X by about 30-fold (Table 2). The rate of factor Xa generation by unstimulated cells treated with anti-VCAM-1•tTF increased from 1.6 to 49.2 ng per 10^4 cells per hour after permeabilization, while that of IL-1 α stimulated cells increased from 3.2 to 98.8 ng per 10^4 cells per hour. The factor Xa generating activity of the permeabilized cells was due to the bound coaguligand rather than to endogenous TF since permeabilized untreated cells or cells treated with control coaguligand of irrelevant specificity had low factor Xa generating activity (2 ng per 10^4 cells per hour).

These results indicate that the coaguligand is able to function more efficiently in the environment of a permeabilized cell. Possibly, permeabilization exposes negatively-charged phospholipids from within the cell that accelerate the formation of the coagulation-initiation complexes, or else prevents the inactivation of such complexes by TFPI.

TABLE 2

Generation of Factor Xa by Anti-VCAM-1•tTF Bound to Intact or Permeabilized

bEnd.3 cells (ng per 10⁴ cells per 60 min.)

| Treatment ^a | Intact cells | | Permeabilized cells ^b | |
|------------------------|-------------------|---------------|----------------------------------|---------------|
| | Control | IL-1 α | Control | IL-1 α |
| Buffer | 0.25 ^c | 0.43 | 0.45 | 2.0 |
| tTF | 0.26 | 0.42 | 0.39 | 2.1 |
| Control IgG•tTF | 0.26 | 0.43 | 0.41 | 2.1 |
| Anti-VCAM-1•tTF | 1.64 | 3.17 | 49.2 | 98.8 |

^a IL-1 α stimulated and unstimulated bEnd.3 cells were incubated with buffer alone or with 4 μ g/ml of tTF, control IgG•tTF or anti-VCAM-1•tTF followed by 60 min. incubation with Proplex T solution at 37°C.

^b Cells were treated with 0.2% saponin 5 min. before addition of Proplex T.

^c Amount of factor Xa was determined as described above. Results are expressed as ng of factor Xa generated per 10⁴ cells per 60 min. The arithmetic mean values from triplicate wells are shown. SE were less than 5 percent of the mean values.

EXAMPLE VI

Tumor Blood Vessel Thrombosis by Anti-VCAM-1 Coaguligand

A. Methods

SCID mice bearing L540 tumors (0.4-0.7 cm³) were injected intravenously with 40 μ g (total protein) of anti-VCAM-1•tTF or R187•tTF. This dose corresponds to 32 μ g of antibody and 8 μ g of tTF. Other animals received equivalent quantities of free antibody, free tTF or a mixture of both. Animals were anesthetized 4 or 24 h later and their blood circulations were perfused with heparinized saline. The tumor and major organs were removed and were fixed in formalin and paraffin-embedded or snap-frozen for cryosectioning. Sections were cut through the center of the tissue or tumor. The number of thrombosed and non-thrombosed blood vessels in 5 cross-sections were counted. The percentage of thrombosed vessels was calculated.

B. Results

1. Thrombosis of Tumor Blood Vessels

This study shows that intravenous administration of the anti-VCAM-1-tTF coaguligand induces selective thrombosis of tumor blood vessels, as opposed to vessels in normal tissues, in tumor-bearing mice.

The anti-VCAM-1-tTF coaguligand was administered to mice bearing subcutaneous L540 tumors of 0.4 to 0.6 cm in diameter. Before coaguligand injection, tumors were healthy, having a uniform morphology lacking regions of necrosis. The tumors were well vascularized and had a complete absence of spontaneously thrombosed vessels or hemorrhages. Within four hours of coaguligand injection, 40-70% of blood vessels were thrombosed, despite the initial staining of only 20-30% of tumor blood vessels shown in Example I. The thrombosed vessels contained occlusive platelet aggregates, packed erythrocytes and fibrin. In several regions, the blood vessels had ruptured, spilling erythrocytes into the tumor interstitium.

By 24 h after coaguligand injection, the blood vessels were still occluded and extensive hemorrhage had spread throughout the tumor. Tumor cells had separated from one another, had pyknotic nuclei and were undergoing cytolysis. By 72 h, advanced necrosis was evident throughout the tumor. Necrosis was clearly present in the intratumoral region of the tumor, where VCAM-1 expression on the vessels was not originally prominent. The coaguligand binding was evidently effective to curtail blood flow in all tumor regions, resulting in widespread tumor destruction. Furthermore, it is likely that the initial coaguligand-induced thrombin deposition results in increased induction of the VCAM-1 target antigen on central vessels, thus amplifying targeting and tumor destruction.

The thrombotic action of anti-VCAM-1-tTF on tumor vessels was antigen specific. None of the control reagents administered at equivalent quantities (tTF alone, anti-VCAM-1 antibody alone, tTF plus anti-VCAM-1 antibody or the control coaguligand of irrelevant specificity) caused thrombosis (Table 3).

5
10
15
20

TABLE 3
Anti-VCAM-1•tTF-Mediated Thrombosis in L540 Tumor Bearing Mice

| Treatment ^a | Thrombosed Vessels (%) ^b | | |
|---|-------------------------------------|----------------|--------------|
| | L540 Tumor | Heart and Lung | Other Organs |
| Saline | 0-2 | 0 | 0 |
| TTF | 0-2 | 0 | 0 |
| Anti-VCAM-1 Antibody | 0-2 | 0 | 0 |
| Anti-VCAM-1 Antibody + tTF | 0-2 | 0 | 0 |
| Control IgG•tTF | 0-2 | 0 | 0 |
| Anti-VCAM-1•tTF (< 0.3 cm ³) ^c | 0-10 | 0 | 0 |
| Anti-VCAM-1•tTF (> 0.3 cm ³) | 40-70 | 0 | 0 |

^a L540 tumor-bearing mice were injected i.v. with one of the following reagents: saline; 8 µg of unconjugated tTF; 32 µg of unconjugated anti-VCAM-1 antibody; mixture of 8 µg of tTF and 32 µg of anti-VCAM-1 antibody; 40 µg of control IgG•tTF coaguligand; or 40 µg of anti-VCAM-1•tTF coaguligand. Animals were sacrificed 4 h after injection. Tissues were removed and fixed in formalin.

^b Histological quantification was performed by counting numbers of thrombosed blood vessels in 5 cross sections of tissue. The number of thrombosed vessels is expressed as a percentage of total vessels. The range of results from three mice is given.

^c L540 tumor bearing mice were divided into two groups (5-8 animals per group) having tumors smaller or larger than 0.3 cm³.

2. Lack of Thrombosis of Normal Blood Vessels

In addition to the thrombosis of tumor blood vessels, this study also shows that intravenous administration of the anti-VCAM-1•tTF coaguligand does not induce thrombosis of blood vessels in normal organs.

Despite expression of VCAM-1 on vessels in the heart and lung of normal or L540 tumor-bearing mice (Table 1), thrombosis did not occur after anti-VCAM-1•tTF coaguligand

administration. No signs of thrombosis, tissue damage or altered morphology were seen in 25 mice injected with 5 to 45 μg of coaguligand 4 or 24 h earlier. There was a normal histological appearance of the heart and lung from the same mouse that had major tumor thrombosis. All other major organs (brain, liver, kidney, spleen, pancreas, intestine, testis) also had unaltered morphology.

Frozen sections of organs and tumors from coaguligand-treated mice gave coincident staining patterns when developed with either the anti-TF antibody, 10H10, or an anti-rat IgG antibody and confirmed that the coaguligand had localized to vessels in the heart, lung and tumor. The intensity of staining was equal to that seen when coaguligand was applied directly to the sections at high concentrations followed by development with anti-TF or anti-rat IgG, indicating that saturation of binding had been attained *in vivo*.

These studies show that binding of coaguligand to VCAM-1 on normal vasculature in heart and lung is not sufficient to induce thrombosis, and that tumor vasculature provides additional factors to support coagulation.

EXAMPLE VII

In Vivo Tumor Destruction by Anti-VCAM-1 Coaguligand

A. Methods

Male CB17 SCID mice were injected subcutaneously with 1×10^7 L540 cells as described above. When the tumors had reached a volume of 0.4-0.6 cm^3 , the mice were injected intravenously with either 20 μg of anti-VCAM-1-tTF, 16 μg anti-VCAM-1 antibody, 4 μg tTF, a mixture of 16 μg of anti-VCAM-1 antibody and 4 μg of tTF, 20 μg control IgG-tTF or saline. In some studies, the treatment was given 3 times, on days 0, 4 and 8. A minimum of 8 animals were treated in each group.

Animals were monitored daily for tumor measurements and body weight. Mice were sacrificed when tumors had reached a diameter of 2 cm^3 , or earlier if tumors showed signs of necrosis or ulceration. Tumor volume was calculated according to the formula: $\pi/6 \times D \times d^2$,

where D is the larger tumor diameter and d is the smaller diameter. Differences in tumor growth rates were tested for statistical significance using a non-parametric test (Mann-Whitney rank sum test) that makes no assumptions about tumor size being normally distributed (Gibbons, 1976).

5 B. Results

The anti-tumor activity of anti-VCAM-1•tTF coaguligand was determined in SCID mice bearing 0.3-0.4 cm³ L540 tumors. The drug was administered i.v. 3 times at intervals of 4 days. The pooled results from 3 separate studies are presented in FIG. 2 and Table 4. Mean tumor volume of anti-VCAM-1•tTF treated mice was significantly reduced at 21 days of treatment (P < 0.001) in comparison to all other groups. Nine of a total of 15 mice treated with the specific coaguligand showed more than 50% reduction in tumor volume. This effect was specific since unconjugated tTF, control IgG coaguligand and mixture of free anti-VCAM-1 antibody and tTF did not affect tumor growth.

00151362-071299
662120-2981560

TABLE 4

Inhibition of Tumor Growth by Anti-VCAM-1•tTF Coaguligand

| Treatment ^a | N | Mean tumor volume (mm ³) ^b | | Tumor Growth Index ^c | P versus saline ^d |
|------------------------|----|--|------------|------------------------------------|---------------------------------|
| | | Day 0 | Day 21 | | |
| Saline | 14 | 331 ± 61 | 2190 ± 210 | 6.91 | - |
| TTF | 13 | 341 ± 22 | 2015 ± 205 | 5.90 | NS |
| Anti-VCAM-1 | 16 | 363 ± 24 | 1920 ± 272 | 5.28 | NS |
| Anti-VCAM-1+tTF | 13 | 349 ± 42 | 2069 ± 362 | 5.92 | NS |
| Control IgG•tTF | 8 | 324 ± 30 | 2324 ± 304 | 7.17 | NS |
| Anti-VCAM-1•tTF | 15 | 365 ± 28 | 1280 ± 130 | 3.50 | < 0.001 |

^a L540 tumor bearing mice were injected i.v. with one of the following reagents: saline; 8 µg of unconjugated tTF; 32 µg of unconjugated anti-VCAM-1 antibody; mixture of 8 µg of tTF and 32 µg of anti-VCAM-1 antibody; 40 µg of control IgG•tTF (R187) coaguligand; or 40 µg of anti-VCAM-1•tTF coaguligand. The treatment was repeated on day 4 and 7 after first injection.

^b Mean tumor volume ± SD.

^c The tumor growth index is the ratio of mean tumor volume on day 21 to mean tumor volume on day 0.

^d Two tailed P values are for differences in tumor volume (day 21) for the treated groups versus the saline group as determined by the Mann-Whitney rank sum test.

EXAMPLE VIII

Phosphatidylserine Expression on Tumor Blood Vessels

A. Methods

1. Antibodies

Anti-phosphatidylserine (anti-PS) and anti-cardiolipin antibodies, both mouse monoclonal IgM antibodies, were produced as described by Rote (Rote *et al.*, 1993). Details of the characterization of the anti-PS and anti-cardiolipin antibodies were also reported by Rote *et al.* (1993, incorporated herein by reference).

2. Detection of PS Expression on Vascular Endothelium

L540 tumor-bearing mice were injected i.v. with 20 µg of either anti-PS or anti-cardiolipin mouse IgM antibodies. After 10 min., mice were anesthetized and their blood circulations were perfused with heparinized saline. Tumors and normal tissues were removed and snap-frozen. Serial sections of organs and tumors were stained with either HRP-labeled anti-mouse IgM for detection of anti-PS antibody or with anti-VCAM-1 antibody followed by HRP-labeled anti-rat Ig.

To preserve membrane phospholipids on frozen sections, the following protocol was developed. Animals were perfused with DPBS containing 2.5 mM Ca²⁺. Tissues were mounted on 3-aminopropyltriethoxysilane-coated slides and were stained within 24 h. No organic solvents, formaldehyde or detergents were used for fixation or washing of the slides. Slides were re-hydrated by DPBS containing 2.5 mM Ca²⁺ and 0.2% gelatin. The same solution was also used to wash sections to remove the excess of reagents. Sections were incubated with HRP-labeled anti-mouse IgM for 3.5 h at room temperature to detect anti-PS IgM.

B. Results

To explain the lack of thrombotic effect of anti-VCAM-1-tTF on VCAM-1 positive vasculature in heart and lungs, the inventors developed a concept of differential PS localization between normal and tumor blood vessels. Specifically, they hypothesized that endothelial cells in normal tissues segregate PS to the inner surface of the plasma membrane phospholipid bilayer, where it is unable to participate in thrombotic reactions; whereas endothelial cells in tumors translocate PS to the external surface of the plasma membrane, where it can support the coagulation action of the coaguligand. PS expression on the cell surface allows coagulation because it enables the attachment of coagulation factors to the membrane and coordinates the assembly of coagulation initiation complexes (Ortel *et al.*, 1992).

The inventors' model of PS translocation to the surface of tumor blood vessel endothelial cells, as developed herein, is surprising in that PS expression does not occur after, and does not

inevitably trigger, cell death. PS expression at the tumor endothelial cell surface is thus sufficiently stable to allow PS to serve as a targetable entity for therapeutic intervention.

To confirm the hypothesis that tumor blood vessel endothelium expresses PS on the luminal surface of the plasma membrane, the inventors used immunohistochemistry to determine the distribution of anti-PS antibody after intravenous injection into L540 tumor bearing mice. Anti-PS antibody localized within 10 min. to the majority of tumor blood vessels, including vessels in the central region of the tumor that can lack VCAM-1. Vessels that were positive for VCAM-1 were also positive for PS. Thus, there is coincident expression of PS on VCAM-1-expressing vessels in tumors.

In the *in vivo* localization studies, none of the vessels in normal organs, including VCAM-1-positive vasculature of heart and lung, were stained, indicating that PS is absent from the external surface of the endothelial cells. In contrast, when sections of normal tissues and tumors were directly stained with anti-PS antibody *in vitro*, no differences were visible between normal and tumor, endothelial or other cell types, showing that PS is present within these cells but only becomes expressed on the surface of endothelial cells in tumors.

The specificity of PS detection was confirmed by two independent studies. First, a mouse IgM monoclonal antibody directed against a different negatively charged lipid, cardiolipin, did not home to tumor or any organs *in vivo*. Second, pretreatment of frozen sections with acetone abolished staining with anti-PS antibody, presumably because it extracted the lipids together with the bound anti-PS antibody.

EXAMPLE IX

Annexin V Blocks Coaguligand Activation of Factor X *In Vitro*

A. Methods

The ability of Annexin V to affect Factor Xa formation induced by coaguligand was determined by a chromogenic assay described above in Example V. IL-1 α -stimulated bEnd.3 cells were incubated with anti-VCAM-tTF and permeabilized by saponin. Annexin V was

added at concentrations ranging from 0.1 to 10 µg/ml and cells were incubated for 30 min. before addition of diluted Proplex T. The amount of Factor Xa generated in the presence or absence of Annexin V was determined as described in Example V. Each treatment was performed in duplicate and repeated at least twice.

5

B. Results

The need for surface PS expression in coaguligand action is further indicated by the inventors' finding that annexin V, which binds to PS with high affinity, blocks the ability of anti-VCAM-1•tTF bound to bEnd.3 cells to generate factor Xa *in vitro*.

10

Annexin V added to permeabilized cells preincubated with anti-VCAM-1•tTF inhibited the formation of factor Xa in a dose-dependent manner (FIG. 3). In the absence of Annexin V, cell-bound coaguligand produced 95 ng of factor Xa per 10,000 cells per 60 min. The addition of increasing amounts of Annexin V (in the µg per ml range) inhibited factor Xa production. At 10 µg per ml, Annexin V inhibited factor Xa production by 58% (FIG. 3). No further inhibition was observed by increasing the concentration of Annexin V during the assay, indicating that annexin V saturated all available binding sites at 10 µg per ml.

09351862-074259

EXAMPLE X

Annexin V Blocks Coaguligand Activity *In Vivo*

A. Methods

The ability of Annexin V to inhibit coaguligand-induced thrombosis *in vivo* was examined in L540 Hodgkin-bearing SCID mice. Tumors were grown in mice as described above in Example II. Two mice per group (tumor size 0.5 cm in diameter) were injected intravenously via the tail vein with one of the following reagents: a) saline; b) 100 µg of Annexin V; c) 40 µg of anti-VCAM-1•tTF; d) 100 µg of Annexin V followed 2 hours later by 40 µg of anti-VCAM-1•tTF.

Four hours after the last injection mice were anesthetized and perfused with heparinized saline. Tumors were removed, fixed with 4% formalin, paraffin-embedded and stained with

hematoxyline-eosin. The number of thrombosed and non-thrombosed blood vessels were counted and the percentage of thrombosis was calculated.

B. Results

5 Annexin V also blocks the activity of the anti-VCAM-1•tTF coaguligand *in vivo*. Groups of tumor-bearing mice were treated with one of the control or test reagents, as described in the methods. The mice were given (a) saline; (b) 100 µg of Annexin V; (c) 40 µg of anti-VCAM-1•tTF coaguligand; or (d) 100 µg of Annexin V followed 2 hours later by 40 µg of anti-VCAM-1•tTF coaguligand. Identical results were obtained in both mice per group.

10

No spontaneous thrombosis, hemorrhages or necrosis were observed in tumors derived from saline-injected mice. Treatment with Annexin V alone did not alter tumor morphology.

10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500

5 In accordance with other data presented herein, 40 µg of anti-VCAM-1•tTF coaguligand caused thrombosis in 70% of total tumor blood vessels. The majority of blood vessels were occluded with packed erythrocytes and clots, and tumor cells were separated from one another. Both coaguligand-induced anti-tumor effects, *i.e.*, intravascular thrombosis and changes in tumor cell morphology, were completely abolished by pre-treating the mice with Annexin V.

20 These findings confirm that the anti-tumor effects of coaguligands are mediated through the blockage of tumor vasculature. These data also demonstrate that PS is essential for coaguligand-induced thrombosis *in vivo*.

EXAMPLE XI

25 Externalized Phosphatidylserine is a Global Marker of Tumor Blood Vessels

A. Methods

PS exposure on tumor and normal vascular endothelium was examined in three animal tumor models: L540 Hodgkin lymphoma, NCI-H358 non-small cell lung carcinoma, and HT 29 colon adenocarcinoma (ATCC). To grow the tumors *in vivo*, 2×10^6 cells were injected into the
30 right flank of SCID mice and allowed to reach 0.8-1.2 cm in diameter. Mice bearing large

tumors (volume above 800 mm³) were injected intravenously *via* the tail vein with 20 µg of either anti-PS or anti-cardiolipin antibodies. The anti-cardiolipin antibody served as a control for all studies since both antibodies are directed against negatively charged lipids and belong to the same class of immunoglobulins (mouse IgM).

5

One hour after injection, mice were anesthetized and their blood circulation was perfused with heparinized saline. Tumors and normal organs were removed and snap-frozen. Frozen sections were stained with anti-mouse IgM-peroxidase conjugate (Jackson Immunoresearch Labs) followed by development with carbazole.

10

B. Results

The anti-PS antibodies specifically homed to the vasculature of all three tumors (HT 29, L540 and NCI-H358) *in vivo*, as indicated by detection of the mouse IgM. The average percentages of vessels stained in the tumors were 80% for HT 29, 30% for L540 and 50% for NCI-H358. Vessels in all regions of the tumors were stained and there was staining both of small capillaries and larger vessels.

No vessel staining was observed with anti-PS antibodies in any normal tissues. In the kidney, tubules were stained both with anti-PS and anti-CL, and this likely relates to the secretion of IgMs by this organ (Table 5). Anti-cardiolipin antibodies were not detected in any tumors or normal tissues, except kidney.

These findings indicate that only tumor endothelium exposes PS to the outer site of the plasma membrane.

25

TABLE 5

Vessel L calization of Anti-PS and Anti-Cardiolipin Abs in Tumor-Bearing Mice*

| Tissue | Anti-PS† | Anti-Cardiolipin‡ |
|------------------|----------|-------------------|
| L540 Cy tumor | ++ | - |
| H358 tumor | ++ | - |
| HT29 tumor | +++ | - |
| Adrenal | - | - |
| Brain Cerebellum | - | - |
| Brain Cortex | - | - |
| Heart | - | - |
| Kidney | -‡ | -‡ |
| Large Intestine | - | - |
| Liver | - | - |
| Lung | - | - |
| Pancreas | - | - |
| Small Intestine | - | - |
| Spleen | - | - |
| Testes | - | - |

*Biodistribution in normal organs of both anti-PS and anti-cardiolipin Abs was identical in all three tumor animal models.

5 †Anti-PS and anti-cardiolipin antibodies were detected on frozen sections using anti-mouse IgM-peroxidase conjugate. - no staining; + weak; ++ moderate; +++ strong, equivalent to pan endothelial marker Meca 32.

‡Tubular staining was observed in the kidneys of both anti-PS and anti-CL recipients.

10 To estimate the time at which tumor vasculature loses the ability to segregate PS to the inner side of the membrane, the inventors examined anti-PS localization in L540 tumors ranging in volume from 140 to 1,600 mm³. Mice were divided into 3 groups according to their tumor size: 140-300, 350-800 and 800-1,600 mm³. Anti-PS Ab was not detected in three mice bearing small L540 tumors (up to 300 mm³). Anti-PS Ab localized in 3 animals of 5 in the group of

intermediate size L540 tumors and in all mice (4 out of 4) bearing large L540 tumors (Table 6). Percent of PS-positive blood vessels from total (identified by pan endothelial marker Meca 32) was 10-20% in the L540 intermediate group and 20-40% in the group of large L540 tumors (Table 6).

5

TABLE 6

PS Externalization Detected in Mid and Large Sized Tumors

| Tumor Size (mm³) | No. Positive Tumors/Total* | %PS-Positive Vessels/Total† |
|------------------------------------|-----------------------------------|------------------------------------|
| 350-800 | 3/5 | 10-20 |
| 850-1,600 | 4/4 | 20-40 |

*Mice bearing L540 Cy tumors were divided into three groups according to tumor size. 20 µg of anti-PS antibodies were injected i.v. and allowed to circulate for 1 hour. Mouse antibodies were detected on frozen sections using anti-mouse IgM-peroxidase conjugate.

†Total number of blood vessels was determined using pan endothelial Ab Meca 32. PS positive and Meca-positive vessels were counted in 4 fields per cross section of tumor. Range of % PS-positive vessels within the same group is shown.

10
15
20
25

EXAMPLE XII

Anti-Tumor Effects of Unconjugated Anti-Phosphatidylserine Antibodies

A. Methods

The effects of anti-PS antibodies were examined in syngeneic and xenogeneic tumor models. For the syngeneic model, 1×10^7 cells of murine colorectal carcinoma Colo 26 (obtained from Dr. Ian Hart, ICRF, London) were injected subcutaneously into the right flank of Balb/c mice. In the xenogeneic model, a human Hodgkin's lymphoma L540 xenograft was established by injecting 1×10^7 cells subcutaneously into the right flank of male CB17 SCID mice. Tumors were allowed to grow to a size of about 0.6-0.9 cm³ before treatment.

Tumor-bearing mice (4 animals per group) were injected i.p. with 20 µg of naked anti-PS antibody (IgM), control mouse IgM or saline. Treatment was repeated 3 times with a 48 hour interval. Animals were monitored daily for tumor measurements and body weight. Tumor

volume was calculated as described in Example VII. Mice were sacrificed when tumors had reached 2 cm³, or earlier if tumors showed signs of necrosis or ulceration.

B. Results

5 The growth of both syngeneic and xenogeneic tumors was effectively inhibited by treatment with naked anti-PS antibodies (FIG. 4A and FIG. 4B). Anti-PS antibodies caused tumor vascular injury, accompanied by thrombosis, and tumor necrosis. The presence of clots and disintegration of tumor mass surrounding blocked blood vessels was evident.

10 Quantitatively, the naked anti-PS antibody treatment inhibited tumor growth by up to 60% of control tumor volume in mice bearing large Colo 26 (FIG. 4A) and L540 (FIG. 4B) tumors. No retardation of tumor growth was found in mice treated with saline or control IgM. No toxicity was observed in mice treated with anti-PS antibodies, with normal organs preserving unaltered morphology, indistinguishable from untreated or saline-treated mice.

15 Tumor regression started 24 hours after the first treatment and tumors continue to decline in size for the next 6 days. This was observed in both syngeneic and immunocompromised tumor models, indicating that the effect was mediated by immune status-independent mechanism(s). Moreover, the decline in tumor burden was associated with the increase of alertness and generally healthy appearance of the animals, compared to control mice bearing tumors larger than 1500 mm³. Tumor re-growth occurred 7-8 days after the first treatment.

20 The results obtained with anti-PS treatment of L540 tumors are further compelling for the following reasons. Notably, the tumor necrosis observed in L540 tumor treatment occurred despite the fact that the percentage of vessels that stained positive for PS in L540 tumors was less than in HT 29 and NCI-H358 tumors. This implies that even more rapid necrosis would likely result when treating other tumor types. Furthermore, L540 tumors are generally chosen as an experimental model because they provide clean histological sections and they are, in fact, known to be resistant to necrosis.

30

EXAMPLE XIII

Anti-Tumor Effects of Annexin Conjugates

The surprising finding that aminophospholipids are stable markers of tumor vasculature also means that antibody-therapeutic agent constructs can be used in cancer treatment. In addition to using antibodies as targeting agents, the inventors reasoned that annexins, and other aminophospholipid-binding proteins, could also be used to specifically deliver therapeutic agents to tumor vasculature. The following data shows the anti-tumor effects that result from the *in vivo* administration of annexin-TF constructs.

10 A. Methods

An annexin V-tTF conjugate was prepared and administered to nu/nu mice with solid tumors. The tumors were formed from human HT29 colorectal carcinoma cells that formed tumors of at least about 1.2 cm³. The annexin V-tTF coaguligand (10 µg) was administered intravenously and allowed to circulate for 24 hours. Saline-treated mice were separately maintained as control animals. After the one day treatment period, the mice were sacrificed and exsanguinated and the tumors and major organs were harvested for analysis.

15 B. Results

The annexin V-tTF conjugate was found to induce specific tumor blood vessel coagulation in HT29 tumor bearing mice. Approximately 55% of the tumor blood vessels in the annexin V-tTF conjugate treated animals were thrombosed following a single injection. In contrast, there was minimal evidence of thrombosis in the tumor vasculature of the control animals.

25 EXAMPLE XIV

Phosphatidylserine Translocation in the Tumor Environment

The discovery of PS as an *in vivo* surface marker unique to tumor vascular endothelial cells prompted the inventors to further investigate the effect of a tumor environment on PS translocation and outer membrane expression. The present example shows that exposing

endothelial cells *in vitro* to certain conditions that mimic those in a tumor duplicates the observed PS surface expression in intact, viable cells.

A. Methods

5 Mouse bEnd.3 endothelial cells were seeded at an initial density of 50,000 cells/well. Twenty-four later cells were incubated with increasing concentrations of H_2O_2 (from 10 μM to 500 μM) for 1 hour at 37°C or left untreated. At the end of the incubation, cells were washed 3 times with PBS containing 0.2% gelatin and fixed with 0.25% glutaraldehyde. Identical wells were either stained with anti-PS IgM or trypsinized and evaluated for viability by the Trypan
10 Blue exclusion test. For the anti-PS staining, after blocking with 2% gelatin for 10 min., cells were incubated with 2 $\mu g/ml$ of anti-PS antibody, followed by detection with anti-mouse IgM-HRP conjugate.

15 Wells seeded with mouse bEnd.3 endothelial cells were also incubated with different effectors and compared to control, untreated wells after the same period of incubation at 37°C. The panel of effectors tested included TNF, LPS, bFGF, IL-1 α , IL-1 β and thrombin. After incubation, cells were washed and fixed and were again either stained with anti-PS IgM or evaluated for viability using the Trypan Blue exclusion test, as described above.

B. Results

1. PS Induction by H_2O_2

25 Exposing endothelial cells to H_2O_2 at concentrations higher than 100 μM caused PS translocation in ~90% cells. However, this was accompanied by detachment of the cells from the substrate and cell viability decreasing to about 50-60%. The association of surface PS expression with decreasing cell viability is understandable, although it is still interesting to note that ~90% PS translocation is observed with only a 50-60% decrease in cell viability.

30 Using concentrations of H_2O_2 lower than 100 μM resulted in significant PS expression without any appreciable reduction in cell viability. For example, PS was detected at the cell surface of about 50% of cells in all H_2O_2 treated wells using H_2O_2 at concentrations as low as

translocation to the endothelial cell surface without disrupting cell integrity has important implications in addition to validating the original *in vivo* data and the inventors' therapeutic approaches.

5 Human, bovine and murine endothelial cells are all known to be PS-negative under normal conditions. Any previously documented PS expression has always been associated with cell damage and/or cell death. This is simply not the case in the present studies, where normal viability is maintained. This shows that PS translocation in tumor vascular endothelium is mediated by biochemical mechanisms unconnected to cell damage. This is believed to be the
10 first demonstration of PS surface expression in morphologically intact endothelial cells and the first indication that PS expression can be disconnected from the apoptosis pathway(s). Returning to the operability of the present invention, these observations again confirm that PS is a sustainable, rather than transient, marker of tumor blood vessels and a suitable candidate for therapeutic intervention.

15 2. PS Expression Does Not Correlate with Cell Activation

The relevance of this *in vitro* data to the tumor environment is also strengthened by the fact that other, general cell activators are without effect on PS translocation in endothelial cells. For example, the inventors tested TNF in similarly controlled studies and found it unable to induce PS surface expression, despite the expected increases in E-selectin and VCAM expression. Likewise, LPS, bFGF, IL-1 α and IL-1 β were all without effect on PS expression in
20 appropriately controlled studies.

3. PS Induction by Thrombin

25 In contrast to the lack of effects of other cell activators, thrombin was observed to increase PS expression, although not to the same extent as H₂O₂. This data is also an integral part of the tumor-induction model of PS expression developed by the present inventors (thrombin-induced PS surface expression in normal tissues would also further coagulation as PS expression coordinates the assembly of coagulation initiation complexes (Ortel *et al.*, 1992)).

The tumor environment is known to be prothrombotic, such that tumor vasculature is predisposed to coagulation (U.S. Patent No. 5,877,289). As thrombin is a product of the coagulation cascade, it is present in tumor vasculature. In fact, the presence of thrombin induces VCAM expression, contributing to the inventors' ability to exploit VCAM as a targetable marker of tumor vasculature (U.S. Patent Nos. 5,855,866; 5,877,289). The present data showing that thrombin also induces PS expression is thus both relevant to targeting aminophospholipids with naked antibodies and therapeutic conjugates, and further explains the beneficial effects of the anti-VCAM coaguligand containing Tissue Factor (Example VII).

10

* * *

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202

REFERENCES

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

- 5 Abrams and Oldham, In: *Monoclonal Antibody Therapy of Human Cancer*, Foon and Morgan (Eds.), Martinus Nijhoff Publishing, Boston, pp. 103-120, 1985.
- Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, 1988.
- 10 Asahara, Chen, Takahashi, Fujikawa, Kearney, Magner, Yancopoulos, Isner, "Tie2 receptor ligands, angiopoietin-1 and angiopoietin-2, modulate VEGF-induced postnatal neovascularization" *Circ. Res.*, 83(3):233-40, 1998.
- 15 Barbas, Kang, Lerner and Benkovic, "Assembly of combinatorial antibody libraries on phage surfaces: the gene III site," *Proc. Natl. Acad. Sci., USA*, 88(18):7978-7982, 1991.
- 20 Berard, Boffa, Karmochkine, Aillaud, Juhan-Vague, Frances, Cacoub, Piette and Harle, "Plasma reactivity to hexagonal II phase phosphatidylethanolamine is more frequently associated with lupus anticoagulant than with antiphosphatidylethanolamine antibodies," *J. Lab. Clin. Med.*, 122(5):601-605, 1993.
- 25 Berman, Mellis, Pollock, Smith, Suh, Heinke, Kowal, Surti, Chess, Cantor, *et al.*, "Content and organization of the human Ig VH locus: definition of three new VH families and linkage to the Ig CH locus," *EMBO J.*, 7(3):727-738, 1988.
- 30 Bervers, Comburius and Zwaal, "The nature of the binding site for prothrombinase at the platelet surface as revealed by lipolytic enzymes," *Eur. J. Biochem.*, 122:81-85, 1982.
- 35 Bervers, Comfurius and Zwaal, "Changes in membrane phospholipid distribution during platelet activation," *Biochim. Biophys. Acta*, 736:57-66, 1983.
- 40 Bervers, Rosing and Zwaal, "Development of procoagulant binding sites on the platelet surface," *Adv. Exp. Med. Biol.*, 192:359-371, 1985.
- 45 Bervers, Galli, Barbui, Comfurius and Zwaal, "Lupus anticoagulant IgG's (LA) are not directed to phospholipids only, but to a complex of lipid-bound human prothrombin," *Thromb. Haemost.*, 66(6):629-32, 1991.
- 50 Bevilacqua, "Endothelial-leukocyte adhesion molecules," *Ann. Rev. Immunol.*, 11:767-804, 1993.
- 55 Blankenberg, Katsikis, Tait, Davis, Naumovski, Ohtsuki, Kopiwoda, Abrams, Darkes, Robbins, Maecker and Strauss, "In vivo detection and imaging of phosphatidylserine expression during programmed cell death," *Proc. Natl. Acad. Sci., USA*, 95:6349-6354, 1998.

Campbell, In: *Monoclonal Antibody Technology, Laboratory Techniques in Biochemistry and Molecular Biology*, Vol. 13, Burden and Von Knippenberg (Eds.), Elsevier, Amsterdam, pp. 75-83, 1984.

- 5 Carnemolla *et al.*, "A tumor-associated fibronectin isoform generated by alternative splicing of messenger RNA precursors," *J. Cell Biol.*, 108:1139-1148, 1989.

Chamley, McKay and Pattison, "Cofactor dependent and cofactor independent anticardiolipin antibodies," *Thromb. Res.*, 61(3):291-9, 1991.

10

Clapp *et al.*, "The 16-kilodalton N-terminal fragment of human prolactin is a potent inhibitor of angiogenesis," *Endocrinology*, 133(3):1292-1299, 1993.

15

Connor, Bucana, Fidler and Schroit, "Differentiation-dependent expression of phosphatidylserine in mammalian plasma membranes: quantitative assessment of outer-leaflet lipid by prothrombinase complex formation," *Proc. Natl. Acad. Sci. USA*, 86(9):3184-3188, 1989.

20

Coughlin *et al.*, "Interleukin-12 and interleukin-18 synergistically induce murine tumor regression which involves inhibition of angiogenesis," *J. Clin. Invest.*, 101(6):1441-1452, 1998.

D'Amato *et al.*, "Thalidomide is an inhibitor of angiogenesis," *Proc. Natl. Acad. Sci. USA*, 91(9):4082-4085, 1994.

25

D'Angelo *et al.*, "Activation of mitogen-activated protein kinases by vascular endothelial growth factor and basic fibroblast growth factor in capillary endothelial cells is inhibited by the antiangiogenic factor 16-kDa N- terminal fragment of prolactin," *Proc. Natl. Acad. Sci. USA*, 92(14):6374-6378, 1995.

30

Dachary-Prigent, Toti, Satta, Pasquet, Uzan and Freyssinet, "Physiopathological significance of catalytic phospholipids in the generation of thrombin," *Seminars In Thrombosis and Hemostasis*, 22:157-164, 1996.

35

Davis and Yancopoulos, "The angiopoietins: Yin and Yang in angiogenesis", *Curr. Top. Microbiol. Immunol.*, 237:173-85, 1999.

de Jong, Geldwerth and Kuypers, "Oxidative damage does not alter membrane phospholipid asymmetry in human erythrocytes," *Am. Chem. Soc.*, 1997.

40

Denekamp, "Vascular attack as a therapeutic strategy for cancer," *Cancer Metastasis Rev.*, 9:267-282, 1990.

DeVore *et al.*, "Phase I Study of the Antineovascularization Drug CM101," *Clin. Cancer Res.*, 3(3):365-372, 1997.

45

- Diehl, Pfreundschuh, Fonatsch, Stein, Falk, Burrichter and Schaadt, "Phenotypic genotypic analysis of Hodgkin's disease derived cell lines: histopathological and clinical implications," *Cancer Surveys*, 4:399-416, 1985.
- 5 Donati, "Cancer and thrombosis: from Phlegmasia *alba dolens* to transgenic mice," *Thromb. Haemost.*, 74:278-281, 1995.
- Drouvalakis and Buchanan, "Phospholipid specificity of autoimmune and drug induced lupus anticoagulants; association of phosphatidylethanolamine reactivity with thrombosis in
10 autoimmune disease," *J. Rheumatol.*, 25(2):290-295, 1998.
- Droz, Patey, Paraf, Chretien and Gogusev, "Composition of extracellular matrix and distribution of cell adhesion molecules in renal cell tumors," *Lab. Invest.*, 71:710-718, 1994.
- 15 Dvorak, Nagy and Dvorak, "Structure of Solid Tumors and Their Vasculature: Implications for Therapy with Monoclonal Antibodies," *Cancer Cells*, 3(3):77-85, 1991.
- Edgington, Mackman, Brand and Ruf, "The Structural Biology of Expression and Function of Tissue Factor," *Thromb. Haemost.*, 66(1):67-79, 1991.
20
- Ferrara, Clapp, Weiner, "The 16K fragment of prolactin specifically inhibits basal or fibroblast growth factor stimulated growth of capillary endothelial cells," *Endocrinology*, 129(2):896-900, 1991.
- 25 Folkman *et al.*, "Angiogenesis inhibition and tumor regression caused by heparin or a heparin fragment in the presence of cortisone," *Science*, 221:719-725, 1983.
- Fotsis *et al.*, "The endogenous oestrogen metabolite 2-methoxyoestradiol inhibits angiogenesis and suppresses tumour growth," *Nature*, 368(6468):237-239, 1994.
30
- Flynn, Byrne, Baglin, Weissberg and Bennett, "Thrombin generation by apoptotic vascular smooth muscle cells," *Blood*, 89(12):4378-4384, 1997.
- Frater-Schroder *et al.*, "Tumor necrosis factor type alpha, a potent inhibitor of endothelial cell growth *in vitro*, is angiogenic *in vivo*," *Proc. Natl. Acad. Sci. USA*, 84(15):5277-5281,
35 1987.
- Frazier, "Thrombospondins," *Curr. Opin. Cell Biol.*, 3(5):792-799, 1991.
- 40 Fries, Williams, Atkins, Newman, Lipscomb and Collins, "Expression of VCAM-1 and E-selectin in an *in vivo* model of endothelial activation," *Am. J. Pathol.*, 143:725-737, 1993.

- Gaffet, Bettache and Bienvenüe, "Transverse redistribution of phospholipids during human platelet activation: evidence for a vectorial outflux specific to aminophospholipids," *Biochem.*, 34:6762-6769, 1995.
- 5 Gagliardi and Collins, "Inhibition of angiogenesis by antiestrogens," *Cancer Res.*, 53(3):533-535, 1993.
- Gagliardi, Hadd, Collins, "Inhibition of angiogenesis by suramin," *Cancer Res.*, 52(18):5073-5075, 1992.
- 10 Gagliardi *et al.*, "Antiangiogenic and antiproliferative activity of suramin analogues," *Cancer Chemother. Pharmacol.*, 41(2):117-124, 1998.
- 15 Galli, Comfurius, Maassen Hemker, de Baets, van Breda-Vriesman, Barbui, Zwaal and Bevers, "Anticardiolipin antibodies (ACA) directed not to cardiolipin but to a plasma protein cofactor," *Lancet*, 335(8705):1544-7, 1990.
- Galli, Barbui, Zwaal, Comfurius and Bevers, "Antiphospholipid antibodies: involvement of protein cofactors," *Haematologica*, 78(1):1-4, 1993.
- 20 Ge and Butcher, "Cloning and expression of a cDNA encoding mouse endoglin, an endothelial cell TGF-beta ligand," *Gene*, 138:201-206, 1994.
- Gefter *et al.*, "A simple method for polyethylene glycol-promoted hybridization of mouse myeloma cells," *Somatic Cell Genet.*, 3:231-236, 1977.
- 25 Gibbons, "Mann-Whitney-Wilcoxon test for two independent samples," In: *Nonparametric methods for quantitative analysis*, J.D.Gibbons (ed.), Holt, Rinehart and Winston, New York, pp. 160, 1976.
- 30 Glennie, *et al.*, "Preparation and performance of bispecific F(ab' gamma)2 antibody containing thioether-linked Fab' gamma fragments," *J. Immunol.*, 139:2367-2375, 1987.
- Goding, In: *Monoclonal Antibodies: Principles and Practice*, 2nd Edition, Academic Press, Orlando, Fl., pp. 60-61, 65-66, 71-74, 1986.
- 35 Good *et al.*, "A tumor suppressor-dependent inhibitor of angiogenesis is immunologically and functionally indistinguishable from a fragment of thrombospondin," *Proc. Natl. Acad. Sci. USA*, 87(17):6624-6628, 1990.
- 40 Grant *et al.*, "Fibronectin fragments modulate human retinal capillary cell proliferation and migration," *Diabetes*, 47(8):1335-1340, 1998.

- Hagemeier *et al.*, "A Monoclonal Antibody Reacting with Endothelial Cells of Budding Vessels in Tumors and Inflammatory Tissues, and Non-Reactive with Normal Adult Tissues," *Int. J. Cancer*, 38:481-488, 1986.
- 5 Hahne, Jager, Isenmann, Hallmann and Vestweber, "Five tumor necrosis factor-inducible cell adhesion mechanisms on the surface of mouse endothelioma cells mediate the binding of leukocytes," *J. Cell Biol.*, 121:655-664, 1993.
- 10 Hampton, Vanags, Porn-Ares and Orrenius, "Involvement of extracellular calcium in phosphatidylserine exposure during apoptosis," *FEBS Lett.*, 399(3):277-282, 1996.
- Haran *et al.*, "Tamoxifen enhances cell death in implanted MCF7 breast cancer by inhibiting endothelium growth," *Cancer Res.*, 54(21):5511-5514, 1994.
- 15 Hasselaar and Sage, "SPARC antagonizes the effect of basic fibroblast growth factor on the migration of bovine aortic endothelial cells," *J. Cell Biochem.*, 49(3):272-283, 1992.
- Hellerqvist *et al.*, "Antitumor effects of GBS toxin: a polysaccharide exotoxin from group B beta-hemolytic streptococcus," *J. Cancer Res. Clin. Oncol.*, 120(1-2):63-70, 1993.
- 20 Hiscox and Jiang, "Interleukin-12, an emerging anti-tumour cytokine," *In Vivo*, 11(2):125-132, 1997.
- 25 Holash *et al.*, "Vessel Cooption, Regression, and Growth in Tumors Mediated by Angiopoietins and VEGF", *Science*, 284:1994-1998, 1999.
- Hori *et al.*, "Differential effects of angiostatic steroids and dexamethasone on angiogenesis and cytokine levels in rat sponge implants," *Br. J. Pharmacol.*, 118(7):1584-1591, 1996.
- 30 Huang, Molema, King, Watkins, Edgington and Thorpe, "Tumor infarction in mice by antibody-directed targeting of tissue factor to tumor vasculature," *Science*, 275:547-550, 1997.
- 35 Huse, Sastry, Iverson, Kang, Alting-Mees, Burton, Benkovic and Lerner, *Science*, "Generation of a large combinatorial library of the immunoglobulin repertoire in phage lambda," 246(4935):1275-1281, 1989.
- Igarashi, Umeda, Tokita, Soo Nam and Inoue, "Effective induction of anti-phospholipid and anticoagulant antibodies in normal mouse," *Thrombosis Res.*, 61:135-148, 1991.
- 40 Ingber *et al.*, "Angioinhibins: Synthetic analogues of fumagillin which inhibit angiogenesis and suppress tumor growth," *Nature*, 48:555-557, 1990.

- Iwamoto *et al.*, "Inhibition of angiogenesis, tumour growth and experimental metastasis of human fibrosarcoma cells HT1080 by a multimeric form of the laminin sequence Tyr-Ile-Gly-Ser-Arg (YIGSR)," *Br. J. Cancer*, 73(5):589-595, 1996.
- 5 Jackson *et al.*, "Stimulation and inhibition of angiogenesis by placental proliferin and proliferin-related protein," *Science*, 266(5190):1581-1584, 1994.
- Jamasbi, Wan and Stoner, "Epitope masking of rat esophageal carcinoma tumor-associated antigen by certain coexisting glycolipid and phospholipid molecules: a potential
10 mechanism for tumor cell escape from the host immune responses," *Cancer Immunol. Immunother.*, 38(2):99-106, 1994.
- Jendraschak and Sage, "Regulation of angiogenesis by SPARC and angiostatin: implications for tumor cell biology," *Semin. Cancer Biol.*, 7(3):139-146, 1996.
- 15 Jones, Dear, Foote, Neuberger and Winter, "Replacing the complementarity-determining regions in a human antibody with those from a mouse," *Nature*, 321(6069):522-525, 1986.
- Julien, Tournier and Tocanne, "Differences in the transbilayer and lateral motions of fluorescent analogs of phosphatidylcholine and phosphatidylethanolamine in the apical plasma
20 membrane of bovine aortic endothelial cells," *Exp. Cell. Res.*, 208(2):387-9, 1993.
- Julien, Tournier and Tocanne, "Basic fibroblast growth factor modulates the aminophospholipid translocase activity present in the plasma membrane of bovine aortic endothelial cells,"
25 *Eur. J. Biochem.*, 230:287-297, 1995.
- Julien, Millot, Tocanne and Tournier, "12-O-Tetradecanoylphorbol-13-Acetate inhibits aminophospholipid translocase activity and modifies the lateral motions of fluorescent phospholipid analogs in the plasma membrane of bovine aortic endothelial cells,"
30 *Experimental Cell Res.*, 234:125-131, 1997.
- Kang, Barbas, Janda, Benkovic and Lerner, "Linkage of recognition and replication functions by assembling combinatorial antibody Fab libraries along phage surfaces," *Proc. Natl. Acad. Sci., U.S.A.*, 88(10):4363-4366, 1991.
- 35 Katsuragawa, Kanzaki, Inoue, Hirano, Mori and Rote, "Monoclonal antibody against phosphatidylserine inhibits *in vitro* human trophoblastic hormone production and invasion," *Biology of Reproduction*, 56:50-58, 1997.
- 40 Kenyon, Browne, D'Amato, "Effects of thalidomide and related metabolites in a mouse corneal model of neovascularization," *Exp. Eye Res.*, 64(6):971-978, 1997.
- Kim, Kwak, Ahn, So, Liu, Koh, Koh, "Molecular cloning and characterization of a novel angiopoietin family protein, angiopoietin-3", *FEBS Lett.*, 443(3):353-6, 1999.
- 45

- 5
10
15
20
25
30
35
40
45
- Kleinman *et al.*, "The laminins: a family of basement membrane glycoproteins important in cell differentiation and tumor metastases," *Vitam. Horm.*, 47:161-186, 1993.
- Kohler and Milstein, "Continuous cultures of fused cells secreting antibody of predefined specificity," *Nature*, 256:495-497, 1975.
- Kohler and Milstein, "Derivation of specific antibody-producing tissue culture and tumor lines by cell fusion," *Eur. J. Immunol.*, 6:511-519, 1976.
- Konieczny, Bobrzecka, Laidler and Rybarska, "The combination of IgM subunits and proteolytic IgG fragment by controlled formation of interchain disulphides," *Haematologia*, 14(1):95-99, 1981.
- Kuzu, Bicknell, Fletcher and Gatter, "Expression of adhesion molecules on the endothelium of normal tissue vessels and vascular tumors," *Lab. Invest.*, 69(3):322-328, 1993.
- Lane, Iruela-Arispe, Sage, "Regulation of gene expression by SPARC during angiogenesis *in vitro*. Changes in fibronectin, thrombospondin-1, and plasminogen activator inhibitor-1," *J. Biol. Chem.*, 267(23):16736-16745, 1992.
- Lee *et al.*, "Inhibition of urokinase activity by the antiangiogenic factor 16K prolactin: activation of plasminogen activator inhibitor 1 expression," *Endocrinology*, 139(9):3696-3703, 1998.
- Leppink, Bishop, Sedmak, Henry, Ferguson, Streeter, Butcher and Orosz, "Inducible expression of an endothelial cell antigen on murine myocardial vasculature in association with interstitial cellular infiltration," *Transplantation*, 48(5):874-877, 1989.
- Levy, Gharavi, Sammaritano, Habina and Lockshin, "Fatty acid chain is a critical epitope for antiphospholipid antibody," *J. Clin. Immunol.*, 10(3):141-145, 1990.
- Lin, Shroyer, Walter, Lyden, Ng and Rote, "Monoclonal IgM antiphosphatidylserine antibody reacts against cytoskeleton-like structures in cultured human umbilical cord endothelial cells," *Am. J. Reprod. Immun.*, 33:97-107, 1995.
- Lin, Buxton, Acheson, Radziejewski, Maisonpierre, Yancopoulos, Channon, Hale, Dewhirst, George, Peters, "Anti-angiogenic gene therapy targeting the endothelium-specific receptor tyrosine kinase Tie2", *Proc. Natl. Acad. Sci., USA*, 95(15):8829-34, 1998.
- Lindner and Borden, "Effects of tamoxifen and interferon-beta or the combination on tumor-induced angiogenesis," *Int. J. Cancer*, 71(3):456-461, 1997.
- Lingen, Polverini, Bouck, "Inhibition of squamous cell carcinoma angiogenesis by direct interaction of retinoic acid with endothelial cells," *Lab. Invest.*, 74(2):476-483, 1996.

- Lingen, Polverini, Bouck, "Retinoic acid and interferon alpha act synergistically as antiangiogenic and antitumor agents against human head and neck squamous cell carcinoma," *Cancer Res.*, 58(23):5551-5558, 1998.
- 5 Liu, Moy, Kim, Xia, Rajasekaran, Navarro, Knudsen and Bander, "Monoclonal antibodies to the extracellular domain of prostate-specific membrane antigen also react with tumor vascular endothelium", *Cancer Res.*, 57:3629-3634, 1997.
- 10 Majewski *et al.*, "Vitamin D3 is a potent inhibitor of tumor cell-induced angiogenesis," *J. Invest. Dermatol. Symp. Proc.*, 1(1):97-101, 1996.
- Mandriota and Pepper, "Regulation of angiopoietin -2 mRNA levels in bovine microvascular endothelial cells by cytokines and hypoxia", *Circ. Res.*, 83(8):852-9, 1998.
- 15 Maneta-Peyret, Bessoule, Geffard and Cassagne, "Demonstration of high specificity antibodies against phosphatidylserine," *J. Immun. Meth.*, 108:123-127, 1988.
- 20 Maneta-Peyret, Freyburger, Bessoule and Cassagne, "Specific immunocytochemical visualization of phosphatidylserine," *J. Immun. Methods*, 122:155-159, 1989.
- 25 Manetti *et al.*, "Synthesis and binding mode of heterocyclic analogues of suramin inhibiting the human basic fibroblast growth factor," *Bioorg. Med. Chem.*, 6(7):947-958, 1998.
- Martin, Reutelingsperger, McGahon, Rader, van Schie, LaFace and Green, "Early redistribution of plasma membrane phosphatidylserine is a general feature of apoptosis regardless of the initiating stimulus: inhibition by overexpression of Bcl-2 and Abl," *J. Exp. Med.*, 182(5):1545-1556, 1995.
- 30 Matsuura, Igarashi, Yasuda, Triplett and Koike, "Anticardiolipin antibodies recognize beta 2-glycoprotein I structure altered by interacting with an oxygen modified solid phase surface," *J. Exp. Med.*, 179(2):457-62, 1994.
- 35 McNeil, Simpson, Chesterman and Krilis, "Anti-phospholipid antibodies are directed against a complex antigen that includes a lipid-binding inhibitor of coagulation: beta 2-glycoprotein I (apolipoprotein H)," *Proc. Natl. Acad. Sci. USA*, 87(11):4120-4, 1990.
- Mills, Brooker and Camerini-Otero, "Sequences of human immunoglobulin switch regions: implications for recombination and transcription," *Nucl. Acids Res.*, 18:7305-7316, 1990.
- 40 Miyake, Medina, Ishihara, Kimoto, Auerbach and Kincade, "VCAM-like adhesion molecule on murine bone marrow stromal cells mediates binding of lymphocyte precursors in culture," *J. Cell. Biol.*, 114:557-565, 1991.

- Moldovan, Moldovan and Simionescu, "Binding of vascular anticoagulant alpha (annexin V) to the aortic intima of the hypercholesterolemic rabbit. An autoradiographic study," *Blood Coagul Fibrinolysis*, 5(6):921-928, 1994.
- 5 Montesano, Pepper, Mohle-Steinlein, Risau, Wagner and Orci, "Increased proteolytic activity is responsible for the aberrant morphogenetic behavior of endothelial cells expressing the middle T oncogene," *Cell*, 62:435-445, 1990.
- 10 Moore *et al.*, "Tumor angiogenesis is regulated by CXC chemokines," *J. Lab. Clin. Med.*, 132(2):97-103, 1998.
- Morrison, Johnson, Herzenberg and Oi, "Chimeric human antibody molecules: mouse antigen-binding domains with human constant region domains," *Proc. Natl. Acad. Sci. USA*, 81(21):6851-6855, 1984.
- 15 Morrison, Wims, Kobrin and Oi, "Production of novel immunoglobulin molecules by gene transfection," *Mt. Sinai J. Med.*, 53(3):175, 1986.
- 20 Morrissey, Fair and Edgington, "Monoclonal antibody analysis of purified and cell-associated tissue factor," *Thromb. Res.*, 52:247-261, 1988.
- Müller, Pomorski, Müller, Zachowski and Herrmann, "Protein-dependent translocation of aminophospholipids and asymmetric transbilayer distribution of phospholipids in the plasma membrane of ram sperm cells," *Biochemistry*, 33:9968-9974, 1994.
- 25 Munro, "Endothelial-leukocyte adhesive interactions in inflammatory diseases," *European Heart Journal*, 14:72-77, 1993.
- 30 Murphy, Joseph, Stephens and Horrocks, "Phospholipid composition of cultured human endothelial cells," *Lipids*, 27(s):150-153, 1992.
- Murray, Clauss, Thurston and Stern, "Tumour-derived factors which induce endothelial tissue factor and enhance the procoagulant response to TNF," *Int. J. Radiat. Biol.*, 60(1-2):273-277, 1991.
- 35 Nagler, Feferman, Shoshan, "Reduction in basic fibroblast growth factor mediated angiogenesis *in vivo* by linomide," *Connect Tissue Res.*, 37(1-2):61-68, 1998.
- 40 Nakamura, Shidara, Kawaguchi, Azuma, Mitsuda, Onishi, Yamaji and Wada, "Lupus anticoagulant autoantibody induces apoptosis in umbilical vein endothelial cells: involvement of annexin V," *Biochem. Biophys. Res. Comm.*, 205(2):1488-1493, 1994.
- 45 Nakamura, Ban, Yamaji, Yoneda and Wada, "Localization of the apoptosis-inducing activity of lupus anticoagulant in an annexin V-binding antibody subset," *J. Clin. Invest.*, 101(9):1951-1959, 1998.

Nawroth, Stern, Kisiel and Bach, "Cellular requirements for tissue factor generation by bovine aortic endothelial cells in culture," *Thromb. Res.*, 40:677-691, 1985.

- 5 Nawroth and Stern, "Modulation of endothelial cell hemostatic properties by tumor necrosis factor," *J. Exp. Med.*, 163:740-745, 1986.

10 Nawroth, Handley, Matsueda, DeWaal, Gerlach, Blohm and Stern, "Tumor necrosis factor/cachectin-induced intravascular fibrin formation in meth A fibrosarcomas," *J. Exp. Med.*, 168:637-647, 1988.

Obringer, Rote and Walter, "Antiphospholipid antibody binding to bilayer-coated glass microspheres," *J. Immun. Methods*, 185:81-93, 1995.

- 15 Ogawa, Shreeniwas, Brett, Clauss, Furie and Stern, "The effect of hypoxia on capillary endothelial cell function: modulation of barrier and coagulant function," *J. Haematology*, 75:517-524, 1990.

20 Ohizumi, Tsunoda, Taniguchi, Saito, Esaki, Makimoto, Wakai, Tsutsumi, Nakagawa, Utoguchi, Kaiho, Ohsugi and Mayumi, "Antibody-based therapy targeting tumor vascular endothelial cells suppresses solid tumor growth in rats," *Biochem. Biophys. Res. Comm.*, 236:493-496, 1997.

25 Oikawa *et al.*, "A highly potent antiangiogenic activity of retinoids," *Cancer Lett.*, 48(2):157-162, 1989.

O'Reilly *et al.*, "Angiostatin: a novel angiogenesis inhibitor that mediates the suppression of metastases by a Lewis lung carcinoma," *Cell*, 79:315-328, 1994.

30 O'Reilly *et al.*, "Endostatin: an endogenous inhibitor of angiogenesis and tumor growth," *Cell*, 88(2):277-285, 1997.

35 Ortel, Devore-Carter, Quinn-Allen and Kane, "Deletion analysis of recombinant human factor V. Evidence for a phosphatidylserine binding site in the second C-type domain," *J. Biol. Chem.*, 267:4189-4198, 1992.

40 Papapetropoulos, Garcia-Cardena, Dengler, Maisonpierre, Yancopoulos, Sessa, "Direct actions of angiopoietin-1 on human endothelium: evidence for network stabilization, cell survival, and interaction with other angiogenic growth factors", *Lab. Invest.*, 79(2):213-23, 1999.

Parmley and Smith, "Antibody-selectable filamentous fd phage vectors: affinity purification of target genes," *Gene*, 73(2):305-318, 1988.

- Patey, Vazeux, Canioni, Potter, Gallatin and Brousse, "Intercellular adhesion molecule-3 on endothelial cells: Expression in tumors but not in inflammatory responses," *Am. J. Pathol.*, 148:465-472, 1996.
- 5 Pepper *et al.*, "Leukemia inhibitory factor (LIF) inhibits angiogenesis *in vitro*," *J. Cell Sci.*, 108(Pt 1):73-83, 1995.
- Qamar, Gharavi, Levy and Lockshin, "Lysophosphatidylethanolamine is the antigen to which apparent antibody to phosphatidylethanolamine binds," *J. Clin. Immunol.*, 10(4):200-203, 10 1990.
- Qu, Conroy, Walker, Wooding and Lucy, "Phosphatidylserine-mediated adhesion of T-cells to endothelial cells," *J. Biochem.*, 317(Pt 2):343-346, 1996.
- 15 Quinn *et al.*, CM101, a polysaccharide antitumor agent, does not inhibit wound healing in murine models," *J. Cancer Res. Clin. Oncol.*, 121(4):253-256, 1995.
- Rao, Tait and Hoang, "Binding of annexin V to a human ovarian carcinoma cell line (OC-2008). Contrasting effects on cell surface factor VIIa/tissue factor activity and prothrombinase activity," *Thromb. Res.*, 67(5):517-531, 1992. 20
- Rauch, Tannenbaum, Tannenbaum, Ramelson, Cullis, Tilcock, Hope and Janoff, "Human hybridoma lupus anticoagulants distinguish between lamellar and hexagonal phase lipid systems," *J. Biol. Chem.*, 261(21):9672-9677, 1986. 25
- Rauch and Janoff, "Phospholipid in the hexagonal II phase is immunogenic: evidence for immunorecognition of nonbilayer lipid phases *in vivo*," *Proc. Natl. Acad. Sci., USA*, 87(11):4112-4, 1990.
- 30 Ravanat, Archipoff, Beretz, Freund, Cazenave and Freyssinet, "Use of annexin-V to demonstrate the role of phosphatidylserine exposure in the maintenance of hemostatic balance by endothelial cells," *Biochem. J.*, 282:7-13, 1992.
- RayChaudhury and D'Amore, "Endothelial cell regulation by transforming growth factor-beta," 35 *J. Cell Biochem.*, 47(3):224-229, 1991.
- Riechmann, Clark, Waldmann and Winter, "Reshaping human antibodies for therapy," *Nature*, 332(6162):323-327, 1988.
- 40 Richer and Lo, "Introduction of human DNA into mouse eggs by injection of dissected human chromosome fragments", *Science* 245, 175-177, 1989.
- Rote, Ng, Dostal-Johnson, Nicholson and Siekman, "Immunologic detection of phosphatidylserine externalization during thrombin-induced platelet activation," 45 *Clin. Immunol. Immunopathol.*, 66:193-200, 1993.

- 5 Rote, Chang, Katsuragawa, Ng, Lyden and Mori, "Expression of phosphatidylserine-dependent antigens on the surface of differentiating BeWo human choriocarcinoma cells," *Am. J. Reprod. Immun.*, 33:114-121, 1995.
- Rote, "Antiphospholipid antibodies and recurrent pregnancy loss," *Am. J. Reprod. Immun.*, 35:394-401, 1996.
- 10 Ruf and Edgington, "Structural biology of tissue factor, the initiator of thrombogenesis *in vivo*," *FASEB J.*, 8:385-390, 1994.
- Ruf, Rehemtulla and Edgington, "Phospholipid-independent and -dependent interactions required for tissue factor receptor and cofactor function," *Biol. Chem.*, 266:2158-2166, 1991.
- 15 Sambrook, Fritsch and Maniatis, *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Press, Cold Spring Harbor, NY, 1989.
- Sakamoto *et al.*, "Heparin plus cortisone acetate inhibit tumor growth by blocking endothelial cell proliferation," *Canc. J.*, 1:55-58, 1986.
- 20 Sang, "Complex role of matrix metalloproteinases in angiogenesis," *Cell Res.*, 8(3):171-177, 1998.
- Schick, Kurica and Chacko, "Location of phosphatidylethanolamine and phosphatidylserine in the human platelet plasma membrane," *J. Clin. Invest.*, 57:1221-1226, 1976.
- 25 Schick, "The organization of aminophospholipids in human platelet membranes: selective changes induced by thrombin", *J. Lab. Clin. Med.*, 91(5):802-10, 1978.
- 30 Schorer, Rick, Swaim and Moldow, "Structural features of endotoxin required for stimulation of endothelial cell tissue factor production; exposure of preformed tissue factor after oxidant-mediated endothelial cell injury," *J. Lab. Clin. Med.*, 106:38-42, 1985.
- 35 Schuurmans Stekhoven, Tijmes, Umeda, Inoue and De Pont, "Monoclonal antibody to phosphatidylserine inhibits Na⁺/K⁺-ATPase activity," *Biochimica et Biophysica Acta*, 1194:155-165, 1994.
- 40 Sheibani and Frazier, "Thrombospondin 1 expression in transformed endothelial cells restores a normal phenotype and suppresses their tumorigenesis," *Proc. Natl. Acad. Sci. USA*, 92(15):6788-6792, 1995.
- 45 Sheu *et al.*, "Inhibition of angiogenesis in vitro and in vivo: comparison of the relative activities of triflavin, an Arg-Gly-Asp-containing peptide and anti-alpha(v)beta3 integrin monoclonal antibody," *Biochim. Biophys. Acta*, 1336(3):445-454, 1997.

Shyu, Manor, Magner, Yancopoulos, Isner, "Direct intramuscular injection of plasmid DNA encoding angiopoietin-1 but not angiopoietin-2 augments revascularization in the rabbit ischemic hindlimb", *Circulation*, 98(19):2081-7, 1998.

- 5 Sideras, Mizuta, Kanamori, Suzuki, Okamoto, Kuze, Ohno, Doi, Fukuhara, Hassan, *et al.*, "Production of sterile transcripts of C gamma genes in an IgM-producing human neoplastic B cell line that switches to IgG-producing cells," *Intl. Immunol.*, 1(6):631-642, 1989.

- 10 Sipos *et al.*, "Inhibition of tumor angiogenesis," *Ann. NY Acad. Sci.*, 732:263-272, 1994.

Sluiter, Pietersma, Lamers and Koster, "Leukocyte adhesion molecules on the vascular endothelium: their role in the pathogenesis of cardiovascular disease and the mechanisms underlying their expression," *J. Cardiol. Pharmacol.*, 22:S37-S44, 1993.

- 15 Smirnov, Triplett, Comp, Esmon and Esmon, "On the role of phosphatidylethanolamine in the inhibition of activated protein C activity by antiphospholipid antibodies," *J. Clin. Invest.*, 95(1):309-16, 1995.

- 20 Soff *et al.*, "Expression of plasminogen activator inhibitor type 1 by human prostate carcinoma cells inhibits primary tumor growth, tumor-associated angiogenesis, and metastasis to lung and liver in an athymic mouse model," *J. Clin. Invest.*, 96(6):2593-2600, 1995.

- 25 Staal-van den Brekel, Thunnissen, Buurman and Wouters, "Expression of E-selectin, intercellular adhesion molecule (ICAM)-1 and vascular cell adhesion molecule (VCAM)-1 in non-small-cell lung carcinoma," *Virchows Arch.*, 428:21-27, 1996.

- 30 Staub, Harris, Khamashta, Savidge, Chahade and Hughes, "Antibody to phosphatidylethanolamine in a patient with lupus anticoagulant and thrombosis," *Ann. Rheum. Dis.*, 48(2):166-169, 1989.

- 35 Stone, Ruf, Miles, Edgington and Wright, "Recombinant soluble human tissue factor secreted by *Saccharomyces cerevisiae* and refolded from *E. coli* inclusion bodies: glycosylation of mutants, activity, and physical characterization," *Biochem. J.*, 310(2):605-614, 1995.

- Stout, Basse, Luhm, Weiss, Wiedmer and Sims, "Scott syndrome erythrocytes contain a membrane protein capable of mediating Ca^{2+} -dependent transbilayer migration of membrane phospholipids," *J. Clin. Invest.*, 99(9):2232-8, 1997.

- 40 Stratmann, Risau, Plate, "Cell type-specific expression of angiopoietin-1 and angiopoietin-2 suggests a role in glioblastoma angiogenesis", *Am. J. Pathol.*, 153(5):1333-9, 1998.

- 45 Sugi and McIntyre, "Autoantibodies to phosphatidylethanolamine (PE) recognize a kininogen-PE complex," *Blood*, 86(8):3083-3089, 1995.

- Sugi and McIntyre, "Phosphatidylethanolamine induces specific conformational changes in the kininogens recognizable by antiphosphatidylethanolamine antibodies," *Thromb. Haemost.*, 76(3):354-360, 1996a.
- 5 Sugi and McIntyre, "Autoantibodies to kininogen- phosphatidylethanolamine complexes augment thrombin-induced platelet aggregation," *Thromb. Res.*, 84(2):97-109, 1996b.
- Sugimura, Donato, Kakar and Scully, "Annexin V as a probe of the contribution of anionic phospholipids to the procoagulant activity of tumor cell surfaces," *Blood Coagul. Fibrinolysis*, 5(3):365-373, 1994.
- 10 Tada *et al.*, "Inhibition of tubular morphogenesis in human microvascular endothelial cells by co-culture with chondrocytes and involvement of transforming growth factor beta: a model for avascularity in human cartilage," *Biochim. Biophys. Acta*, 1201(2):135-142, 1994.
- 15 Takano *et al.*, "Suramin, an anticancer and angiostatic agent, inhibits endothelial cell binding of basic fibroblast growth factor, migration, proliferation, and induction of urokinase-type plasminogen activator," *Cancer Res.*, 54(10):2654-2660, 1994.
- 20 Tanaka *et al.*, "Viral vector-mediated transduction of a modified platelet factor 4 cDNA inhibits angiogenesis and tumor growth," *Nat. Med.*, 3(4):437-442, 1997.
- Tanaka, Mori, Sakamoto, Makuuchi, Sugimachi, Wands, "Biologic significance of angiopoietin - 2 expression in human hepatocellular carcinoma," *J. Clin. Invest.*, 103(3):341-5, 1999.
- 25 Thornhill, Kyan-Aung and Haskard, "IL-4 increases human endothelial cell adhesiveness for T cells but not for neutrophils," *J. Immunol.*, 144:3060-3065, 1990.
- 30 Thorpe *et al.*, "Heparin-Steroid Conjugates: New Angiogenesis Inhibitors with Antitumor Activity in Mice," *Cancer Res.*, 53:3000-3007, 1993.
- Tolsma *et al.*, "Peptides derived from two separate domains of the matrix protein thrombospondin-1 have anti-angiogenic activity," *J. Cell Biol.*, 122(2):497-511, 1993.
- 35 Toti, Satta, Fressinaud, Meyer and Freyssinet, "Scott syndrome, characterized by impaired transmembrane migration of procoagulant phosphatidylserine and hemorrhagic complications, is an inherited disorder," *Blood*, 87(4):1409-1415, 1996.
- 40 Trudell, Ardies and Anderson, "Antibodies raised against trifluoroacetyl-protein adducts bind to N-trifluoroacetyl-phosphatidylethanolamine in hexagonal phase phospholipid micelles," *J. Pharmacol. Exp. Ther.*, 257(2):657-662, 1991a.
- 45 Trudell JR, Ardies, Green and Allen, "Binding of anti-acetaldehyde IgG antibodies to hepatocytes with an acetaldehyde-phosphatidylethanolamine adduct on their surface," *Alcohol Clin. Exp. Res.*, 15(2):295-299, 1991b.

- Tryggvason, "The laminin family," *Curr. Opin. Cell Biol.*, 5(5):877-882, 1993.
- 5 Umeda, Igarashi, Nam and Inoue, "Effective production of monoclonal antibodies against phosphatidylserine: Stereo-specific recognition of phosphatidylserine by monoclonal antibody," *J. Immun.*, 143(7):2273-2279, 1989.
- 10 Utsugi, Schroit, Connor, Bucana and Fidler, "Elevated expression of phosphatidylserine in the outer membrane leaflet of human tumor cells and recognition by activated human blood monocytes," *Cancer Res.*, 51(11):3062-3066, 1991.
- 15 Valenzuela, Griffiths, Rojas, Aldrich, Jones, Zhou, McClain, Copeland, Gilbert, Jenkins, Huang, Papadopoulos, Maisonpierre, Davis, Yancopoulos, "Angiopoietins 3 and 4: diverging gene counterparts in mice and humans", *Proc. Natl. Acad. Sci., USA*, 96(5):1904-9, 1999.
- van Dijk, Warnaar, van Eendenburg, Thienpont, Braakman, Boot, Fleuren and Bolhuis, "Induction of tumor-cell lysis by bi-specific monoclonal antibodies recognizing renal-cell carcinoma and CD3 antigen," *Int. J. Cancer*, 43:344-349, 1989.
- Van Heerde, Poort, van T. Veer, Reutelingsperger and de Groot, "Binding of recombinant annexin V to endothelial cells: effect of annexin V binding on endothelial-cell-mediated thrombin formation," *J. Biochem.*, 302:305-312, 1994.
- 25 Vermes, Haanes, Steffens-Nakken and Reutelingsperger, "A novel assay for apoptosis. Flow cytometric detection of phosphatidylserine expression on early apoptotic cells using fluorescein labeled Annexin V," *J. Immunol. Methods*, 184(1):39-51, 1995.
- 30 Vitetta *et al.*, "Phase I immunotoxin trial in patients with B-cell lymphoma," *Cancer Res.*, 51:4052-4058, 1991.
- Vlachoyiannopoulos, Beigbeder, Duelanes, Youinou, Hunt, Krilis and Moutsopoulos, "Antibodies to phosphatidylethanolamine in antiphospholipid syndrome and systemic lupus erythematosus: their correlation with anticardiolipin antibodies and beta 2 glycoprotein-I plasma levels," *Autoimmunity*, 16(4):245-249, 1993.
- 35 Vogt, Ng and Rote, "A model for the antiphospholipid antibody syndrome: Monoclonal antiphosphatidylserine antibody induces intrauterine growth restriction in mice," *Am. J. Obstet. Gynecol.*, 174:700-707, 1996.
- 40 Vogt, Ng and Rote, "Antiphosphatidylserine antibody removes Annexin V and facilitates the binding prothrombin at the surface of a choriocarcinoma model of trophoblast differentiation," *Am. J. Obstet. Gynecol.*, 177:964-972, 1997.

Volpert, Lawler, Bouck, "A human fibrosarcoma inhibits systemic angiogenesis and the growth of experimental metastases via thrombospondin-1," *Proc. Natl. Acad. Sci. USA*, 95(11):6343-6348, 1998.

- 5 Vukanovic *et al.*, "Antiangiogenic effects of the quinoline-3-carboxamide linomide," *Cancer Res.*, 53(8):1833-1837, 1993.

10 Waltenberger *et al.*, "Suramin is a potent inhibitor of vascular endothelial growth factor. A contribution to the molecular basis of its antiangiogenic action," *J. Mol. Cell Cardiol.*, 28(7):1523-1529, 1996.

Wamil *et al.*, "Soluble E-selectin in cancer patients as a marker of the therapeutic efficacy of CM101, a tumor-inhibiting anti-neovascularization agent, evaluated in phase I clinical trail," *J. Cancer Res. Clin. Oncol.*, 123(3):173-179, 1997.

15

Wells, "Starving cancer into submission" *Chem. Biol.*, 5(4):R87-88, 1998.

Weiss, Young, LoBuglio, Slivka and Nimeh, "Role of Hydrogen Peroxide in Neutrophil-Mediated Destruction of Cultured Endothelial Cells," *J. Clin. Invest.*, 68:714-721, 1981.

20

White, Haandier, Smith, Hill and Lehman, In: *Principles of Biochemistry, 6th Edition*, McGraw-Hill, Inc. N.Y., Chapter 3 and pp 48-54, 1978.

25 Williamson and Schlegel, "Back and forth: the regulation and function of transbilayer phospholipid movement in eukaryotic cells," *Molec. Mem. Biol.*, 11:199-216, 1994.

Winter and Milstein, "Man-made antibodies," *Nature*, 349:293-299, 1991.

30 Wolff *et al.*, "Dexamethasone inhibits glioma-induced formation of capillary like structures *in vitro* and angiogenesis *in vivo*," *Klin. Padiatr.*, 209(4):275-277, 1997.

Yamada, Moldow, Sacks, Craddock, Boogaens and Jacob, "Deleterious Effects of Endotoxin on Cultured Endothelial Cells: An *in vitro* Model of Vascular injury," *Inflammation*, 5:115-116, 1981.

35

Yamamura *et al.*, "Effect of Matrigel and laminin peptide YIGSR on tumor growth and metastasis," *Semin. Cancer Biol.*, 4(4):259-265, 1993.

40

Yoon *et al.*, "Inhibitory effect of Korean mistletoe (*Viscum album coloratum*) extract on tumour angiogenesis and metastasis of haematogenous and non- haematogenous tumour cells in mice," *Cancer Lett*, 97(1):83-91, 1995.

45 Yoshida *et al.*, "Suppression of hepatoma growth and angiogenesis by a fumagillin derivative TNP470: possible involvement of nitric oxide synthase," *Cancer Res.*, 58(16):3751-3756, 1998.

